

MODELLING MODIFIED ATMOSPHERE PACKAGING FOR FRUITS AND VEGETABLES USING MEMBRANE SYSTEMS

Gabi Escuela¹, Thomas Hinze², Peter Dittrich¹, Stefan Schuster²

*Friedrich Schiller University Jena, ¹Bio Systems Analysis Group and ²Department of Bioinformatics, Jena, Germany
{gabi.escuola, thomas.hinze, Peter.Dittrich, Stefan.Schu}@uni-jena.de*

Mario Moreno-Álvarez

*U.N.E. Simón Rodríguez, Laboratorio de Biomoléculas, Canoabo, Venezuela
morenoalvarez@cantv.net*

Keywords: P systems, Modified Atmosphere Packaging, fruit modelling, membrane systems, respiration, temperature.

Abstract: As living materials, post-harvested fruits and vegetables continue their metabolic activity, exhibiting progressive biochemical changes. Optimisation of environmental conditions during storage of these fresh commodities is required in order to increase their shelf life. In this work we use P systems to abstract molecular interactions that occur between plant organ, film and surrounding atmosphere factors involved in fresh fruit and vegetable package designs. The proposed model constitutes a general framework to simulate the dynamical behaviour of these systems, specially due to gas exchanges and temperature fluctuations. Moreover, the model can be extended introducing other variables and processes that affect quality of such produces. This can be considered, to the best of our knowledge, the first contribution of Membrane Computing in Food Engineering.

1 INTRODUCTION

Membrane systems (Păun and Rozenberg, 2002), also called P systems, had emerged to assist in the modelling of systems of concurrent reactions taking place in compartments, so as occur in biological systems. In this paper we use P systems as membrane structures delimiting compartments that contain multisets of objects representing molecules. Compartments configuration changes over time (evolve) according to given rules that represent biochemical reactions and diffusions. In contrast to ODE-based approaches, each single molecule within the entire system is represented explicitly as individual entity. Capturing aspects of structural dynamics (changes in the membrane structure as well as in the composition of complex molecules) is seen as an advantageous feature of P systems. Inclusion of reaction kinetics into this formalism can be done by discretised kinetic laws (Hinze et al., 2006). We applied this mathematical formalism to a real known problem in fruits and vegetables post-harvest processing.

Fresh fruits and vegetables are living materials that continue to respire after harvesting exhibiting progressive biochemical changes. Food Engineering methods to preserve freshness of post-harvest

produces include low temperature storage and special packaging technologies, mainly Modified Atmosphere Packaging (MAP). MAP of fresh fruits and vegetables refers to the technique of enveloping the produce in a sealed container of polymeric film in order to modify the O_2 and CO_2 concentrations inside the package, reducing metabolic activity and increasing shelf life (Paul and Clarke, 2002).

Designing MAP systems is a complex task that involves considerations about many interrelated environmental (as temperature and atmosphere composition), biological and package technology factors. Basic biological processes are respiration, transpiration, ethylene production and compositional changes due to metabolism. The variability of responses to internal and external signals depends on the characteristic of each plant organ type, developmental stage and physiological condition. In addition, much of the behaviour of a MAP system at cellular level are not fully understood. As examples we can refer to the little knowledge about the effect of CO_2 on the activity of respiratory enzymes (Ho et al., 2008). Moreover, the contribution of the biochemical changes that alters physical properties of cell walls and tissues modifying the texture of the produce is not known in detail (Gross et al., 2004). On the other hand, the mechanism of ethylene

signal transduction that coordinates fruit ripening processes, is another aspect subject to study (Alexander and Grierson, 2002).

The difficulty to test different combination of gases and temperatures and the complexity of experimental setup for MAP systems had lead to the development of various mathematical models (Exama et al., 1993; Tijssens et al., 2001; Paul and Clarke, 2002; Ho et al., 2008) and software (Mahajan et al., 2007). In the literature, many respiration models are empirical fits of experimental data, based on one particular type and variety of fruit or vegetable, and most of them are based on the principles of enzyme kinetics and are represented using ODEs (for reviews see (Fonseca et al., 2002; Rodriguez-Aguilera and Oliveira, 2009)). However, there exists some lack on studies about the dynamical behaviour of these systems in terms of changes in environmental conditions, so as produce composition and physiology due to developmental processes (Fonseca et al., 2002). It is worth mentioning that post-harvested fruits and vegetables, unlike other living materials, can be considered as less robust systems, as their responses on environmental fluctuations depends mostly on their actual configuration of biochemical components. In this context, some authors (Génard et al., 2007; Mahajan et al., 2007) have considered the potential benefits of a systematic analysis or process-based modelling approach for fruits and vegetables. Considering that understanding the reaction network underlying MAP systems can give food experts more knowledge about emergent properties of packaged fruits and vegetables, we propose a framework based on membrane systems that abstracts basic biochemical reactions that occur in MAP systems. In the future, the proposed model can serve as a predictive tool to simulate changes in fresh produce on the molecular level, due to changes in environmental conditions.

This paper is organized as follows: in Section 2 we present a P system framework for MAP, including the description of components, reaction kinetics and evolution of the system. Section 3 shows an application of the framework considering a package under modified atmosphere containing two produces. Finally, in Section 4 we point out some benefits of using our framework and future extensions of it.

2 A P SYSTEM-BASED MAP

We abstract a fruit or vegetable as a graph of cells or modules, like tissue P systems (Martín-Vide et al., 2003). Each cell represents a compartment that contains species, and at a specific time, the contents of the

compartment determine the cell configuration. This serves as a mechanism to differentiate one cell from other, given the possibility to creating diverse tissue types, as occurs for example in fruits epicarp, mesocarp and endocarp tissues (Génard et al., 2007). Additionally, as gas consumption-production occur inside the cells, at the mitochondria level, and is stated that gas diffusion between cells depends on the geometry of the produce (Ho et al., 2008), differences in gas content in cells that conform a determinate region can adequately be represented. This is also in accordance to the idea that the ripening process usually starts in one region of a fruit and spreads to neighbouring regions, due to ethylene diffusion starting from promoter cells (Alexander and Grierson, 2002). Produces into the package are represented as a population of membranes, giving the advantage that the model can deal with distinct fruits and vegetables within the same film, or the same produce in distinct developmental stages, varieties and/or presentations. Figure 1 shows as example, the schematic representation for such a system. In the next section we present the formal specification of our model.




Figure 1: A schematic representation for the MAP system model. In this case, two produces share a package: $plant_1$ is formed by three connected cells, and $plant_2$ is formed by a single cell. Arrows represent paths for molecules (spheres) diffusions

Multiset Prerequisites

Let A be an arbitrary set and \mathbb{N} the set of natural numbers including zero. A multiset over A is a mapping $F : A \rightarrow \mathbb{N} \cup \{\infty\}$. $F(a)$, also denoted as $[a]_F$, specifies the multiplicity of $a \in A$ in F . Multisets can be written as an element-wise enumeration of the form $\{(a_1, F(a_1)), (a_2, F(a_2)), \dots\}$ since $\forall (a, b_1), (a, b_2) \in F : b_1 = b_2$. The support $\text{supp}(F) \subseteq A$ of F is defined by $\text{supp}(F) = \{a \in A \mid F(a) > 0\}$. A multiset F over A is said to be empty iff $\forall a \in A : F(a) = 0$. The cardinality $|F|$ of F over A is $|F| = \sum_{a \in A} F(a)$. Let F_1 and F_2 be multisets over A . F_1 is a subset of F_2 , denoted as $F_1 \subseteq F_2$, iff $\forall a \in A : (F_1(a) \leq F_2(a))$. Multisets F_1 and F_2

are equal iff $F_1 \subseteq F_2 \wedge F_2 \subseteq F_1$. The intersection $F_1 \cap F_2 = \{(a, F(a)) \mid a \in A \wedge F(a) = \min(F_1(a), F_2(a))\}$, the multiset sum $F_1 \uplus F_2 = \{(a, F(a)) \mid a \in A \wedge F(a) = F_1(a) + F_2(a)\}$, and the multiset difference $F_1 \ominus F_2 = \{(a, F(a)) \mid a \in A \wedge F(a) = \max(F_1(a) - F_2(a), 0)\}$ form multiset operations. Multiplication of a multiset $F = \{(a, F(a)) \mid a \in A\}$ with a scalar c , denoted $c \cdot F$, is defined by $\{(a, c \cdot F(a)) \mid a \in A\}$.

P system components

Let $\mathbb{N}_+ = \mathbb{N} \setminus \{0\}$ be the set of natural numbers without zero, and $m, n \in \mathbb{N}_+$. We define a P system for a MAP system as a construct:

$$\Pi_{\text{MAP}} = (\mu, S, \text{plant}_1, \dots, \text{plant}_m, G, L_0, D_1, \dots, D_d, f_1, \dots, f_d, \Delta\tau)$$

where:

- $\mu = ([\![\![\text{cell}_{1,1} \dots \text{cell}_{1,m_1}] \text{plant}_1 \dots [\![\text{cell}_{m,1} \dots \text{cell}_{m,m_m}] \text{plant}_m] \text{package}] \text{package})$ is the spatial system structure composed of three inner levels: package, plants, and cells,
- S is a set of chemical species,
- $\text{plant}_1, \dots, \text{plant}_m$ represent the produces into the package,
- G is a set of global parameters,
- $L_0 : S \rightarrow \mathbb{N}$ is a multiset of axioms representing the initial molecular configuration,
- D_v is a diffusion (communication) rule among package and external environment ($v = 1, \dots, d$),
- $f_v : (S \rightarrow \mathbb{N}) \rightarrow \mathbb{N}$ is a kinetic function attached to diffusion rule D_v ,
- $\Delta\tau \in \mathbb{R}_+$ is the time discretisation interval.

A diffusion rule D_v can be of the form $[s] \rightarrow [s]$ for molecules $s \in S$ leaving the package and released to the external environment, and $[s] \rightarrow [s]$ for molecules entering the package, respectively.

Furthermore, each plant_i is defined as a tuple:

$$\text{plant}_i = (N_i, E_i, G_i, D_{i,1}, \dots, D_{i,d_i}, f_{i,1}, \dots, f_{i,d_i})$$

where:

- $N_i = \{\text{cell}_{i,1}, \dots, \text{cell}_{i,m_i}\}$ defines a set of cells within plant i ,
- $E_i \subseteq N_i \times N_i$ specifies a set of directed edges (diffusion channels between cells),
- G_i is a set of plant (organ) specific parameters,
- $D_{i,\kappa}$ represents a diffusion rule inside plant i and between plant i and package ($\kappa = 1, \dots, d_i$),
- $f_{i,\kappa}$ is a kinetic function attached to diffusion rule $D_{i,\kappa}$.

Here, a diffusion rule can be of the form $[s]_{\text{cell}_{p,q}} \rightarrow [s]_{\text{cell}_{p,q}}$ for molecules $s \in S$ leaving $\text{cell}_{p,q}$ and spread out into the package. A rule of the form $[s]_{\text{cell}_{p,q}} \rightarrow [s]_{\text{cell}_{p,q}}$ describes molecules entering $\text{cell}_{p,q}$ from the package. Finally, a rule of the form $[s]_{\text{cell}_{p,q}} \rightarrow [s]_{\text{cell}_{x,y}}$ formulates the directed transport of molecule s along the edge $(\text{cell}_{p,q}, \text{cell}_{x,y}) \in E_i$.

Each $\text{cell}_{i,j}$ is defined as a tuple

$$\text{cell}_{i,j} = (L_{i,j,0}, R_{i,j,1}, \dots, R_{i,j,r_{i,j}}, f_{i,j,1}, \dots, f_{i,j,r_{i,j}})$$

where:

- $L_{i,j,0} : S \rightarrow \mathbb{N}$ is a multiset of axioms representing its initial molecular configuration,
- $R_{i,j,k} = (A_{i,j,k}, B_{i,j,k})$ with $A_{i,j,k} : S \rightarrow \mathbb{N}$ (multiset of reactants) and $B_{i,j,k} : S \rightarrow \mathbb{N}$ (multiset of products) specifies a reaction rule including its stoichiometric factors,
- $f_{i,j,k} : (S \rightarrow \mathbb{N}) \rightarrow \mathbb{N}$ is a function corresponding to kinetics of reaction $R_{i,j,k}$.

System evolution

A P system of the form Π_{MAP} evolves by successive progression of its configuration at discrete points in time $t \in \mathbb{N}$ for what we assume a global clock. Two consecutive dates t and $t + 1$ specify a time span $\Delta\tau$. A system step at time t consists of three modification stages carried out from outer to inner spatial components of the system. Firstly, the diffusion between package and its environment is considered. To this end, the rules D_1 up to D_d are employed. Afterwards, the diffusion between package and cells as well as the intracellular diffusion is utilised by employing the rules $D_{i,\kappa}$ for each plant $i = 1, \dots, m$. The last modification stage concerns application of the reaction rules specified in each cell. To cope with conflicts that can occur if the available amount of substrate cannot satisfy all matching diffusion and reaction rules, we prioritise all rules by their index: $D_1 > D_2 > \dots > D_d$. Moreover, for each plant i : $D_{i,1} > D_{i,2} > \dots > D_{i,d_i}$ and for each cell i, j : $R_{i,j,1} > R_{i,j,2} > \dots > R_{i,j,r_{i,j}}$. Thus, we keep determinism of the system evolution and enable mass conservation. An alternative method for coping with conflicts is randomisation in selection and sequentialisation of diffusion and reaction rules.

The application of an arbitrary rule is organised into two consecutive steps. The first step identifies all molecules from the rule's left hand side acting as sources for diffusion or reactants. These molecules are removed from the current configurations. Corresponding molecules from the right hand side (destinations in case of diffusion and products in case of reactions) are then added. We formulate discretised reaction-diffusion kinetics by specification of scalar functions $f : M \rightarrow \mathbb{N}$ based on a multiset $M : S \rightarrow \mathbb{N}$. Each function f converts the current configuration (L_t or $L_{i,j,t}$), a multiset of objects, into the number of turns for application of the corresponding diffusion or reaction rule. Here, kinetic laws $\hat{f}(s)$ for each species $s \in S$ employ the multiplicity of its occurrences to formulate the corresponding reaction rate. For updating the entire system configuration, we define an iteration scheme as shown in Figure 2.

Stage 1 (diffusion between package and external environment):

$\forall \alpha = 1, \dots, d$

| diffusion rule | conditions | action |
|--|--|---|
| $D_{\alpha} = [\sigma] \rightarrow \llbracket \sigma \rrbracket$ | $(\sigma \in S) \wedge (\{(\sigma, f_{\alpha})\} \subseteq L_t)$ | $L_t := L_t \ominus \{(\sigma, f_{\alpha})\}$ |
| $D_{\alpha} = \llbracket \sigma \rrbracket \rightarrow [\sigma]$ | $(\sigma \in S)$ | $L_t := L_t \uplus \{(\sigma, f_{\alpha})\}$ |

with $f_{\alpha}(L_t) = \lfloor k_{\alpha}(G) \cdot \Delta\tau \cdot \hat{f}(|L_t \cap \{(\sigma, \infty)\}|) \rfloor$

Stage 2 (diffusion between plant cells and package):

$\forall i = 1, \dots, m$

$\forall \alpha = 1, \dots, d_i$

| diffusion rule | conditions | action |
|--|---|---|
| $D_{i,\alpha} = [\sigma]_{cell_{i,j}} \rightarrow \llbracket \sigma \rrbracket_{cell_{i,j}}$ | $(\sigma \in S) \wedge (cell_{i,j} \in N_i) \wedge (\{(\sigma, f_{i,\alpha})\} \subseteq L_{i,j,t})$ | $L_{i,j,t} := L_{i,j,t} \ominus \{(\sigma, f_{i,\alpha})\}$ $L_t := L_t \uplus \{(\sigma, f_{i,\alpha})\}$ |
| $D_{i,\alpha} = \llbracket \sigma \rrbracket_{cell_{i,j}} \rightarrow [\sigma]_{cell_{i,j}}$ | $(\sigma \in S) \wedge (cell_{i,j} \in N_i) \wedge (\{(\sigma, f_{i,\alpha})\} \subseteq L_t)$ | $L_t := L_t \ominus \{(\sigma, f_{i,\alpha})\}$ $L_{i,j,t} := L_{i,j,t} \uplus \{(\sigma, f_{i,\alpha})\}$ |
| $D_{i,\alpha} = [\sigma]_{cell_{i,j}} \rightarrow [\sigma]_{cell_{i,k}}$ | $(\sigma \in S) \wedge (k \neq j) \wedge (cell_{i,j} \in N_i) \wedge (cell_{i,k} \in N_i)$ $\wedge ((cell_{i,j}, cell_{i,k}) \in E_i) \wedge (\{(\sigma, f_{i,\alpha})\} \subseteq L_{i,j,t})$ | $L_{i,j,t} := L_{i,j,t} \ominus \{(\sigma, f_{i,\alpha})\}$ $L_{i,k,t} := L_{i,k,t} \uplus \{(\sigma, f_{i,\alpha})\}$ |

with $f_{i,\alpha}(L_t) = \lfloor k_{i,\alpha}(G, G_i) \cdot \Delta\tau \cdot \hat{f}(|L_t \cap \{(\sigma, \infty)\}|) \rfloor$

Stage 3 (reactions occurring within each cell):

$\forall i = 1, \dots, m$

$\forall j = 1, \dots, n_i$

$\forall \alpha = 1, \dots, r_{i,j}$

| reaction rule | conditions | action |
|---|---|--|
| $R_{i,j,\alpha} = (A_{i,j,\alpha}, B_{i,j,\alpha})$ | $f_{i,j,\alpha} \cdot A_{i,j,\alpha} \subseteq L_{i,j,t}$ | $L_{i,j,t} := L_{i,j,t} \ominus f_{i,j,\alpha} \cdot A_{i,j,\alpha}$ $\uplus f_{i,j,\alpha} \cdot B_{i,j,\alpha}$ |

with $f_{i,j,\alpha}(L_{i,j,t}) = \left\lfloor k_{i,j,\alpha}(G, G_i) \cdot \Delta\tau \prod_{\forall c \in \text{supp}(A_{i,j,\alpha}) : (R_{i,j,\alpha} = (A_{i,j,\alpha}, B_{i,j,\alpha}))} \hat{f}(|L_{i,j,t} \cap \{(c, \infty)\}|)^{|A_{i,j,\alpha} \cap \{(c, \infty)\}|} \right\rfloor$

Increment time t :

$L_{t+1} := L_t$

$\forall i = 1, \dots, m$

$\forall j = 1, \dots, n_i$


$L_{i,j,t+1} := L_{i,j,t}$

Figure 2: Iteration scheme for the temporal evolution of Π_{MAP} system

3 SIMULATION

As a first application, we introduced as rules into the model only the basic processes involved in a MAP design: respiration and fermentation, so as gas diffusion between membranes. Respiration rate can be expressed in terms of O_2 consumed or CO_2 produced. The respiratory quotient (RQ), the ratio of CO_2 produced to O_2 consumed, ranges from about 0.7 to 1.4 depending on the substrate and its metabolic state (if the substrate is a lipid, $RQ < 1$, and $RQ > 1$ for organic acids)(Fonseca et al., 2002). When carbohydrates are aerobically respired, the RQ is near 1, and the reaction is represented by Eq. (1). The influence of gas composition on respiration rates of produce has been widely represented by Michaelis-Menten-type

equation (Fonseca et al., 2002). In this context, respiration rate is considered as a function of concentration in terms of enzymatic reaction, with O_2 in the place of substrate and the product CO_2 acting as inhibitor.



Transpiration occurs due to the fact that fruits and vegetables internal atmosphere is saturated with water vapour, while external atmosphere contains lesser. Therefore, water loss rate depends on the external and internal water vapour pressure gradient.

Temperature dependence over respiratory rate and over film permeability was represented using Arrhenius equation (Eq. 2).

$$k = F \times e^{-E_a/R \times T} \quad (2)$$

where E_a is the activation energy, expressed in joule per mol, defined as the energy that must be overcome for a chemical reaction to occur; R is the gas constant ($\approx 8.314 \text{ J} \cdot \text{K}^{-1} \text{ mol}^{-1}$), T the absolute temperature, F is the pre-exponential factor that represents the total number of molecular collisions per second; and k corresponds to the number of collisions per second that result in a reaction. This can be related to the probabilistic approach to P systems introduced by (Ardelean and Cavaliere, 2003) in order to obtain more biological-like models. In this context, the Arrhenius exponential term can be viewed as the probability per time unit that the reaction takes place.

In order to apply our model, we simulate the dynamical behaviour of an instance of a Π_{MAP} with two hypothetical fruits as it is shown in Fig. 1, using continuous film and passive MAP as package techniques. Rules that use symbol \rightleftharpoons between reactants and products must be interpreted as reversible reactions. Into the formalism described in Fig. 2, a rule of the form $D_\alpha = [\sigma] \rightleftharpoons [\sigma]$, for example, consists in the following two rules, in order of application: $[\sigma] \rightarrow [\sigma]$ and $[\sigma] \rightarrow [\sigma]$.

$$\begin{aligned}
S &= \{CO_2, Ethanol, Glucose, H_2O, O_2\} \\
G &= \{M, T, R, A, E, CO_2ex, H_2Oex, O_2ex, EaCO_2, EaO_2, pCO_2, pO_2, pH_2O\} \\
N_1 &= \{cell_{1,1}, cell_{1,2}, cell_{1,3}\} \\
N_2 &= \{cell_{2,1}\} \\
G_i &: \{ErO_2, rO_2, ErCO_2f, rCO_2f\} \forall i \in \{1, 2\} \\
D_1 &: \llbracket package \rrbracket O_2 \rightleftharpoons \llbracket package \rrbracket O_2 \\
f_1(L_t) &= \left[\frac{A}{E} \cdot (O_2ex - L_t(O_2)) \cdot pO_2 \cdot e^{-\frac{EaO_2}{R \cdot T}} \right] \\
D_2 &: [CO_2]_{package} \rightleftharpoons \llbracket package \rrbracket CO_2 \\
f_2(L_t) &= \left[\frac{A}{E} \cdot (L_t(CO_2) - CO_2ex) \cdot pCO_2 \cdot e^{-\frac{EaCO_2}{R \cdot T}} \right] \\
D_3 &: [H_2O]_{package} \rightleftharpoons \llbracket package \rrbracket H_2O \\
f_3(L_t) &= \left[\frac{A}{E} \cdot (L_t(H_2O) - H_2Oex) \cdot pH_2O \right] \\
D_{1,1} &: \llbracket cell_{1,1} \rrbracket O_2 \rightleftharpoons [O_2]_{cell_{1,1}} \quad f_{1,1}(L_t) = k_{1,1} \cdot L_t(O_2) \\
D_{1,2} &: [CO_2]_{cell_{1,1}} \rightleftharpoons \llbracket cell_{1,1} \rrbracket CO_2 \quad f_{1,2}(L_t) = k_{1,2} \cdot L_t(CO_2) \\
D_{1,3} &: [H_2O]_{cell_{1,1}} \rightleftharpoons \llbracket cell_{1,1} \rrbracket H_2O \quad f_{1,3}(L_t) = k_{1,3} \cdot L_t(H_2O) \\
D_{1,4} &: [O_2]_{cell_{1,1}} \rightleftharpoons [O_2]_{cell_{1,2}} \quad f_{1,4}(L_t) = k_{1,4} \cdot L_t(O_2) \\
D_{1,5} &: [CO_2]_{cell_{1,2}} \rightleftharpoons [CO_2]_{cell_{1,1}} \quad f_{1,5}(L_t) = k_{1,5} \cdot L_t(CO_2) \\
D_{1,6} &: [H_2O]_{cell_{1,2}} \rightleftharpoons [H_2O]_{cell_{1,1}} \quad f_{1,6}(L_t) = k_{1,6} \cdot L_t(H_2O) \\
D_{1,7} &: [O_2]_{cell_{1,2}} \rightleftharpoons [O_2]_{cell_{1,3}} \quad f_{1,7}(L_t) = k_{1,7} \cdot L_t(O_2) \\
D_{1,8} &: [CO_2]_{cell_{1,3}} \rightleftharpoons [CO_2]_{cell_{1,2}} \quad f_{1,8}(L_t) = k_{1,8} \cdot L_t(CO_2) \\
D_{1,9} &: [H_2O]_{cell_{1,3}} \rightleftharpoons [H_2O]_{cell_{1,2}} \quad f_{1,9}(L_t) = k_{1,9} \cdot L_t(H_2O) \\
D_{2,1} &: \llbracket cell_{2,1} \rrbracket O_2 \rightleftharpoons [O_2]_{cell_{2,1}} \quad f_{2,1}(L_t) = k_{2,1} \cdot L_t(O_2) \\
D_{2,2} &: [CO_2]_{cell_{2,1}} \rightleftharpoons \llbracket cell_{2,1} \rrbracket CO_2 \quad f_{2,2}(L_t) = k_{2,2} \cdot L_t(CO_2) \\
D_{2,3} &: [H_2O]_{cell_{2,1}} \rightleftharpoons \llbracket cell_{2,1} \rrbracket H_2O \quad f_{2,3}(L_t) = k_{2,3} \cdot L_t(H_2O) \\
R_{i,j,1} &: Glucose + 6 O_2 \rightarrow 6 CO_2 + 6 H_2O \quad \forall i \in \{1, 2\} \wedge j \in \{1, 2, 3\} \\
f_{i,j,1}(L_{i,j,t}) &= \left[\frac{L_{i,j,t}(Glucose)}{\Theta_{i,j,1,1} + L_{i,j,t}(Glucose)} \cdot \frac{L_{i,j,t}(O_2)^6}{\Theta_{i,j,1,2} + L_{i,j,t}(O_2)^6} \cdot \frac{M_i}{3} \cdot rO_2 \cdot e^{-\frac{ErO_2}{R \cdot T}} \right] \\
R_{i,j,2} &: Glucose \rightarrow 2 Ethanol + 2 CO_2 \quad \forall i \in \{1, 2\} \wedge j \in \{1, 2, 3\} \\
f_{i,j,2}(L_{i,j,t}) &= \left[\frac{L_{i,j,t}(Glucose)}{\Theta_{i,j,2,1} + L_{i,j,t}(Glucose)} \cdot M_i \cdot rCO_2f \cdot e^{-\frac{ErCO_2f}{R \cdot T}} \right]
\end{aligned}$$

Temperature is represented by T and expressed in Kelvin (K). O_2 , CO_2 and H_2O abundances in the outside are represented by O_2ex , CO_2ex and H_2Oex respectively. A and E symbolise the surface area in cm^2 and the thickness of the packaging film in mil ($1mil = 0.00254cm$). pO_2 and pCO_2 represent the reference film permeability in $mL \cdot mil \cdot cm^2 \cdot hr^{-1} \cdot atm^{-1}$ for O_2 , CO_2 and H_2O , respectively. EaO_2 and $EaCO_2$ symbolise the permeability activation energy expressed in $J \cdot mol^{-1}$ for O_2 and CO_2 , respectively. M_i symbolises mass of the produce i in kg . For simplicity, we assume that each cell in a produce has the same mass. rO_2 and rCO_2f corresponds to the preexponential factor for produce respiration and fermentation in $mL \cdot kg^{-1} \cdot hr^{-1}$. ErO_2 and $ErCO_2f$ represent the respiration and fermentation activation energy for the produce expressed in $J \cdot mol^{-1}$. Most of the values for these symbols were taken from the literature (Exama et al., 1993). Symbols O_2 , CO_2 , H_2O , $Ethanol$ and $Glucose$ represent amounts of species O_2 , CO_2 , H_2O , $Ethanol$ and $Glucose$. Initial values for these symbols in each compartment and the rest of the parameters were assigned empirically.

Figure 3 shows the corresponding courses of $plant_1$ internal gas composition, resulting from following parameter setting for the discrete iteration scheme: $A = 100$, $E = 1$, $M_i = 0.1$, $pO_2i = 1620000$, $EaO_2i = 43100$, $pCO_2i = 238000$, $EaCO_2i = 34300$, $rO_2i = rCO_2fi = 3 \times 10^{14}$, $ErO_2i = ErCO_2f = 70700$, $pH_2Oi = 1$, $\Theta_{i,j,1,k} = 1$, for $i \in \{1, 2\}$ and $j \in \{1, 2, 3\}$ and $k \in \{1, 2\}$; $k_{1,j} = 0.2$ for $j \in \{1, \dots, 9\}$, $k_{2,j} = 0.2$ for $j \in \{1, \dots, 3\}$. A fixed value $T = 277.15$ was considered for a constant temperature scenario, and transient values for $273.15 \leq T \leq 293.15$ were obtained through a sigmoid function to represent changes in temperature over time in another scenario. Simulations have been performed using Copasi (Hoops et al., 2006). Differences in internal gas composition of $plant_1$ have been observed during time due to the interplay between cellular respiration and fermentation processes and intercellular diffusion. Those differences could determine the form of maturation of the produce, in this case, from the center to the skin. An equilibrium is reached in the package gas composition, while respiration rates of the produces diminished.

4 CONCLUSIONS

Using a membrane based model for MAP, we presented a framework that is able to abstract packaging for different fruit and vegetable types, varieties or developmental stages. Respiration of the produce




Figure 3: Dynamical behaviour for gas composition for $plant_1$ in constant and varying temperature scenarios

is considered as the basic process when modelling MAP, and predictions about the dynamical behaviour of such systems can be improved taking into account environmental, biological and technical factors. Our approach allows extensions including other low level processes, such as ethylene signaling pathway, cell/tissue rupture due to produce cutting and transport of other molecules, that can be easily modeled using P systems. When the formalism showed in Figure 2 is hidden in a software, the specification is intuitive and accessible for an expert focussing on MAP modelling. Finally, the quality of the packaged produce (taste, texture, colour and appearance) is based on some subjective consumer evaluation. These traits are based on specific product properties, such as sugar content, volatile production and cell wall structure (Tijksens et al., 2001), and therefore can be introduced into the model through reactions, as a mechanism to obtain more knowledge about the impact of packaging conditions over product quality.

ACKNOWLEDGEMENTS

GE is supported by Universidad Simón Bolívar (Caracas, Venezuela) and Deutscher Akademischer Austausch Dienst (DAAD) Grant A/08/94489.

REFERENCES

- Alexander, L. and Grierson, D. (2002). Ethylene biosynthesis and action in tomato: a model for climacteric fruit ripening. *J. Exp. Bot.*, 53(377):2039–2055.
- Ardelean, I. and Cavaliere, M. (2003). Modelling biological processes by using a probabilistic P system software. *Natural Computing*, 2(2):173–197.
- Exama, A., Arul, J., Lencki, R., Lee, L., and Toupin, C. (1993). Suitability of plastic films for modified atmosphere packaging of fruits and vegetables. *J Food Sci*, 58(6):1365–1370.
- Fonseca, S., Oliveira, F., and Brecht, J. (2002). Modelling respiration rate of fresh fruits and vegetables for modified atmosphere packages: a review. *J Food Eng*, 52(2):99–119.
- Génard, M., Bertin, N., Borel, C., Bussières, P., Gautier, H., Habib, R., Lechaudel, M., Lecomte, A., Lescourret, F., Lobit, P., and Quilot, B. (2007). Towards a virtual fruit focusing on quality: modelling features and potential uses. *J Exp Bot*, 58(5):917–928.
- Gross, K., Wang, C., and Saltveit, M. (2004). The commercial storage of fruits, vegetables, and florist and nursery stocks. *Agriculture Handbook*, 66.
- Hinze, T., Lenser, T., and Dittrich, P. (2006). A protein substructure based P system for description and analysis of cell signalling networks. *Lect Notes Comput Sci*, 4361:409–423.
- Ho, Q., Verboven, P., Verlinden, B., Lammertyn, J., Vandewalle, S., and Nicolai, B. (2008). A continuum model for metabolic gas exchange in pear fruit. *PLoS Computational Biology*, 4(3):e1000023.
- Hoops, S., Sahle, S., Gauges, R., Lee, C., Pahle, J., Simus, N., Singhal, M., Xu, L., Mendes, P., and Kummer, U. (2006). Copasi—a complex pathway simulator. *Bioinformatics*, 22(24):3067–3074.
- Mahajan, P., Oliveira, F., J., M., and Frias, J. (2007). Development of user-friendly software for design of modified atmosphere packaging for fresh and fresh-cut produce. *Innovative Food Science and Emerging Technologies*, 8(1):84–92.
- Martín-Vide, C., Păun, G., Pazos, J., and Rodríguez-Patón, A. (2003). Tissue P systems. *Theor Comput Sci*, 296(2):295–326.
- Paul, D. and Clarke, R. (2002). Modelling of modified atmosphere packaging based on designs with a membrane and perforations. *J Membr Sci*, 208(1–2):269–283.
- Păun, G. and Rozenberg, G. (2002). A guide to membrane computing. *Theor Comput Sci*, 287(73–100):73–100.
- Rodríguez-Aguilera, R. and Oliveira, J. (2009). Review of design engineering methods and applications of active and modified atmosphere packaging systems. *Food Engineering Reviews*, 1(1):66–83.
- Tijksens, L., Hertog, M., and Nicolai, B. (2001). *Food Process Modelling*. Woodhead Pub.