

COMPUTATION OF QUASI-LOCAL EFFECTIVE DIFFUSION TENSORS AND CONNECTIONS TO THE MATHEMATICAL THEORY OF HOMOGENIZATION*

D. GALLISTL[†] AND D. PETERSEIM[‡]

Abstract. This paper aims at bridging existing theories in numerical and analytical homogenization. For this purpose the multiscale method of Målqvist and Peterseim [*Math. Comp.*, 83 (2014), pp. 2583–2603], which is based on orthogonal subspace decomposition, is reinterpreted by means of a discrete integral operator acting on standard finite element spaces. The exponential decay of the involved integral kernel motivates the use of a diagonal approximation and, hence, a localized piecewise constant coefficient. In a periodic setting, the computed localized coefficient is proved to coincide with the classical homogenization limit. An a priori error analysis shows that the local numerical model is appropriate beyond the periodic setting when the localized coefficient satisfies a certain homogenization criterion, which can be verified a posteriori. The results are illustrated in numerical experiments.

Key words. numerical homogenization, multiscale method, upscaling, a priori error estimates, a posteriori error estimates

AMS subject classifications. 65N12, 65N15, 65N30, 73B27, 74Q05

DOI. 10.1137/16M1088533

1. Introduction. Consider the prototypical elliptic model problem

$$-\operatorname{div} A_\varepsilon \nabla u = f,$$

where the diffusion coefficient A_ε encodes microscopic features on some characteristic length scale ε . Homogenization is a tool of mathematical modeling to identify reduced descriptions of the macroscopic response of such multiscale models in the limit as ε tends to zero. It turns out that suitable limits represented by the so-called effective or homogenized coefficient exist in fairly general settings in the framework of G -, H -, or two-scale convergence [37, 10, 31, 32, 3]. In general, the effective coefficient is not explicitly given but is rather the result of an implicit representation based on cell problems. This representation usually requires structural assumptions on the sequence of coefficients A_ε such as local periodicity and scale separation [6]. Under such assumptions, efficient numerical methods for the approximate evaluation of the homogenized model are available, e.g., the heterogeneous multiscale method (HMM) [11, 1] or the two-scale finite element method [29].

In contrast to this idealized setting of analytical homogenization, in practice one is often concerned with one coefficient A with heterogeneities on multiple nonseparable scales and a corresponding sequence of scalable models can hardly be identified or may not be available at all. That is why we are interested in the computation of effective

*Received by the editors August 8, 2016; accepted for publication (in revised form) May 30, 2017; published electronically November 2, 2017.

<http://www.siam.org/journals/mms/15-4/M108853.html>

Funding: The first author was supported by Deutsche Forschungsgemeinschaft (DFG) through CRC 1173. The second author acknowledges support by DFG in Priority Program 1748, “Reliable Simulation Techniques in Solid Mechanics” (PE2143/2-1).

[†]Institut für Angewandte und Numerische Mathematik, Karlsruher Institut für Technologie, 76049 Karlsruhe, Germany (gallistl@kit.edu).

[‡]Institut für Mathematik, Universität Augsburg, 86135 Augsburg, Germany (daniel.peterseim@math.uni-augsburg.de).

representations of very rough coefficients beyond structural assumptions such as scale separation and local periodicity. In recent years, many numerical attempts have been developed that conceptually do not rely on analytical homogenization results for rough cases. Among them are the multiscale finite element method [22, 13], metric-based upscaling [34], hierarchical matrix compression [17, 19], the flux-norm approach [8], generalized finite elements based on spectral cell problems [5, 12], the AL basis [16, 38], rough polyharmonic splines [35], iterative numerical homogenization [26], and gamblets [33].

Another construction based on concepts of orthogonal subspace decomposition and the solution of localized microscopic cell problems was given in [28] and later optimized in [21, 20, 14, 36]. The method is referred to as the localized orthogonal decomposition (LOD) method. The approach is inspired by ideas of the variational multiscale method [23, 24, 27]. As most of the methods above, the LOD constructs a basis representation of some finite-dimensional operator-dependent subspace with superior approximation properties rather than computing an upscaled coefficient. The effective model is then a discrete one represented by the corresponding stiffness matrix and possibly right-hand side. The computation of an effective coefficient is, however, often favorable and this paper reinterprets and modifies the LOD method in this regard.

To this end, we revisit the LOD method of [28]. The original method employs finite element basis functions that are modified by a fine-scale correction with a slightly larger support. We show that it is possible to rewrite the method by means of a discrete integral operator acting on standard finite element spaces. The discrete operator is of nonlocal nature and is not necessarily associated with a differential operator on the energy space $H_0^1(\Omega)$ (for the physical domain $\Omega \subseteq \mathbb{R}^d$). The observation scale H is associated with some quasi-uniform mesh \mathcal{T}_H of width H . We are able to show that there is a discrete effective nonlocal model represented by an integral kernel $\mathcal{A}_H \in L^\infty(\Omega \times \Omega; \mathbb{R}^{d \times d})$ such that the problem is well-posed on a finite element space V_H with similar constants and satisfies

$$\sup_{f \in L^2(\Omega) \setminus \{0\}} \frac{\|u(f) - u_H(f)\|_{L^2(\Omega)}}{\|f\|_{L^2(\Omega)}} \lesssim \sup_{f \in L^2(\Omega) \setminus \{0\}} \inf_{v_H \in V_H} \frac{\|u(f) - v_H\|_{L^2(\Omega)}}{\|f\|_{L^2(\Omega)}} + H^2.$$

Based on the exponential decay of that kernel \mathcal{A}_H away from the diagonal, we propose a quasi-local and sparse formulation as an approximation. The storage requirement for the quasi-local kernel is $\mathcal{O}(H^{-d} |\log H|)$.

For an even stronger compression to $\mathcal{O}(H^{-d})$ information, one can replace \mathcal{A}_H by a local and piecewise constant tensor field A_H . It turns out that this localized effective coefficient A_H coincides with the homogenized coefficient of classical homogenization in the periodic case provided that the structure of the coefficient is slightly stronger than only periodic and that the mesh is suitably aligned with the periodicity pattern. In this sense, the results of this paper bridge the multiscale method of [28] with classical analytical techniques and numerical methods such as HMM. With regard to the recent reinterpretation of the multiscale method in [25], the paper even connects all the way from analytic homogenization to the theory of iterative solvers.

This new representation of the multiscale method turns out to be particularly attractive for computational stochastic homogenization [15]. A further advantage of our numerical techniques when compared with classical analytical techniques is that they are still applicable in the general nonperiodic case, where the local numerical model yields reasonable results whenever a certain quantitative homogenization criterion is satisfied, which can be checked a posteriori through a computable model

error estimator. Almost optimal convergence rates can be proved under reasonable assumptions on the data.

The structure of this article is as follows. After the preliminaries on the model problem and notation from section 2, we review the LOD method of [28] and introduce the quasi-local effective discrete coefficients in section 3. In section 4, we present the error analysis for the localized effective coefficient. Section 5 studies the particular case of a periodic coefficient. We present numerical results in section 6.

Standard notation on Lebesgue and Sobolev spaces applies throughout this paper. The notation $a \lesssim b$ abbreviates $a \leq Cb$ for some constant C that is independent of the mesh-size but may depend on the contrast of the coefficient A ; $a \approx b$ abbreviates $a \lesssim b \lesssim a$. The symmetric part of a quadratic matrix M is denoted by $\text{sym}(M)$.

2. Model problem and notation. This section describes the model problem and some notation on finite element spaces.

2.1. Model problem. Let $\Omega \subseteq \mathbb{R}^d$ for $d \in \{1, 2, 3\}$ be an open Lipschitz polytope. We consider the prototypical model problem

$$(2.1) \quad -\operatorname{div}(A\nabla u) = f \quad \text{in } \Omega, \quad u|_{\partial\Omega} = 0.$$

The coefficient $A \in L^\infty(\Omega; \mathbb{R}^{d \times d})$ is assumed to be symmetric and to satisfy the following uniform spectral bounds:

$$(2.2) \quad 0 < \alpha \leq \operatorname{ess\,inf}_{x \in \Omega} \inf_{\xi \in \mathbb{R}^d \setminus \{0\}} \frac{\xi \cdot (A(x)\xi)}{\xi \cdot \xi} \leq \operatorname{ess\,sup}_{x \in \Omega} \sup_{\xi \in \mathbb{R}^d \setminus \{0\}} \frac{\xi \cdot (A(x)\xi)}{\xi \cdot \xi} \leq \beta.$$

The symmetry of A is not essential for our analysis and is assumed for simpler notation. The weak form employs the Sobolev space $V := H_0^1(\Omega)$ and the bilinear form a defined, for any $v, w \in V$, by

$$a(v, w) := (A\nabla v, \nabla w)_{L^2(\Omega)}.$$

Given $f \in L^2(\Omega)$ and the linear functional

$$F : V \rightarrow \mathbb{R} \quad \text{with } F(v) := \int_{\Omega} f v \, dx \quad \text{for any } v \in V,$$

the weak form seeks $u \in V$ such that

$$(2.3) \quad a(u, v) = F(v) \quad \text{for all } v \in V.$$

2.2. Finite element spaces. Let \mathcal{T}_H be a quasi-uniform regular triangulation of Ω and let V_H denote the standard P_1 finite element space, that is, the subspace of V consisting of piecewise first-order polynomials.

Given any subdomain $S \subseteq \overline{\Omega}$, define its neighborhood via

$$\mathbf{N}(S) := \operatorname{int}(\cup\{T \in \mathcal{T}_H : T \cap \overline{S} \neq \emptyset\}).$$

Furthermore, we introduce for any $m \geq 2$ the patch extensions

$$\mathbf{N}^1(S) := \mathbf{N}(S) \quad \text{and} \quad \mathbf{N}^m(S) := \mathbf{N}(\mathbf{N}^{m-1}(S)).$$

Throughout this paper, we assume that the coarse-scale mesh \mathcal{T}_H belongs to a family of quasi-uniform triangulations. The global mesh-size reads $H := \max\{\operatorname{diam}(T) : T \in \mathcal{T}_H\}$.

Note that the shape-regularity implies that there is a uniform bound $C(m)$ on the number of elements in the m th-order patch, $\text{card}\{K \in \mathcal{T}_H : K \subseteq \overline{N^m(T)}\} \leq C(m)$ for all $T \in \mathcal{T}_H$. The constant $C(m)$ depends polynomially on m . The set of interior $(d - 1)$ -dimensional hyper faces of \mathcal{T}_H is denoted by \mathcal{F}_H . For a piecewise continuous function φ , we denote the jump across an interior edge by $[\varphi]_F$, where the index F will sometimes be omitted for brevity. The space of piecewise constant $d \times d$ matrix fields is denoted by $P_0(\mathcal{T}_H; \mathbb{R}^{d \times d})$.

Let $I_H : V \rightarrow V_H$ be a surjective quasi-interpolation operator that acts as a H^1 -stable and L^2 -stable quasi-local projection in the sense that $I_H \circ I_H = I_H$ and that for any $T \in \mathcal{T}_H$ and all $v \in V$ there holds

$$(2.4) \quad H^{-1}\|v - I_H v\|_{L^2(T)} + \|\nabla I_H v\|_{L^2(T)} \leq C_{I_H} \|\nabla v\|_{L^2(N(T))},$$

$$(2.5) \quad \|I_H v\|_{L^2(T)} \leq C_{I_H} \|v\|_{L^2(N(T))}.$$

Since I_H is a stable projection from V to V_H , any $v \in V$ is quasi-optimally approximated by $I_H v$ in the $L^2(\Omega)$ norm as well as in the $H^1(\Omega)$ norm. One possible choice is to define $I_H := E_H \circ \Pi_H$, where Π_H is the L^2 projection onto the space $P_1(\mathcal{T}_H)$ of piecewise affine (possibly discontinuous) functions and E_H is the averaging operator that maps $P_1(\mathcal{T}_H)$ to V_H by assigning to each free vertex the arithmetic mean of the corresponding function values of the neighboring cells, that is, for any $v \in P_1(\mathcal{T}_H)$ and any free vertex z of \mathcal{T}_H ,

$$(2.6) \quad (E_H(v))(z) = \sum_{\substack{T \in \mathcal{T}_H \\ \text{with } z \in T}} v|_T(z) / \text{card}\{K \in \mathcal{T}_H : z \in K\}.$$

This choice of I_H is employed in our numerical experiments.

3. Nonlocal effective coefficient. We introduce a modified version of the LOD method of [28, 21] and its localization. We give a new interpretation by means of a nonlocal effective coefficient and present an a priori error estimate.

3.1. A modified LOD method. Let $W := \ker I_H \subseteq V$ denote the kernel of I_H . Given any $T \in \mathcal{T}_H$ and $j \in \{1, \dots, d\}$, the element corrector $q_{T,j} \in W$ is the solution of the variational problem

$$(3.1) \quad a(w, q_{T,j}) = \int_T \nabla w \cdot (Ae_j) \, dx \quad \text{for all } w \in W.$$

Here e_j is the j th standard Cartesian unit vector in \mathbb{R}^d . The gradient of any $v_H \in V_H$ has the representation

$$\nabla v_H = \sum_{T \in \mathcal{T}_H} \sum_{j=1}^d (\partial_j v_H|_T) e_j.$$

Given any $v_H \in V_H$, define the corrector $\mathcal{C}v_H$ by

$$(3.2) \quad \mathcal{C}v_H = \sum_{T \in \mathcal{T}_H} \sum_{j=1}^d (\partial_j v_H|_T) q_{T,j}.$$

We remark that for any $v_H \in V_H$ the gradient ∇v_H is piecewise constant and, thus, $\mathcal{C}v_H$ is a finite linear combination of the element correctors $q_{T,j}$. It is readily verified that, for any $v_H \in V_H$, $\mathcal{C}v_H$ is the a -orthogonal projection on W , i.e.,

$$(3.3) \quad a(w, v_H - \mathcal{C}v_H) = 0 \quad \text{for all } w \in W.$$

Clearly, by (3.3), the projection $\mathcal{C}v \in W$ is well-defined for any $v \in V$. The representation (3.2) for discrete functions will, however, be useful in this work.

The LOD method in its version from [28] seeks $\bar{u}_H \in V_H$ such that

$$(3.4) \quad a((1 - \mathcal{C})\bar{u}_H, (1 - \mathcal{C})v_H) = F((1 - \mathcal{C})v_H) \quad \text{for all } v_H \in V_H.$$

By (3.3), it is clear that this is equivalent to

$$(3.5) \quad a(\bar{u}_H, (1 - \mathcal{C})v_H) = F((1 - \mathcal{C})v_H) \quad \text{for all } v_H \in V_H.$$

A variant of this multiscale method employs a problem-independent right-hand side and seeks $u_H \in V_H$ such that

$$a((1 - \mathcal{C})u_H, (1 - \mathcal{C})v_H) = F(v_H) \quad \text{for all } v_H \in V_H$$

or, equivalently,

$$(3.6) \quad a(u_H, (1 - \mathcal{C})v_H) = F(v_H) \quad \text{for all } v_H \in V_H.$$

3.2. Localization of the corrector problems. Here, we briefly describe the localization technique of [28]. It was shown in [28] and [21, Lemma 4.9] that the method is localizable in the sense that any $T \in \mathcal{T}_H$ and any $j \in \{1, \dots, d\}$ satisfy

$$(3.7) \quad \|\nabla q_{T,j}\|_{L^2(\Omega \setminus \mathbb{N}^m(T))} \lesssim \exp(-cm) \|e_j\|_{L^2(T)}, \quad m \geq 0.$$

The exponential decay from (3.7) suggests to localize the computation (3.1) of the corrector belonging to an element $T \in \mathcal{T}_H$ to a smaller domain, namely, the extended element patch $\Omega_T := \mathbb{N}^\ell(T)$ of order ℓ . The nonnegative integer ℓ is referred to as the *oversampling parameter*. Let $W_{\Omega_T} \subseteq W$ denote the space of functions from W that vanish outside Ω_T . On the patch, in analogy to (3.1), for any $v_H \in V_H$, any $T \in \mathcal{T}_H$, and any $j \in \{1, \dots, d\}$, the function $q_{T,j}^{(\ell)} \in W_{\Omega_T}$ solves

$$(3.8) \quad \int_{\Omega_T} \nabla w \cdot (A \nabla q_{T,j}^{(\ell)}) dx = \int_T \nabla w \cdot (A e_j) dx \quad \text{for all } w \in W_{\Omega_T}.$$

Given $v_H \in V_H$, we define the corrector $\mathcal{C}^{(\ell)}v_H \in W$ by

$$(3.9) \quad \mathcal{C}^{(\ell)}v_H = \sum_{T \in \mathcal{T}_H} \sum_{j=1}^d (\partial_j v_H|_T) q_{T,j}^{(\ell)}.$$

A practical variant of (3.6) is to seek $u_H^{(\ell)} \in V_H$ such that

$$(3.10) \quad a(u_H^{(\ell)}, (1 - \mathcal{C}^{(\ell)})v_H) = F(v_H) \quad \text{for all } v_H \in V_H.$$

This procedure is indispensable for actual computations and the effect of the truncation of the domain on the error of the multiscale method was analyzed in [28] and [21]. We will provide the error analysis for the method (3.10) in subsection 3.4.

3.3. Definition of the quasi-local effective coefficient. In this subsection, we do not make any specific choice for the oversampling parameter ℓ . In particular, the analysis covers the case that all element patches Ω_T equal the whole domain Ω . We denote the latter case formally by $\ell = \infty$.

We reinterpret the left-hand side of (3.10) as a nonlocal operator acting on standard finite element functions. To this end, consider any $u_H, v_H \in V_H$. We have

$$a(u_H, (1 - \mathcal{C}^{(\ell)})v_H) = \int_{\Omega} \nabla u_H \cdot (A \nabla v_H) dx - \int_{\Omega} \nabla u_H \cdot (A \mathcal{C}^{(\ell)} \nabla v_H) dx.$$

The second term can be expanded with (3.9) as

$$\begin{aligned} & \int_{\Omega} \nabla u_H \cdot (A \nabla \mathcal{C}^{(\ell)} v_H) dx \\ &= \sum_{T \in \mathcal{T}_H} \sum_{k=1}^d (\partial_k v_H|_T) \int_{\Omega} \nabla u_H \cdot (A \nabla q_{T,k}^{(\ell)}) dx \\ &= \sum_{K, T \in \mathcal{T}_H} \int_K \nabla u_H \cdot \left(\sum_{k=1}^d \int_K (A(y) \nabla q_{T,k}^{(\ell)}(y)) dy (\partial_k v_H|_T) \right) dx \\ &= \sum_{K, T \in \mathcal{T}_H} |K| |T| \nabla u_H|_K \cdot (\mathcal{K}_{T,K} \nabla v_H|_T) \end{aligned}$$

for the matrix $\mathcal{K}_{T,K}^{(\ell)}$ defined for any $K, T \in \mathcal{T}_H$ by

$$(\mathcal{K}_{T,K}^{(\ell)})_{j,k} := \frac{1}{|T| |K|} e_j \cdot \int_K A \nabla q_{T,k}^{(\ell)} dx.$$

Define the piecewise constant matrix field over $\mathcal{T}_H \times \mathcal{T}_H$, for $T, K \in \mathcal{T}_H$ by

$$\mathcal{A}_H^{(\ell)}|_{T,K} := \frac{\delta_{T,K}}{|K|} \int_T A dx - \mathcal{K}_{T,K}^{(\ell)}$$

(where δ is the Kronecker symbol) and the bilinear form $\mathbf{a}^{(\ell)}$ on $V_H \times V_H$ by

$$\mathbf{a}^{(\ell)}(v_H, z_H) := \int_{\Omega} \int_{\Omega} \nabla v_H(y) \cdot (\mathcal{A}_H^{(\ell)}(x, y) \nabla z_H(x)) dy dx \quad \text{for any } v_H, z_H \in V_H.$$

We obtain for all $v_H, z_H \in V_H$ that

$$(3.11) \quad a(v_H, (1 - \mathcal{C}^{(\ell)})z_H) = \mathbf{a}^{(\ell)}(v_H, z_H).$$

Remark 3.1 (notation). For simplices $T, K \in \mathcal{T}_H$ with $x \in T$ and $y \in K$, we will sometimes write $\mathcal{K}^{(\ell)}(x, y)$ instead of $\mathcal{K}_{T,K}^{(\ell)}$ (with analogous notation for $\mathcal{A}^{(\ell)}$).

Next, we state the equivalence of two multiscale formulations.

PROPOSITION 3.2. *A function $u_H^{(\ell)} \in V_H$ solves (3.10) if and only if it solves*

$$(3.12) \quad \mathbf{a}^{(\ell)}(u_H^{(\ell)}, v_H) = F(v_H).$$

Proof. This follows directly from the representation (3.11). □

Remark 3.3. For $d = 1$ and I_H the standard nodal interpolation operator, the corrector problems localize to one element and the presented multiscale approach coincides with various known methods (homogenization, multiscale finite element method (MSFEM)). The resulting effective coefficient $\mathcal{A}_H^{(\ell)}$ is diagonal and, thus, local. This is no longer the case for $d \geq 2$.

3.4. Error analysis. This subsection presents an error estimate for the error produced by the method (3.10) (and so by the method (3.12)). We begin by briefly summarizing some results from [28].

LEMMA 3.4. *Let $u \in V$ solve (2.3) and $\bar{u}_H \in V_H$ solve (3.4). Then we have the following properties:*

- (i) \bar{u}_H coincides with the quasi interpolation of u , i.e., $\bar{u}_H = I_H u$.
- (ii) The Galerkin orthogonality $a(u - (1 - \mathcal{C})I_H u, (1 - \mathcal{C})v_H) = 0$ for all $v_H \in V_H$ is satisfied.
- (iii) The error satisfies $\|\nabla(u - (1 - \mathcal{C})\bar{u}_H)\|_{L^2(\Omega)} \lesssim H\|f\|_{L^2(\Omega)}$.

Proof. See [28] for proofs. □

We define the following worst-case best-approximation error:

$$(3.13) \quad \mathbf{wcb}\mathbf{a}(A, \mathcal{T}_H) := \sup_{g \in L^2(\Omega) \setminus \{0\}} \inf_{v_H \in V_H} \frac{\|u(g) - v_H\|_{L^2(\Omega)}}{\|g\|_{L^2(\Omega)}},$$

where for $g \in L^2(\Omega)$, $u(g) \in V$ solves (2.3) with right-hand-side g . Standard interpolation and stability estimates show that always $\mathbf{wcb}\mathbf{a}(A, \mathcal{T}_H) \lesssim H$, but it may behave better in certain regimes. For example, in a periodic homogenization problem with some small parameter ε and some smooth homogenized solution $u_0 \in H^2(\Omega)$, the best-approximation error is dominated by the best-approximation error of u_0 in the regime $H \lesssim \sqrt{\varepsilon}$ where it scales like H^2 . By contrast, the error is typically not improved in the regime $\sqrt{\varepsilon} \gtrsim H \gtrsim \varepsilon$. This nonlinear behavior of the best-approximation error in the preasymptotic regime is prototypical for homogenization problems with scale separation and explains why the rough bound H is suboptimal.

The following result states an L^2 error estimate for the method (3.6). The result is surprising because the perturbation of the right-hand side seems to be of order H at first glance. In cases of scale separation the quadratic rate is indeed observed in the regime $H \lesssim \sqrt{\varepsilon}$ and cannot be explained by naive estimates.

PROPOSITION 3.5. *The solutions $u \in V$ to (2.3) and $u_H \in V_H$ to (3.6) for right-hand-side $f \in L^2(\Omega)$ satisfy the following error estimate:*

$$\frac{\|u - u_H\|_{L^2(\Omega)}}{\|f\|_{L^2(\Omega)}} \lesssim H^2 + \mathbf{wcb}\mathbf{a}(A, \mathcal{T}_H).$$

Proof. Let $f \in L^2(\Omega) \setminus \{0\}$ and let $\bar{u}_H \in V_H$ solve (3.5). We begin by analyzing the error $e_H := u_H - \bar{u}_H$. Let $z \in V$ denote the solution to

$$a(v, z) = (e_H, I_H v)_{L^2(\Omega)} \quad \text{for all } v \in V.$$

To see that the right-hand side is indeed represented by an L^2 function, note that I_H is continuous on $L^2(\Omega)$ and, hence, the right-hand side has a Riesz representative $\tilde{e} \in L^2(\Omega)$ such that $(e_H, I_H v)_{L^2(\Omega)} = (\tilde{e}, v)_{L^2(\Omega)}$. In particular, z solves (2.3) with right-hand-side \tilde{e} . Its L^2 norm is bounded with (2.5) as follows:

$$\|\tilde{e}\|_{L^2(\Omega)}^2 = (e_H, I_H \tilde{e})_{L^2(\Omega)} \lesssim \|e_H\|_{L^2(\Omega)} \|\tilde{e}\|_{L^2(\Omega)};$$

hence

$$(3.14) \quad \|\tilde{e}\|_{L^2(\Omega)} \lesssim \|e_H\|_{L^2(\Omega)}.$$

We note that, for any $w \in W$, we have $a(w, z) = (e_H, I_H w)_{L^2(\Omega)} = 0$. Thus, we have $a(e_H, \mathcal{C}z) = a(\mathcal{C}e_H, z) = 0$. With $(1 - \mathcal{C})z = (1 - \mathcal{C})I_H z$ we conclude

$$(3.15) \quad \|e_H\|_{L^2(\Omega)}^2 = a(e_H, z) = a(e_H, (1 - \mathcal{C})I_H z).$$

Elementary algebraic manipulations with the projection I_H show that

$$-\mathcal{C}I_H z = (1 - I_H)((1 - \mathcal{C})I_H z - z) + (1 - I_H)z.$$

The relation (3.15) and the solution properties (3.5) and (3.6), thus, lead to

$$(3.16) \quad \|e_H\|_{L^2(\Omega)}^2 = F(\mathcal{C}I_H z) = |F((1 - I_H)((1 - \mathcal{C})I_H z - z)) + F((1 - I_H)z)|.$$

We proceed by estimating the two terms on the right-hand side of (3.16) separately. For the second term in (3.16), the L^2 -best-approximation property of I_H and (3.14) reveal

$$(3.17) \quad \begin{aligned} |F((1 - I_H)z)| &\lesssim \|f\|_{L^2(\Omega)} \|\tilde{e}\|_{L^2(\Omega)} \inf_{v_H \in V_H} \frac{\|z - v_H\|_{L^2(\Omega)}}{\|\tilde{e}\|_{L^2(\Omega)}} \\ &\lesssim \|f\|_{L^2(\Omega)} \|e_H\|_{L^2(\Omega)} \mathbf{wcb}a(A, \mathcal{T}_H). \end{aligned}$$

For the first term in (3.16), we obtain with the stability of I_H and the Cauchy inequality that

$$|F((1 - I_H)((1 - \mathcal{C})I_H z - z))| \lesssim \|f\|_{L^2(\Omega)} \|z - (1 - \mathcal{C})I_H z\|_{L^2(\Omega)}.$$

Let $\tilde{g} := z - (1 - \mathcal{C})I_H z$ and let $\zeta \in V$ denote the solution to

$$a(\zeta, v) = (\tilde{g}, v)_{L^2(\Omega)} \quad \text{for all } v \in V.$$

As stated in Lemma 3.4(i), the function $I_H z \in V_H$ is the Galerkin approximation to z with method (3.4) with right-hand-side \tilde{e} . We, thus, have by symmetry of a and the Galerkin orthogonality from Lemma 3.4(ii) that

$$\begin{aligned} \|z - (1 - \mathcal{C})I_H z\|_{L^2(\Omega)}^2 &= a(\zeta, z - (1 - \mathcal{C})I_H z) \\ &= a(\zeta - (1 - \mathcal{C})I_H \zeta, z - (1 - \mathcal{C})I_H z). \end{aligned}$$

Continuity of a and Lemma 3.4(iii) reveal that this is bounded by

$$H^2 \|\tilde{g}\|_{L^2(\Omega)} \|\tilde{e}\|_{L^2(\Omega)} = H^2 \|z - (1 - \mathcal{C})I_H z\|_{L^2(\Omega)} \|\tilde{e}\|_{L^2(\Omega)}.$$

Altogether, with (3.16),

$$\frac{\|e_H\|_{L^2(\Omega)}}{\|f\|_{L^2(\Omega)}} \lesssim H^2 + \mathbf{wcb}a(A, \mathcal{T}_H).$$

Since

$$\frac{\|u - \bar{u}_H\|_{L^2(\Omega)}}{\|f\|_{L^2(\Omega)}} \lesssim \mathbf{wcb}a(A, \mathcal{T}_H)$$

(which follows from the fact that $\bar{u}_H = I_H u$), the triangle inequality concludes the proof. \square

With similar arguments it is possible to prove that the coupling $\ell \approx |\log H|$ is sufficient to derive the error bound

$$(3.18) \quad \|u - u_H^{(\ell)}\|_{L^2(\Omega)} \lesssim (H^2 + \mathbf{wcb}\mathbf{a}(A, \mathcal{T}_H)) \|f\|_{L^2(\Omega)}.$$

The proof is based on a similar argument as in Proposition 3.5: Since the L^2 distance of $u - \bar{u}_H^{(\ell)}$ is controlled by the right-hand side of (3.18) [21], where $\bar{u}_H^{(\ell)}$ solves a modified version of (3.10) with right-hand-side $F((1 - \mathcal{C}^{(\ell)})v_H)$, it is sufficient to control $u_H^{(\ell)} - \bar{u}_H^{(\ell)}$ in the L^2 norm. This can be done with a duality argument similar to that from the proof of Proposition 3.5. The additional tool needed therein is the fact that

$$\|\nabla(\mathcal{C} - \mathcal{C}^{(\ell)})I_H z\|_{L^2(\Omega)} \lesssim \exp(-c\ell)C(\ell)\|\nabla z\|_{L^2(\Omega)}$$

for the dual solution z (see [21, Proof of Theorem 4.13] for an outline of a proof), where $C(\ell)$ is an overlap constant depending polynomially on ℓ . The choice of $\ell \approx |\log H|$ therefore leads to (3.18). The details are omitted here and the reader is referred to [28, 21, 36, 25].

4. Local effective coefficient. Throughout this section we consider oversampling parameters chosen as $\ell \approx |\log H|$.

4.1. Definition of the local effective coefficient. The exponential decay motivates us to approximate the nonlocal bilinear form $\mathbf{a}^{(\ell)}(\cdot, \cdot)$ by a quadrature-like procedure: Define the piecewise constant coefficient $A_H^{(\ell)} \in P_0(\mathcal{T}_H; \mathbb{R}^{d \times d})$ by

$$A_H^{(\ell)}|_T := \int_T A \, dx - \sum_{K \in \mathcal{T}_H} |K| \mathcal{K}_{T,K}^{(\ell)}.$$

and the bilinear form $\tilde{\mathbf{a}}^{(\ell)}$ on $V \times V$ by

$$\tilde{\mathbf{a}}^{(\ell)}(u, v) := \int_{\Omega} \nabla u \cdot (A_H^{(\ell)} \nabla v) \, dx.$$

Remark 4.1. In analogy to classical periodic homogenization, the local effective coefficient $A_H^{(\ell)}$ can be written as

$$\begin{aligned} (A_H^{(\ell)})_{j,k}|_T &= |T|^{-1} \int_{\Omega_T} e_j \cdot (A(\chi_T e_k - \nabla q_{T,k}^{(\ell)})) \\ &= |T|^{-1} \int_{\Omega_T} (e_j - \nabla q_{T,j}^{(\ell)}) \cdot (A(\chi_T e_k - \nabla q_{T,k}^{(\ell)})) \end{aligned}$$

for the characteristic function χ_T of T and the slightly enlarged averaging domain Ω_T . See section 5 for further analogies to homogenization theory in the periodic case.

The localized multiscale method is to seek $\tilde{u}_H^{(\ell)} \in V_H$ such that

$$(4.1) \quad \tilde{\mathbf{a}}^{(\ell)}(\tilde{u}_H^{(\ell)}, v_H) = F(v_H) \quad \text{for all } v_H \in V_H.$$

The unique solvability of (4.1) is not guaranteed a priori. It must be checked a posteriori whether positive spectral bounds α_H, β_H on $A_H^{(\ell)}$ exist in the sense of (2.2). Throughout this paper we assume that such bounds exist, that is, we assume that there exist positive numbers α_H, β_H such that

$$(4.2) \quad \alpha_H |\xi|^2 \leq \xi \cdot (A_H^{(\ell)}(x)\xi) \leq \beta_H |\xi|^2$$

for all $\xi \in \mathbb{R}^d$ and almost all $x \in \Omega$.

4.2. Error analysis. The goal of this section is to establish an error estimate for the error

$$\|u - \tilde{u}_H^{(\ell)}\|_{L^2(\Omega)}.$$

Let $u_H^{(\ell)} \in V_H$ solve (3.10). Then the error estimate (3.18) leads to the a priori error estimate

$$(4.3) \quad \|u - u_H^{(\ell)}\|_{L^2(\Omega)} \lesssim (H^2 + \mathbf{wcb}a(A, \mathcal{T}_H)) \|f\|_{L^2(\Omega)}.$$

We employ the triangle inequality and merely estimate the difference $\|u_H^{(\ell)} - \tilde{u}_H^{(\ell)}\|_{L^2(\Omega)}$.

With the finite localization parameter ℓ , the quasi-local coefficient $\mathcal{A}^{(\ell)}$ is sparse in the sense that $\mathcal{A}^{(\ell)}(x, y) = 0$ whenever $|x - y| > C\ell H$. We note the following lemma, which will be employed in the error analysis.

LEMMA 4.2. *Given some $x \in \Omega$ with $x \in T$ for some $T \in \mathcal{T}_H$, we have*

$$\|\mathcal{K}^{(\ell)}(x, y)\|_{L^2(\Omega, dy)} \lesssim H^{-d/2}.$$

Proof. From the definition of $\mathcal{K}^{(\ell)}$, the boundedness of A , and the Hölder inequality we obtain for any $j, k \in \{1, \dots, d\}$ that

$$|(\mathcal{K}_{T,K}^{(\ell)})_{j,k}| \lesssim \frac{1}{|T||K|} \|\nabla q_{T,k}\|_{L^1(K)} \lesssim \frac{1}{|T||K|^{1/2}} \|\nabla q_{T,k}\|_{L^2(K)}.$$

Hence, we conclude with the stability of problem (3.8) and $\|e_k\|_{L^2(T)}^2 = |T|$ that

$$\|\mathcal{K}^{(\ell)}(x, y)\|_{L^2(\Omega, dy)}^2 = \sum_{K \in \mathcal{T}_H} |K| |\mathcal{K}_{T,K}^{(\ell)}|^2 = |T|^{-2} \|\nabla q_{T,k}\|_{L^2(\Omega)}^2 \lesssim H^{-d}.$$

This implies the assertion. □

In what follows, we abbreviate

$$(4.4) \quad \rho := CH|\log H|$$

for some appropriately chosen constant C .

PROPOSITION 4.3 (error estimate I). *Assume that (4.2) is satisfied. Let $u_H^{(\ell)} \in V_H$ solve (3.12) and let $\tilde{u}_H^{(\ell)}$ solve (4.1). Then,*

$$\|\nabla(u_H^{(\ell)} - \tilde{u}_H^{(\ell)})\|_{L^2(\Omega)} \lesssim H^{-d/2} \left\| \|\nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x)\|_{L^2(B_\rho(x), dy)} \right\|_{L^2(\Omega, dx)}.$$

Proof. Denote $e_H := \tilde{u}_H^{(\ell)} - u_H^{(\ell)}$. In the idealized case, $\ell = \infty$, the orthogonality (3.3) and relation (3.11) show that

$$\|\nabla(1 - \mathcal{C}^{(\ell)})e_H\|_{L^2(\Omega)}^2 \lesssim \mathbf{a}^{(\ell)}(e_H, e_H).$$

The case $\ell \gtrsim |\log H|$ again follows ideas from [28] with the exponential-in- ℓ closeness of \mathcal{C} and \mathcal{C}_ℓ and is merely sketched here. From the stability of I_H and the properties of the fine-scale projection $\mathcal{C}^{(\ell)}$ we observe (with contrast-dependent constants)

$$\begin{aligned} \|\nabla e_H\|_{L^2(\Omega)}^2 &= \|\nabla I_H e_H\|_{L^2(\Omega)}^2 = \|\nabla I_H(1 - \mathcal{C}^{(\ell)})e_H\|_{L^2(\Omega)}^2 \\ &\lesssim \|\nabla(1 - \mathcal{C}^{(\ell)})e_H\|_{L^2(\Omega)}^2 \\ &\lesssim \mathbf{a}^{(\ell)}(e_H, e_H) + \exp(-c\ell) \|\nabla e_H\|_{L^2(\Omega)}^2 \end{aligned}$$

for some constant $c > 0$. Hence, with positive constants C_1, C_2 ,

$$\|\nabla e_H\|_{L^2(\Omega)}^2 \leq C_1 \mathbf{a}^{(\ell)}(e_H, e_H) + C_2 \exp(-c\ell) \|\nabla e_H\|_{L^2(\Omega)}^2.$$

If, for some sufficiently large r , the parameter ℓ is chosen to satisfy $\ell \geq r|\log H|$ such that $C_2 \exp(-c\ell) \leq 1/2$, then the second term on the right-hand side can be absorbed. Thus, we proceed with (3.12) and (4.1) as

$$\|\nabla e_H\|_{L^2(\Omega)}^2 \lesssim \mathbf{a}^{(\ell)}(\tilde{u}_H^{(\ell)} - u_H^{(\ell)}, e_H) = \mathbf{a}^{(\ell)}(\tilde{u}_H^{(\ell)}, e_H) - \tilde{\mathbf{a}}^{(\ell)}(\tilde{u}_H^{(\ell)}, e_H).$$

The right-hand side can be rewritten as

$$\begin{aligned} & \mathbf{a}^{(\ell)}(\tilde{u}_H^{(\ell)}, e_H) - \tilde{\mathbf{a}}^{(\ell)}(\tilde{u}_H^{(\ell)}, e_H) \\ &= \int_{\Omega} \int_{\Omega} (\nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x)) \cdot \left[\mathcal{A}_H^{(\ell)}(x, y) \nabla e_H(x) \right] dy dx \\ & \quad + \int_{\Omega} \left[\tilde{u}_H^{(\ell)}(x) \cdot \left(\int_{\Omega} \mathcal{A}_H^{(\ell)}(x, y) dy - A_H^{(\ell)}(x) \right) \nabla e_H(x) \right] dx. \end{aligned}$$

The second term vanishes by definition of $A_H^{(\ell)}$. Hence, the combination of the preceding arguments with the Cauchy inequality leads to

$$\|\nabla e_H\|_{L^2(\Omega)}^2 \lesssim \|\nabla e_H\|_{L^2(\Omega)} \left\| \int_{B_{\rho}(x)} \mathcal{A}_H^{(\ell)}(x, y)^* (\nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x)) dy \right\|_{L^2(\Omega, dx)},$$

where it was used that $\mathcal{A}_H^{(\ell)}(x, y) = 0$ whenever $|x - y| > \rho$. Note that $(\nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x)) = 0$ for all x and y that belong to the same element $T \in \mathcal{T}_H$. Thus, $\mathcal{A}_H^{(\ell)}(x, y)$ in the above expression can be replaced by $\mathcal{K}_H^{(\ell)}(x, y)$. This and division by $\|\nabla e_H\|_{L^2(\Omega)}$ lead to

$$(4.5) \quad \|\nabla e_H\|_{L^2(\Omega)} \lesssim \sqrt{\int_{\Omega} \left| \int_{B_{\rho}(x)} \mathcal{K}_H^{(\ell)}(x, y)^* (\nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x)) dy \right|^2 dx}.$$

This term can be bounded with the Cauchy inequality and Lemma 4.2 by

$$\begin{aligned} & \sqrt{\int_{\Omega} \left\| \mathcal{K}_H^{(\ell)}(x, y) \right\|_{L^2(B_{\rho}(x), dy)} \left\| \nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x) \right\|_{L^2(B_{\rho}(x), dy)} \right|^2 dx} \\ & \lesssim H^{-d/2} \left\| \left\| \nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x) \right\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)}. \end{aligned}$$

This finishes the proof. □

It is worth noting that the error bound in Proposition 4.3 can be evaluated without knowledge of the exact solution. Hence, Proposition 4.3 can be regarded as an a posteriori error estimate. Formula (4.5) could also be an option if it is available. We expect Proposition 4.3 to be rather sharp. Below we provide the main a priori error estimate, Proposition 4.5, which is fundamental for the mentioned link between analytical and numerical homogenization. The following technical lemma is required.

LEMMA 4.4 (existence of a regularized coefficient). *Let $A_H \in P_0(\mathcal{T}_H; \mathbb{R}^{d \times d})$ be a piecewise constant field of $d \times d$ matrices that satisfies the spectral bounds (4.2).*

Then there exists a Lipschitz continuous coefficient $A_H^{reg} \in W^{1,\infty}(\Omega; \mathbb{R}^{d \times d})$ satisfying the following three properties: (1) The piecewise integral mean is conserved, i.e.,

$$\int_T A_H^{reg} dx = \int_T A_H dx \quad \text{for all } T \in \mathcal{T}_H.$$

(2) The eigenvalues of $\text{sym}(A_H^{reg})$ lie in the interval $[\alpha_H/2, 2\beta_H]$. (3) The derivative satisfies the bound

$$\|\nabla A_H^{reg}\|_{L^\infty(\Omega)} \leq C\eta(A_H)$$

for some constant C that depends on the shape-regularity of \mathcal{T}_H and for the expression

$$(4.6) \quad \eta(A_H) := H^{-1} \|[A_H]\|_{L^\infty(\mathcal{F}_H)} (1 + \alpha_H^{-1} \|[A_H]\|_{L^\infty(\mathcal{F}_H)}).$$

Here $[\cdot]$ defines the interelement jump and \mathcal{F}_H denotes the set of interior hyperfaces of \mathcal{T}_H .

Proof. Consider a refined triangulation \mathcal{T}_L resulting from L uniform refinements of \mathcal{T}_H . In particular, the mesh-size in \mathcal{T}_L is of the order $2^{-L}H$. Let $E_L A_H$ denote the \mathcal{T}_L -piecewise affine and continuous function that takes at every interior vertex the arithmetic mean of the nodal values of A_H on the adjacent elements of \mathcal{T}_L (similar to (2.6)). Clearly, for this convex combination the eigenvalues of $\text{sym}(E_L A_H)$ range within the interval $[\alpha_H, \beta_H]$. It is not difficult to prove that, for any $T \in \mathcal{T}_H$,

$$(4.7) \quad \int_T |A_H - E_L A_H| dx \lesssim 2^{-L} \|[A_H]\|_{L^\infty(\mathcal{F}_H(\omega_T))}$$

as well as

$$(4.8) \quad \|A_H - E_L A_H\|_{L^\infty(T)} \lesssim \|[A_H]\|_{L^\infty(\mathcal{F}_H(\omega_T))}.$$

Here, $\mathcal{F}_H(\omega_T)$ denotes the set of interior hyperfaces of \mathcal{T}_H that share a point with T . Let, for any $T \in \mathcal{T}_H$, $b_T \in H_0^1(T)$ denote a positive polynomial bubble function with $\int_T b_T dx = 1$ and $\|b_T\|_{L^\infty(T)} \approx 1$. The regularized coefficient $A_H^{reg} = E_L(A_H) + b_T \int_T (A_H - E_L(A_H)) dx$ has, for any $T \in \mathcal{T}_H$, the integral mean $\int_T A_H^{reg} dx = \int_T A_H dx$. For any $\xi \in \mathbb{R}^d$ with $|\xi| = 1$ and any $T \in \mathcal{T}_H$, the estimate (4.7) shows

$$\begin{aligned} \left| \xi \cdot \int_T (A_H - E_L A_H) dx b_T \xi \right| &\leq \left| \int_T (A_H - E_L A_H) dx b_T \right| \\ &\leq C 2^{-L} \|[A_H]\|_{L^\infty(\mathcal{F}_H(\omega_T))}. \end{aligned}$$

If L is chosen to be of the order $\lceil \log(\alpha_H^{-1} C \|[A_H]\|_{L^\infty(\mathcal{F}_H)}) \rceil$ (for small jumps of A_H it can be chosen of order 1), then

$$\left| \xi \cdot \int_T (A_H - E_L A_H) dx b_T \xi \right| \leq \alpha/2.$$

This and the triangle inequality prove the claimed spectral bound on $\text{sym}(A_H^{reg})$. For the bound on the derivative of A_H^{reg} , let $t \in \mathcal{T}_L$ and $T \in \mathcal{T}_H$ such that $t \subseteq T$. The diameter of t is of order $2^{-L}H$. Since $\|\nabla b_T\|_{L^\infty(T)} \lesssim H^{-1}$, the triangle and inverse inequalities therefore yield with the above choice of L (note that $\nabla(A_H|_T) = 0$)

$$\begin{aligned} \|\nabla A_H^{reg}\|_{L^\infty(t)} &\lesssim \|\nabla(A_H - E_L(A_H))\|_{L^\infty(t)} + H^{-1} \|A_H - E_L(A_H)\|_{L^\infty(t)} \\ &\lesssim H^{-1} \|[A_H]\|_{L^\infty(\mathcal{F}_H(\omega_T))} (1 + \alpha_H^{-1} \|[A_H]\|_{L^\infty(\mathcal{F}_H(\omega_T))}). \end{aligned}$$

This proves the assertion. □

By Lemma 4.4, there exists a coefficient $A_H^{reg} \in W^{1,\infty}(\Omega)$ such that $A_H^{(\ell)}$ is the piecewise L^2 projection of A_H^{reg} onto the piecewise constants. Let $u^{reg} \in V$ solve

$$(4.9) \quad \int_{\Omega} \nabla u^{reg} \cdot (A_H^{reg} \nabla v) \, dx = F(v) \quad \text{for all } v \in V.$$

In particular, \tilde{u}_H is the finite element approximation to u^{reg} . In the following, s refers to the $H^{1+s}(\Omega)$ regularity index of a function. Recall that the $H^{1+s}(\Omega)$ norm [2] of some function v is given by

$$(4.10) \quad \|v\|_{H^{1+s}(\Omega)} = \left[\|v\|_{H^1(\Omega)}^2 + \int_{\Omega} \int_{\Omega} \frac{|\nabla v(x) - \nabla v(y)|^2}{|x - y|^{d+2s}} \, dy \, dx \right]^{1/2}.$$

We have the following error estimate.

PROPOSITION 4.5 (error estimate II). *Let $\ell \approx |\log H|$ and assume that (4.2) is satisfied. Let $u_H^{(\ell)}$ solve (3.10) and let $\tilde{u}_H^{(\ell)}$ solve (4.1). Assume furthermore that the solution u^{reg} to (4.9) belongs to $H^{1+s}(\Omega)$ for some $0 < s \leq 1$. Then,*

$$\|\nabla(u_H^{(\ell)} - \tilde{u}_H^{(\ell)})\|_{L^2(\Omega)} \lesssim H^s |\log H|^{s+d/2} (1 + \eta(A_H^{(\ell)}))^s \|f\|_{L^2(\Omega)}.$$

Proof. Recall the estimate from Proposition 4.3,

$$\|\nabla(u_H^{(\ell)} - \tilde{u}_H^{(\ell)})\|_{L^2(\Omega)} \lesssim H^{-d/2} \left\| \|\nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)}.$$

To bound the norm on the right-hand side, we denote $e := \nabla(\tilde{u}_H^{(\ell)} - u^{reg})$ and infer with the triangle inequality

$$(4.11) \quad \begin{aligned} & \left\| \|\nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)} \\ & \leq \left\| \|e(y)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)} \\ & \quad + \left\| \|\nabla u^{reg}(y) - \nabla u^{reg}(x)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)} \\ & \quad + \left\| \|e(x)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)}. \end{aligned}$$

The square of the first term on the right-hand side of (4.11) satisfies

$$\begin{aligned} \left\| \|e(y)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)}^2 &= \int_{\Omega} \int_{B_{\rho}(x)} |e(y)|^2 \, dy \, dx \\ &= \int_{\Omega} \int_{\{x \text{ with } y \in B_{\rho}(x)\}} |e(y)|^2 \, dx \, dy \lesssim \rho^d \|e\|_{L^2(\Omega)}^2. \end{aligned}$$

Similarly, the third term on the right-hand side of (4.11) satisfies

$$\left\| \|e(x)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)}^2 = \int_{\Omega} \int_{B_{\rho}(x)} |e(x)|^2 \, dy \, dx \lesssim \rho^d \|e\|_{L^2(\Omega)}^2.$$

The second term on the right-hand side of (4.11) reads for any $0 < s < 1$ as

$$\begin{aligned} & \left\| \|\nabla u^{reg}(y) - \nabla u^{reg}(x)\|_{L^2(B_{\rho}(x), dy)} \right\|_{L^2(\Omega, dx)} \\ & = \rho^{(d+2s)/2} \left(\int_{\Omega} \int_{B_{\rho}(x)} \frac{|\nabla u^{reg}(x) - \nabla u^{reg}(y)|^2}{\rho^{d+2s}} \, dy \, dx \right)^{1/2} \\ & \lesssim \rho^{(d+2s)/2} \|u^{reg}\|_{H^{1+s}(\Omega)}. \end{aligned}$$

Here we have used the representation (4.10) and the fact that the value of the double integral increases when, first, in the denominator ρ is replaced by $|x - y|$ and thereafter the integration domain of the inner integral is replaced by Ω . In conclusion,

$$\begin{aligned} & \left\| \left\| \nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x) \right\|_{L^2(B_\rho(x), dy)} \right\|_{L^2(\Omega, dx)} \\ & \lesssim \rho^{d/2} \|e\|_{L^2(\Omega)} + \rho^{(d+2s)/2} \|u^{reg}\|_{H^{1+s}(\Omega)}. \end{aligned}$$

Since $\tilde{u}_H^{(\ell)}$ is the finite element approximation to ∇u^{reg} , standard a priori error estimates for the Galerkin projection yield

$$\|e\|_{L^2(\Omega)} \lesssim H^s \|u^{reg}\|_{H^{1+s}(\Omega)}.$$

Thus,

$$(4.12) \quad \left\| \left\| \nabla \tilde{u}_H^{(\ell)}(y) - \nabla \tilde{u}_H^{(\ell)}(x) \right\|_{L^2(B_\rho(x), dy)} \right\|_{L^2(\Omega, dx)} \lesssim \rho^{(d+2s)/2} \|u^{reg}\|_{H^{1+s}(\Omega)}.$$

If u^{reg} belongs to $H^2(\Omega)$, then the results of [18, 9, 30] lead to

$$(4.13) \quad \begin{aligned} \|u^{reg}\|_{H^2(\Omega)} & \lesssim \|A_H^{reg}\|_{W^{1,\infty}(\Omega)} (\|f\|_{L^2(\Omega)} + \|u^{reg}\|_{H^1(\Omega)}) \\ & \lesssim \|A_H^{reg}\|_{W^{1,\infty}(\Omega)} \|f\|_{L^2(\Omega)}. \end{aligned}$$

The assertion in $H^{1+s}(\Omega)$ can be proved with an operator interpolation argument. Indeed, as shown in [18], the operator $-\operatorname{div}(A^{reg}\nabla\cdot)$ maps $H^2(\Omega) \cap H_0^1(\Omega)$ to a closed subspace Y_1 of $L^2(\Omega)$. Let T denote the solution operator, which maps Y_1 to $X_1 := H^2(\Omega)$ and furthermore maps $Y_0 := L^2(\Omega)$ to $X_0 := H^1(\Omega)$. The real method of Banach space interpolation [7] shows that $H^{1+s}(\Omega) = [X_0, X_1]_{s,2}$, which together with the H^1 stability of the problem and (4.13) proves

$$\|u^{reg}\|_{H^{1+s}(\Omega)} \lesssim \|A_H^{reg}\|_{W^{1,\infty}(\Omega)}^s \|f\|_{L^2(\Omega)}.$$

The combination with Lemma 4.4 proves

$$\|u^{reg}\|_{H^{1+s}(\Omega)} \lesssim (1 + \eta(A_H^{(\ell)}))^s \|f\|_{L^2(\Omega)}.$$

The combination with Proposition 4.3 and (4.12) proves

$$\begin{aligned} \|\nabla(u_H^{(\ell)} - \tilde{u}_H^{(\ell)})\|_{L^2(\Omega)} & \lesssim H^{-d/2} \rho^{(d+2s)/2} \|u^{reg}\|_{H^{1+s}(\Omega)} \\ & \lesssim H^s |\log H|^{s+d/2} (1 + \eta(A_H^{(\ell)}))^s \|f\|_{L^2(\Omega)}. \end{aligned}$$

This implies the assertion. □

Remark 4.6 (homogenization indicator). If the relations

$$H^{-1} \|[A_H^{(\ell)}]\|_{L^\infty(\mathcal{F}_H)} \lesssim 1 \quad \text{and} \quad \alpha_H^{-1} H \lesssim 1$$

are satisfied, then the multiplicative constant in Proposition 4.5 is of moderate size. Hence, we interpret $\eta(A_H^{(\ell)})$ as a homogenization indicator and the above relations as a *homogenization criterion*.

Remark 4.7 (local mesh-refinement). We furthermore remark that local versions of $\eta(A_H^{(\ell)})$ involving the jump information $H^{-1} \|[A_H^{(\ell)}]\|_{L^\infty(F)}$ for interior interfaces F may be used as refinement indicators for local mesh-adaptation. This possibility, however, shall not be further discussed here.

Remark 4.8 (global homogenized coefficient). If the global variations of $A_H^{(\ell)}$ are small in the sense that there are positive constants c_1, c_2 such that, almost everywhere,

$$c_1|\xi|^2 \leq \xi \cdot (A_H^{(\ell)}\xi) \leq c_2|\xi|^2 \quad \text{for any } \xi \in \mathbb{R}^d$$

holds with $|c_2 - c_1| \lesssim H$, then $A_H^{(\ell)}$ can be replaced by $f_\Omega A_H^{(\ell)} dx$ without effecting the accuracy.

The combination of Proposition 4.5 with (4.3) leads to the following a priori error estimate. The parameter s therein is determined by the elliptic regularity of the model problem with a $W^{1,\infty}(\Omega)$ coefficient.

THEOREM 4.9. *Let $\ell \approx |\log H|$ and assume that (4.2) is satisfied. Let u solve (2.3) and let $\tilde{u}_H^{(\ell)}$ solve (4.1). Assume furthermore that the solution u^{reg} to (4.9) belongs to $H^{1+s}(\Omega)$ for some $0 < s \leq 1$. Then,*

$$\|u - \tilde{u}_H^{(\ell)}\|_{L^2(\Omega)} \lesssim \left(H + H^s |\log H|^{s+d/2} (1 + \eta(A_H^{(\ell)}))^s \right) \|f\|_{L^2(\Omega)}.$$

In particular, under the homogenization criterion from Remark 4.6, a convergence rate is achieved. If the domain is convex, then s can be chosen as $s = 1$, i.e., the convergence rate is linear up to a logarithmic factor.

Proof. This follows from combining Proposition 4.5 with (4.3), the triangle inequality, and the Friedrichs inequality. If the domain is convex, elliptic regularity theory [18, 9, 30] shows that $s = 1$ is an admissible choice. \square

Remark 4.10. We emphasize that $\eta(A_H^{(\ell)})$ is not an error estimator for the discretization error. It rather indicates whether the local discrete model is appropriate. If $\eta(A_H^{(\ell)})$ is close to zero, then the multiplicative constant on the right-hand side of the formula in Theorem 4.9 is of reasonable magnitude.

5. The periodic setting. In this section we justify the use of the local effective coefficient A_H in the periodic setting. We show that the procedure in its idealized form with $\ell = \infty$ recovers the classical periodic homogenization limit. We denote by $V := H^1_{\#}(\Omega)/\mathbb{R}$ the space of periodic H^1 functions with vanishing integral mean over Ω . We assume Ω to be a polytope allowing for periodic boundary conditions. We adopt the notation of section 3; in particular, $W \subseteq V$ is the kernel of the quasi-interpolation I_H , V_H is the space of piecewise affine globally continuous functions of V , and $\mathcal{C}^{(\ell)}$, a , $\tilde{a}^{(\ell)}$, $\mathbf{a}^{(\ell)}$, $\mathcal{A}_H^{(\ell)}$, $A_H^{(\ell)}$, $\mathcal{K}^{(\ell)}$ are defined as in section 3 with the underlying space V being $H^1_{\#}(\Omega)/\mathbb{R}$. We assume that the domain Ω matches with integer multiples of the period. We assume the triangulation \mathcal{T}_H to match with the periodicity pattern. For simplicial partitions this implies further symmetry assumptions. In particular, periodicity with respect to a uniform rectangular grid is not sufficient. Instead we require further symmetry within the triangulated macro-cells; see Example 5.1 for an illustration. This property will be required in the proof of Proposition 5.2 below. In particular, not every periodic coefficient may meet this requirement. Also, generating such a triangulation requires knowledge about the length of the period.

Example 5.1. Figure 1 displays a periodic coefficient and a matching triangulation.

We remark that the error estimate (3.18) and Proposition 4.5 hold in this case as well. Due to the periodic boundary conditions, the auxiliary solution u^{reg} utilized in the proof of Proposition 4.5 has the smoothness $u^{reg} \in H^2(\Omega)$ so that those estimates




FIG. 1. Periodic coefficients with respect to a square grid and triangulations: nonmatching (left) and matching (right).

are valid with $s = 1$. In the periodic setting, further properties of $A_H^{(\ell)}$ can be derived. First, it is not difficult to prove that the coefficient $A_H^{(\ell)}$ is globally constant. The following result states that, in the idealized case $\ell = \infty$, the coefficient $A_H^{(\ell)}$ is even independent of the mesh-size H and coincides with the classical homogenization limit, where for any $j = 1, \dots, d$, the corrector $\hat{q}_j \in H_{\#}^1(\Omega)/\mathbb{R}$ is the solution to

$$(5.1) \quad \operatorname{div} A(\nabla \hat{q}_j - e_j) = 0 \text{ in } \Omega \text{ with periodic boundary conditions.}$$

PROPOSITION 5.2. *Let A be periodic and let \mathcal{T}_H be uniform and aligned with the periodicity pattern of A and let V, W be spaces with periodic boundary conditions. Then, for any $T \in \mathcal{T}_H$, the idealized coefficient $A_H^{(\infty)}|_T$ coincides with the homogenized coefficient from the classical homogenization theory. In particular, $A_H^{(\infty)}$ is globally constant and independent of H .*

Proof. Let $T \in \mathcal{T}_H$ and $j, k \in \{1, \dots, d\}$. The definitions of $A_H^{(\infty)}|_T$ and $\mathcal{K}^{(\infty)}$ lead to

$$(5.2) \quad \begin{aligned} \int_T A_{jk} \, dx - (A_H^{(\infty)}|_T)_{jk} &= |T|^{-1} \sum_{K \in \mathcal{T}_H} \int_K e_j \cdot (A \nabla q_{T,k}) \, dx \\ &= |T|^{-1} \int_{\Omega} e_j \cdot (A \nabla q_{T,k}) \, dx. \end{aligned}$$

The sum over all element correctors defined by $q_k := \sum_{T \in \mathcal{T}_H} q_{T,k}$ solves

$$(5.3) \quad a(w, q_k) = (\nabla w, A e_k)_{L^2(\Omega)} \quad \text{for all } w \in W.$$

The definitions of $q_{T,k}$ and q_k and the symmetry of A lead to

$$(5.4) \quad \begin{aligned} |T|^{-1} \int_{\Omega} e_j \cdot (A \nabla q_{T,k}) \, dx &= |T|^{-1} \int_{\Omega} \nabla q_j \cdot (A \nabla q_{T,k}) \, dx \\ &= \int_T e_k \cdot (A \nabla q_j) \, dx. \end{aligned}$$

Let $v \in V$. We have $(v - I_H v) \in W$ and therefore by (5.3) that

$$\begin{aligned} \int_{\Omega} \nabla v \cdot (A(\nabla q_j - e_j)) \, dx &= \int_{\Omega} (\nabla I_H v) \cdot (A(\nabla q_j - e_j)) \, dx \\ &= \sum_{K \in \mathcal{T}_H} \int_K (\nabla I_H v) \, dx \cdot \int_K A(\nabla q_j - e_j) \, dx, \end{aligned}$$

where for the last identity it was used that $\nabla I_H v$ is constant on each element. By periodicity we have that $\int_K A(\nabla q_j - e_j) dx = \int_\Omega A(\nabla q_j - e_j) dx$ for any $K \in \mathcal{T}_H$. Therefore, for all $v \in V$,

$$\int_\Omega \nabla v \cdot (A(\nabla q_j - e_j)) dx = \int_\Omega (\nabla I_H v) dx \cdot \int_\Omega A(\nabla q_j - e_j) dx = 0$$

due to the periodic boundary conditions of $I_H v$. Hence, the difference $\nabla q_j - e_j$ satisfies (5.1). This is the corrector problem from classical homogenization theory and, thus, the proof is concluded by the above formulae (5.2)–(5.4). Indeed, by symmetry of A ,

$$(A_H^{(\infty)})_{j,k}|_T = \int_T A_{jk} dx - \int_T e_k \cdot (A \nabla q_j) dx = \int_T (e_j - \nabla q_j) \cdot A e_k dx. \quad \square$$

Remark 5.3. For Dirichlet boundary conditions, the method is different from the classical periodic homogenization as it takes the boundary conditions into account.

The remaining parts of this section are devoted to an L^2 error estimate for the classical homogenization limit. Let the coefficient $A = A_\varepsilon$ be periodic, oscillating on the scale ε . Let H be the observation scale represented by the mesh-size of the finite element mesh. We couple H to ε so that the ratio H/ε is constant. Recall from Proposition 5.2 that the idealized coefficient $A_H^{(\infty)} = A_0$ for a constant coefficient A_0 that is independent of H . It is known (see, e.g., [4]) that, in the present case of a symmetric coefficient, A_0 satisfies the bounds (4.2). Denote, for any ε , by $u_\varepsilon \in V$ the solution to

$$(5.5) \quad \int_\Omega \nabla u_\varepsilon \cdot (A_\varepsilon \nabla v) dx = F(v) \quad \text{for all } v \in V.$$

Denote by $u_0 \in V$ the solution to

$$(5.6) \quad \int_\Omega \nabla u_0 \cdot (A_0 \nabla v) dx = F(v) \quad \text{for all } v \in V.$$

In periodic homogenization theory, the function u_0 is called the homogenized solution. The aim is to estimate $\|u_0 - u_\varepsilon\|_{L^2(\Omega)}$ in terms of ε . The following perturbation result is required.

LEMMA 5.4 (perturbed coefficient). *Let H and ε be coupled so that H/ε is constant. Let the localization parameter ℓ be chosen of order $\ell \approx |\log H|$. Then,*

$$\|A_H^{(\infty)} - A_H^{(\ell)}\|_{L^\infty(\Omega)} \lesssim H.$$

There exist $\varepsilon_0 > 0$ and $0 < \alpha' \leq \beta' < \infty$ such that for all $\varepsilon \leq \varepsilon_0$

$$\alpha' |\xi|^2 \leq \xi \cdot (A_H^{(\ell)}(x)\xi) \leq \beta' |\xi|^2$$

for all $\xi \in \mathbb{R}^d$ and almost all $x \in \Omega$.

Proof. Remark 4.1 shows that $A_H^{(\infty)}$ and $A_H^{(\ell)}$ are given on any $T \in \mathcal{T}_H$ through

$$(A_H^{(\infty)})_{j,k}|_T = |T|^{-1} \int_\Omega e_j \cdot (A(\chi_T e_k - \nabla q_{T,k}))$$

and

$$(A_H^{(\ell)})_{j,k}|_T = |T|^{-1} \int_{\Omega_T} e_j \cdot (A(\chi_T e_k - \nabla q_{T,k}^{(\ell)}))$$

for any $j, k \in \{1, \dots, d\}$. Thus,

$$\begin{aligned} |(A_H^{(\infty)})_{j,k}|_T - (A_H^{(\ell)})_{j,k}|_T &= |T|^{-1} \int_{\Omega} e_j \cdot (A(\nabla(q_{T,k} - q_{T,k}^{(\ell)}))) \\ &\leq |T|^{-1} \|A^{1/2} e_j\|_{L^2(\Omega)} \|A^{1/2} \nabla(q_{T,k} - q_{T,k}^{(\ell)})\|_{L^2(\Omega)} \\ &\leq |T|^{-1} \|A^{1/2} \nabla(q_{T,k} - q_{T,k}^{(\ell)})\|_{L^2(\Omega)}. \end{aligned}$$

It is shown in [21, proof of Corollary 4.11] that

$$\|A^{1/2} \nabla(q_{T,k} - q_{T,k}^{(\ell)})\|_{L^2(\Omega)} \lesssim \exp(-c\ell) |T|^{1/2}.$$

In conclusion, the choice $\ell \approx |\log H|$ implies the first stated estimate. The second stated result follows from a perturbation argument because it is known [4] that $A_H^{(\infty)} = A_0$ satisfies (2.2). \square

The following result recovers the classical homogenization limit $u_\varepsilon \rightarrow u_0$ strongly in L^2 as $\varepsilon \rightarrow 0$. In particular, it quantifies the convergence speed and states that for $f \in L^2(\Omega)$ an almost linear rate is achieved.

PROPOSITION 5.5 (quantified homogenization limit). *Let Ω be convex, let $u_\varepsilon \in V$ solve (5.5), and let $u_0 \in V$ solve (5.6). For any $\varepsilon \leq \varepsilon_0$ (for ε_0 from Lemma 5.4) we have*

$$\|u_\varepsilon - u_0\|_{L^2(\Omega)} \lesssim \varepsilon |\log \varepsilon|^{1+d/2} \|f\|_{L^2(\Omega)}.$$

Proof. As before, we couple H and ε such that H/ε is constant. We denote by $u_H^{(\ell)} \in V_H$ the solution to (3.10), by $\tilde{u}_H^{(\ell)} \in V_H$ the solution to (4.1), and by $\tilde{u}_H^{(\infty)} \in V_H$ the solution to (4.1) with the choice $\ell = \infty$, where in all problems A is replaced by A_ε . Note that Lemma 5.4 implies stability of the discrete system (4.1) and thereby unique existence of $\tilde{u}_H^{(\ell)}$. We employ the triangle inequality to split the error as follows:

$$\begin{aligned} (5.7) \quad \|u_\varepsilon - u_0\|_{L^2(\Omega)} &\lesssim \|u_\varepsilon - u_H^{(\ell)}\|_{L^2(\Omega)} + \|u_H^{(\ell)} - \tilde{u}_H^{(\ell)}\|_{L^2(\Omega)} \\ &\quad + \|\tilde{u}_H^{(\ell)} - \tilde{u}_H^{(\infty)}\|_{L^2(\Omega)} + \|\tilde{u}_H^{(\infty)} - u_0\|_{L^2(\Omega)}. \end{aligned}$$

Estimate (3.18) allows us to bound the first term on the right-hand side of (5.7) as

$$\|u_\varepsilon - u_H^{(\ell)}\|_{L^2(\Omega)} \lesssim \varepsilon \|f\|_{L^2(\Omega)}.$$

The second term on the right-hand side of (5.7) was bounded in Proposition 4.5. With the Friedrichs inequality the result reads

$$\|u_H^{(\ell)} - \tilde{u}_H^{(\ell)}\|_{L^2(\Omega)} \lesssim \varepsilon |\log \varepsilon|^{1+d/2} \|f\|_{L^2(\Omega)},$$

where it was used that $\eta(A_H^{(\ell)}) = 0$ because $A_H^{(\ell)}$ is spatially constant. In order to bound the third term on the right-hand side of (5.7) we use the stability of the discrete problems and the perturbation result of Lemma 5.4 to deduce

$$\|\tilde{u}_H^{(\ell)} - \tilde{u}_H^{(\infty)}\|_{L^2(\Omega)} \lesssim \|A_H^{(\infty)} - A_H^{(\ell)}\|_{L^\infty(\Omega)} \|f\|_{L^2(\Omega)} \lesssim \varepsilon \|f\|_{L^2(\Omega)}.$$

For the fourth term on the right-hand side of (5.7) it is enough to note that $\tilde{u}_H^{(\infty)}$ is the Galerkin approximation of u_0 in V_H , which satisfies

$$\|\tilde{u}_H^{(\infty)} - u_0\|_{L^2(\Omega)} \lesssim \varepsilon^2 \|f\|_{L^2(\Omega)}$$

on convex domains. The combination of the foregoing estimates concludes the proof. \square

6. Numerical illustration. In this section, we present numerical experiments on the unit square domain $\Omega = (0, 1)^2$ with homogeneous Dirichlet boundary conditions. We consider the following worst-case error (referred to as the L^2 error) as error measure:

$$\sup_{f \in L^2(\Omega) \setminus \{0\}} \frac{\|u(f) - u_{\text{discrete}}(f)\|_{L^2(\Omega)}}{\|f\|_{L^2(\Omega)}},$$

where $u(f)$ is the exact solution to (2.3) with right-hand side f and $u_{\text{discrete}}(f)$ a discrete approximation (standard FEM or local effective coefficient or quasi-local effective coefficient or L^2 -best approximation). The error quantity is approximated by solving an eigenvalue problem on the reference mesh.

6.1. First experiment: Convergence rates. Consider the scalar coefficient A ,

$$A(x_1, x_2) = \left(\frac{11}{2} + \sin\left(\frac{2\pi}{\varepsilon_1} x_1\right) \sin\left(\frac{2\pi}{\varepsilon_1} x_2\right) + 4 \sin\left(\frac{2\pi}{\varepsilon_2} x_1\right) \sin\left(\frac{2\pi}{\varepsilon_2} x_2\right) \right)^{-1}$$

with $\varepsilon_1 = 2^{-3}$ and $\varepsilon_2 = 2^{-5}$. We consider a sequence of uniformly refined meshes of mesh-size $H = \sqrt{2} \times 2^{-1}, \dots, \sqrt{2} \times 2^{-6}$. The corrector problems are solved on a reference mesh of width $h = \sqrt{2} \times 2^{-9}$. The localization (or oversampling) parameter is chosen as $\ell = 2$. Figure 2 displays the coefficient A . The four components of the reconstructed coefficient $A_H^{(\ell)}$ for $H = \sqrt{2} \times 2^{-6}$ are displayed in Figure 3. Figure 4 compares the L^2 errors of the standard FEM, the FEM with the local effective coefficient, the method with the quasi-local effective coefficient, and the L^2 -best approximation in dependence of H . For comparison, also the error of the MSFEM from [13] is displayed. As expected, the error of the FEM is of order $\mathcal{O}(1)$ because the coefficient is not resolved by the mesh-size H . The error for the quasi-local effective coefficient is close to the best approximation. The local effective coefficient leads to comparable errors on coarse meshes. On the finest mesh, where the coefficient is almost resolved, the error deteriorates. This effect, referred to as the “resonance effect,” will be studied in the second numerical experiment. Table 1 lists the values of the




FIG. 2. The scalar coefficient A for the first experiment.




FIG. 3. Matrix entries of the reconstructed localized coefficient $(A_H^{(\ell)})$ in the first experiment for $H = \sqrt{2} \times 2^{-6}$.




FIG. 4. Convergence history under uniform mesh refinement.

estimator $\eta(A_H^{(\ell)})$ as well as the bounds α_H and β_H on $(A_H^{(\ell)})$. The estimator $\eta(A_H^{(\ell)})$ is small on the first meshes, which corresponds to an effective coefficient close to a constant. The estimator increases for the meshes approaching the resonance regime. The values of the coefficient A range in the interval $[\alpha, \beta] = [0.096, 1.55]$. In this example, the discrete bounds α_H, β_H stay in this interval.

6.2. Second experiment: Resonance effects. In this experiment we investigate so-called resonance effects of our homogenization procedure. These effects occur because, unlike in section 5, in the present case we deal with Dirichlet boundary conditions as well as meshes that do not satisfy requirements in the spirit of Example 5.1. We consider a fixed mesh of width $H = \sqrt{2} \times 2^{-4}$ and the scalar coefficient

TABLE 1

Values of the estimator $\eta(A_H^{(\ell)})$ and the bounds α_H and β_H on A_H for the first experiment. The values of the coefficient A range in the interval $[\alpha, \beta] = [0.096, 1.55]$.

H	$\eta(A_H^{(\ell)})$	α_H	β_H
$\sqrt{2} \times 2^{-1}$	3.2108e-02	1.9223e-01	2.0786e-01
$\sqrt{2} \times 2^{-2}$	1.1267e-02	1.9568e-01	1.9954e-01
$\sqrt{2} \times 2^{-3}$	1.4765e-02	1.9579e-01	1.9986e-01
$\sqrt{2} \times 2^{-4}$	5.3952e-01	1.8323e-01	2.1992e-01
$\sqrt{2} \times 2^{-5}$	1.7199e+00	1.6909e-01	2.3257e-01
$\sqrt{2} \times 2^{-6}$	1.5538e+01	1.4070e-01	3.0277e-01




FIG. 5. Resonance effect: normalized (by L^2 -best error) errors of FEM, local effective model and quasi-local effective model; and values of the estimator $\eta(A_H^{(\ell)})$.

$$A(x_1, x_2) = \left(5 + 4 \sin\left(\frac{2\pi}{\varepsilon}x_1\right) \sin\left(\frac{2\pi}{\varepsilon}x_2\right) \right)^{-1}$$

for a sequence of parameters $\varepsilon = 2^0, 2^{-1}, \dots, 2^{-6}$. The coefficient $(A_H^{(\ell)})$ was computed with the same reference mesh and the same oversampling parameter as in the first experiment. Figure 5 displays the L^2 errors normalized by the L^2 error of the L^2 -best approximation. On the third mesh, where H and ε have the same order of magnitude, the local effective coefficient leads to a larger error compared to the coarser meshes (where the coefficient is resolved by H) and finer meshes, where H is much coarser than ε and the effective coefficient is close to a constant. We observe that the values of the estimator $\eta(A_H^{(\ell)})$ are large in the resonance regime where also the error of the method the local effective coefficient is large. For smaller values of ε , the values of $\eta(A_H^{(\ell)})$ are close to zero, which indicates that the homogenization criterion from Remark 4.6 is satisfied; cf. also Remark 4.10.

Acknowledgments. The main parts of this paper were written at the Institut für Numerische Simulation (Bonn). The authors thank the Hausdorff Institute for Mathematics in Bonn for the kind hospitality of during the trimester program on multiscale problems.

REFERENCES

- [1] A. ABDULLE, W. E. B. ENGQUIST, AND E. VANDEN-EIJNDEN, *The heterogeneous multiscale method*, Acta Numer., 21 (2012), pp. 1–87.
- [2] R. A. ADAMS, *Sobolev Spaces*, Pure Appl. Math., Academic Press, New York, 1975.
- [3] G. ALLAIRE, *Homogenization and two-scale convergence*, SIAM J. Math. Anal., 23 (1992), pp. 1482–1518.
- [4] G. ALLAIRE, *Mathematical approaches and methods*, in Homogenization and Porous Media, U. Hornung, ed., Interdiscip. Appl. Math. 6, Springer, New York, 1997, pp. 225–250, 259–275.
- [5] I. BABUSKA AND R. LIPTON, *Optimal local approximation spaces for generalized finite element methods with application to multiscale problems*, Multiscale Model. Simul., 9 (2011), pp. 373–406.
- [6] A. BENSOUSSAN, J.-L. LIONS, AND G. PAPANICOLAOU, *Asymptotic Analysis for Periodic Structures*, North-Holland, Amsterdam, 1978.
- [7] J. BERGH AND J. LÖFSTRÖM, *Interpolation Spaces. An Introduction*, Grundlehren Math. Wiss. 223, Springer-Verlag, Berlin, 1976.
- [8] L. BERLYAND AND H. OWHADI, *Flux norm approach to finite dimensional homogenization approximations with non-separated scales and high contrast*, Arch. Ration. Mech. Anal., 198 (2010), pp. 677–721.
- [9] M. DAUGE, *Elliptic Boundary Value Problems on Corner Domains. Smoothness and Asymptotics of Solutions*, Springer-Verlag, Berlin, 1988.
- [10] E. DE GIORGI, *Sulla convergenza di alcune successioni d'integrali del tipo dell'area*, Rend. Mat., 8 (1975), pp. 277–294.
- [11] W. E AND B. ENGQUIST, *The heterogeneous multiscale methods*, Commun. Math. Sci., 1 (2003), pp. 87–132.
- [12] Y. EFENDIEV, J. GALVIS, AND T. Y. HOU, *Generalized multiscale finite element methods (GMs-FEM)*, J. Comput. Phys., 251 (2013), pp. 116–135.
- [13] Y. EFENDIEV AND T. Y. HOU, *Multiscale Finite Element Methods*, Surv. Tutor. Appl. Math. Sci. 4, Springer, New York, 2009.
- [14] D. GALLISTL AND D. PETERSEIM, *Stable multiscale Petrov-Galerkin finite element method for high frequency acoustic scattering*, Comput. Methods Appl. Mech. Eng., 295 (2015), pp. 1–17.
- [15] D. GALLISTL AND D. PETERSEIM, *Numerical Stochastic Homogenization by Quasiloal Effective Diffusion Tensors*, arXiv:1702.08858, 2017.
- [16] L. GRASEDYCK, I. GREFF, AND S. SAUTER, *The AL basis for the solution of elliptic problems in heterogeneous media*, Multiscale Model. Simul., 10 (2012), pp. 245–258.
- [17] I. GREFF AND W. HACKBUSCH, *Numerical method for elliptic multiscale problems*, J. Numer. Math., 16 (2008), pp. 119–138.
- [18] P. GRISVARD, *Elliptic Problems in Nonsmooth Domains*, Monogr. Stud. Math. 24, Pitman, Boston, MA, 1985.
- [19] W. HACKBUSCH, *Hierarchical Matrices: Algorithms and Analysis*, Springer-Verlag, Berlin, 2015.
- [20] P. HENNING, P. MORGENSTERN, AND D. PETERSEIM, *Multiscale partition of unity*, in Meshfree Methods for Partial Differential Equations VII, M. Griebel and M. A. Schweitzer, eds., Lect. Notes Comput. Sci. Eng. 100, Springer, New York, 2015, pp. 185–204.
- [21] P. HENNING AND D. PETERSEIM, *Oversampling for the multiscale finite element method*, Multiscale Model. Simul., 11 (2013), pp. 1149–1175.
- [22] T. Y. HOU AND X.-H. WU, *A multiscale finite element method for elliptic problems in composite materials and porous media*, J. Comput. Phys., 134 (1997), pp. 169–189.
- [23] T. J. R. HUGHES, G. R. FELIÓO, L. MAZZEI, AND J.-B. QUINCY, *The variational multiscale method—a paradigm for computational mechanics*, Comput. Methods Appl. Mech. Engrg., 166 (1998), pp. 3–24.
- [24] T. J. R. HUGHES AND G. SANGALLI, *Variational multiscale analysis: The fine-scale Green's function, projection, optimization, localization, and stabilized methods*, SIAM J. Numer. Anal., 45 (2007), pp. 539–557.
- [25] R. KORNUBER, D. PETERSEIM, AND H. YSERENTANT, *An Analysis of a Class of Variational Multiscale Methods Based on Subspace Decomposition*, arXiv:1608.04081, 2016.
- [26] R. KORNUBER AND H. YSERENTANT, *Numerical homogenization of elliptic multiscale problems by subspace decomposition*, Multiscale Model. Simul., 14 (2016), pp. 1017–1036.
- [27] A. MÅLQVIST, *Multiscale methods for elliptic problems*, Multiscale Model. Simul., 9 (2011), pp. 1064–1086.

- [28] A. MÅLQVIST AND D. PETERSEIM, *Localization of elliptic multiscale problems*, Math. Comp., 83 (2014), pp. 2583–2603.
- [29] M. ANA-MARIA AND S. CHRISTOPH, *Two-scale fem for homogenization problems*, ESAIM Math. Model. Numer. Anal., 36 (2002), pp. 537–572.
- [30] J. M. MELENK, *hp-Finite Element Methods for Singular Perturbations*, Lecture Notes in Math. 1796, Springer-Verlag, Berlin, 2002.
- [31] F. MURAT AND L. TARTAR, *H-convergence*, in Proceedings of Séminaire d'Analyse Fonctionnelle et Numérique de l'Université d'Alger, 1978.
- [32] G. NGUETSENG, *A general convergence result for a functional related to the theory of homogenization*, SIAM J. Math. Anal., 20 (1989), pp. 608–623.
- [33] H. OWHADI, *Multigrid with rough coefficients and multiresolution operator decomposition from hierarchical information games*, SIAM Rev., 59 (2017), pp. 99–149.
- [34] H. OWHADI AND L. ZHANG, *Metric-based upscaling*, Comm. Pure Appl. Math., 60 (2007), pp. 675–723.
- [35] H. OWHADI, L. ZHANG, AND L. BERLYAND, *Polyharmonic homogenization, rough polyharmonic splines and sparse super-localization*, ESAIM Math. Model. Numer. Anal., 48 (2014), pp. 517–552.
- [36] D. PETERSEIM, *Variational multiscale stabilization and the exponential decay of fine-scale correctors*, in Building Bridges: Connections and Challenges in Modern Approaches to Numerical Partial Differential Equations, G. R. Barrenechea, F. Brezzi, A. Cangiani, and E. H. Georgoulis, eds., Lect. Notes Comput. Sci. Eng. 114, Springer, New York, 2016, pp. 341–367.
- [37] S. SPAGNOLO, *Sulla convergenza di soluzioni di equazioni paraboliche ed ellittiche*, Ann. Sc. Norm. Super. Pisa, 22 (1968), pp. 571–597.
- [38] M. WEYMUTH AND S. SAUTER, *An adaptive local (AL) basis for elliptic problems with complicated discontinuous coefficients*, PAMM, 15 (2015), pp. 605–606.