

# LORENTZ SPACES WITH VARIABLE EXPONENTS

HENNING KEMPKA AND JAN VYBÍRAL

ABSTRACT. We introduce Lorentz spaces  $L_{p(\cdot),q}(\mathbb{R}^n)$  and  $L_{p(\cdot),q(\cdot)}(\mathbb{R}^n)$  with variable exponents. We prove several basic properties of these spaces including embeddings and the identity  $L_{p(\cdot),p(\cdot)}(\mathbb{R}^n) = L_{p(\cdot)}(\mathbb{R}^n)$ . We also show that these spaces arise through real interpolation between  $L_{p(\cdot)}(\mathbb{R}^n)$  and  $L_\infty(\mathbb{R}^n)$ . Furthermore, we answer in a negative way the question posed in [11] whether the Marcinkiewicz interpolation theorem holds in the frame of Lebesgue spaces with variable integrability.

## 1. INTRODUCTION

Lorentz spaces were introduced in [23, 24] as a generalization of classical Lebesgue spaces and have become a standard tool in mathematical analysis, cf. [3, 6, 4, 15]. For an introduction to Lorentz spaces we refer e.g. to [31, Chapter V], [4, Chapter 4] or [15, Chapter 1].

One of the main ingredients of the theory of Lorentz spaces is the celebrated Marcinkiewicz interpolation theorem, which states that under certain conditions one can deduce the strong boundedness of a sublinear operator  $T$  on the interpolation spaces provided that the operator is weakly bounded at the endpoints of the interpolation pair. This approach was used for example in the classical book of Stein [30] to prove the boundedness of the Hardy-Littlewood maximal operator on  $L_p(\mathbb{R}^n)$  for  $1 < p \leq \infty$ .

Another classical topic we shall touch in our work are the Lebesgue spaces  $L_{p(\cdot)}(\mathbb{R}^n)$  of variable integrability. The study of this class of function spaces goes back to Orlicz [27]. After the survey paper of Kováčik and Rákosník [22], there has been an enormous interest in these spaces (and in Sobolev spaces  $W_{p(\cdot)}^1(\Omega)$  built on  $L_{p(\cdot)}(\Omega)$ ) especially in connection with the application in modeling of electrorheological fluids [28]. Moreover, the spaces  $L_{p(\cdot)}(\mathbb{R}^n)$  possess interesting applications in the theory of PDE's, variational calculus, financial mathematics and image processing. A recent overview of this vastly growing field is given in [13]. A fundamental breakthrough concerning spaces of variable integrability was the observation that, under certain regularity assumptions on  $p(\cdot)$ , the Hardy-Littlewood maximal operator is also bounded on  $L_{p(\cdot)}(\mathbb{R}^n)$ , see [10]. This result has been generalized to wider classes of exponents  $p(\cdot)$  in [9], [26] and [12]. Unfortunately, it turned out that the standard proof of Stein [30] for spaces with constant indices breaks down and completely different methods had to be used to achieve this result, see [13].

---

The first author acknowledges the financial support by the DFG grants HA 2794/5-1 and KE 1847/1-1.

The second author acknowledges the support by the DFG Research Center MATHEON "Mathematics for key technologies" in Berlin.

The main aim of this paper is to return to this topic and to study the validity of the Marcinkiewicz interpolation theorem in the frame of Lebesgue spaces with variable integrability. For this reason, we first explore the possibility of extending the definition of Lorentz spaces to the setting of variable integrability exponents. We show, that there is really a natural way to define the Lorentz spaces  $L_{p(\cdot),q(\cdot)}(\mathbb{R}^n)$ , which extends the scale of Lebesgue spaces with variable exponents, i.e.  $L_{p(\cdot),p(\cdot)}(\mathbb{R}^n) = L_{p(\cdot)}(\mathbb{R}^n)$  for  $p(\cdot) = q(\cdot)$ . Later on, we study the interpolation properties of this new scale of spaces. Especially, we show in Theorem 8 that

$$(L_{p(\cdot)}(\mathbb{R}^n), L_\infty(\mathbb{R}^n))_{\theta,q} = L_{\tilde{p}(\cdot),q}(\mathbb{R}^n)$$

for  $0 < \theta < 1$  and

$$\frac{1}{\tilde{p}(\cdot)} = \frac{1-\theta}{p(\cdot)}.$$

Finally, we discuss the validity of the Marcinkiewicz interpolation theorem in the context of this new scale of function spaces, an open question posed in [11]. It turns out that the answer is negative and we provide a detailed counterexample to this conjecture.

The structure of the paper is as follows. Section 2 collects classical definitions of Lorentz spaces with constant indices and of Lebesgue spaces with variable integrability. Furthermore, the definition of Lorentz spaces with variable integrability is given. After collecting some basic properties of this new scale of function spaces in Section 3, we study the real interpolation properties of this scale in Section 4. Section 5 is devoted to Marcinkiewicz interpolation and contains the counterexample to [11, Question 2.8]. Finally, Section 6 collects some possible research directions and open problems.

At the end of this introduction we would like to mention that another definition of Lorentz spaces  $\mathcal{L}^{p(\cdot),q(\cdot)}(\mathbb{R}^n)$  with variable exponents was recently given in [14] with [19] and [18] as forerunners. Their definition works with non-increasing rearrangement and two variable exponents  $p(\cdot), q(\cdot) : [0, \infty) \rightarrow [1, \infty]$ . Due to this effect, the important and natural identity  $\mathcal{L}^{p(\cdot),p(\cdot)}(\mathbb{R}^n) = L_{p(\cdot)}(\mathbb{R}^n)$  does not hold in this scale of variable Lorentz spaces. This is a consequence of the definition of  $L_{p(\cdot)}(\mathbb{R}^n)$  where the variable exponent  $p(\cdot)$  is defined on  $\mathbb{R}^n$  and not on  $[0, \infty)$ . We return to this topic in detail in Remark 2.

## 2. OLD AND NEW DEFINITIONS

In this section we collect the very well known definitions of classical Lorentz spaces (Section 2.2) and Lebesgue spaces of variable exponents (Section 2.1). Finally, in Section 2.3, we provide the definition of Lorentz spaces with variable exponents. For simplicity, we start in Definition 2 with the more intuitive spaces  $L_{p(\cdot),q}(\mathbb{R}^n)$ . The Lorentz spaces  $L_{p(\cdot),q(\cdot)}(\mathbb{R}^n)$  with both exponents variable are introduced shortly after in Definition 3.

**2.1. Lebesgue spaces with variable exponents.** Let us now recall the definition of the variable Lebesgue spaces  $L_{p(\cdot)}(\mathbb{R}^n)$ . A measurable function  $p : \mathbb{R}^n \rightarrow (0, \infty]$  is called a variable exponent function if it is bounded away from zero. For a set  $A \subset \mathbb{R}^n$  we denote  $p_A^+ = \operatorname{ess\,sup}_{x \in A} p(x)$  and  $p_A^- = \operatorname{ess\,inf}_{x \in A} p(x)$ ; we use the abbreviations  $p^+ = p_{\mathbb{R}^n}^+$  and  $p^- = p_{\mathbb{R}^n}^-$ . The variable exponent Lebesgue space

$L_{p(\cdot)}(\mathbb{R}^n)$  consists of all measurable functions  $f$  such that there exist an  $\lambda > 0$  such that the modular

$$\varrho_{L_{p(\cdot)}(\mathbb{R}^n)}(f/\lambda) = \int_{\mathbb{R}^n} \varphi_{p(x)} \left( \frac{|f(x)|}{\lambda} \right) dx$$

is finite, where

$$(1) \quad \varphi_p(t) = \begin{cases} t^p & \text{if } p \in (0, \infty), \\ 0 & \text{if } p = \infty \text{ and } t \leq 1, \\ \infty & \text{if } p = \infty \text{ and } t > 1. \end{cases}$$

This definition is nowadays standard and was used also in [1, Section 2.2] and [13, Definition 3.2.1].

If we define  $\mathbb{R}_\infty^n = \{x \in \mathbb{R}^n : p(x) = \infty\}$  and  $\mathbb{R}_0^n = \mathbb{R}^n \setminus \mathbb{R}_\infty^n$ , then the Luxemburg norm of a function  $f \in L_{p(\cdot)}(\mathbb{R}^n)$  is given by

$$\begin{aligned} \|f\|_{L_{p(\cdot)}(\mathbb{R}^n)} &= \inf\{\lambda > 0 : \varrho_{L_{p(\cdot)}(\mathbb{R}^n)}(f/\lambda) \leq 1\} \\ &= \inf \left\{ \lambda > 0 : \int_{\mathbb{R}_0^n} \left( \frac{|f(x)|}{\lambda} \right)^{p(x)} dx \leq 1 \text{ and } |f(x)| \leq \lambda \text{ for a.e. } x \in \mathbb{R}_\infty^n \right\}. \end{aligned}$$

It constitutes a norm if  $p(\cdot) \geq 1$ , but it is always a quasi-norm if at least  $p^- > 0$ . Furthermore, the spaces  $L_{p(\cdot)}(\mathbb{R}^n)$  are complete, hence they are (quasi-) Banach spaces if  $p^- > 0$ , see [22] for details and further properties. We denote the class of all measurable functions  $p : \mathbb{R}^n \rightarrow (0, \infty]$  such that  $p^- > 0$  by  $\mathcal{P}(\mathbb{R}^n)$  and the corresponding modular is denoted by  $\varrho_{p(\cdot)}$  instead of  $\varrho_{L_{p(\cdot)}(\mathbb{R}^n)}$ .

**2.2. Classical Lorentz spaces.** Next, we recall the definition of classical Lorentz spaces as it can be found in [4] or [15]. This definition makes use of the so-called *non-increasing rearrangement*  $f^*$  of a function  $f$ . For a measurable function  $f : \mathbb{R}^n \rightarrow \mathbb{C}$ , we define first the *distribution function*  $\mu_f : [0, \infty) \rightarrow [0, \infty]$  by

$$\mu_f(s) = \mu\{x \in \mathbb{R}^n : |f(x)| > s\}, \quad s \geq 0,$$

where  $\mu$  denotes the Lebesgue measure on  $\mathbb{R}^n$ .

The distribution function  $\mu_f$  provides information about the size of  $f$  but not about the local behavior of  $f$ . The (generalized) inverse function to the distribution function is called *non-increasing rearrangement*  $f^* : [0, \infty) \rightarrow [0, \infty]$  and is defined by

$$f^*(t) = \inf\{s > 0 : \mu_f(s) \leq t\}.$$

Equipped with these tools, we are now ready to give the definition of the classical Lorentz spaces with constant indices.

**Definition 1.** Given a measurable function  $f$  on  $\mathbb{R}^n$  and real parameters  $0 < p, q \leq \infty$ , we define

$$\|f\|_{L_{p,q}(\mathbb{R}^n)} = \begin{cases} \left( \int_0^\infty \left( t^{1/p} f^*(t) \right)^q \frac{dt}{t} \right)^{1/q}, & \text{if } q < \infty \\ \sup_{t>0} t^{1/p} f^*(t), & \text{if } q = \infty. \end{cases}$$

The space of all measurable  $f : \mathbb{R}^n \rightarrow \mathbb{C}$  with  $\|f\|_{L_{p,q}(\mathbb{R}^n)} < \infty$  is denoted by  $L_{p,q}(\mathbb{R}^n)$ . The spaces are complete and they are normable for  $1 < p < \infty$  and  $1 \leq q \leq \infty$ , see [15, Theorem 1.4.11 and Exercise 1.4.3].

The use of non-increasing rearrangement makes it rather difficult to extend Definition 1 to variable exponents  $p(\cdot), q(\cdot) : \mathbb{R}^n \rightarrow (0, \infty]$ . It is very well known that the spaces  $L_{p(\cdot)}(\mathbb{R}^n)$  are not translation invariant (see Proposition 3.6.1 in [13]) and therefore the membership of  $f$  in  $L_{p(\cdot)}(\mathbb{R}^n)$  cannot be characterized by any condition on  $f^*$  only.

To avoid this obstacle, we look for an equivalent characterization of Lorentz spaces  $L_{p,q}(\mathbb{R}^n)$  which does not make use of the notion of non-increasing rearrangement. Therefore we calculate for  $p, q < \infty$  using Fubini's theorem and the substitution  $s^{p/q} := t$  (cf. Proposition 1.4.9 in [15])

$$\begin{aligned}
\|f\|_{L_{p,q}(\mathbb{R}^n)} &= \left( \int_0^\infty \left( t^{1/p} f^*(t) \right)^q \frac{dt}{t} \right)^{1/q} = \left( \int_0^\infty \frac{p}{q} f^*(s^{p/q})^q ds \right)^{1/q} \\
&= \left( \frac{p}{q} \right)^{1/q} \left( \int_0^\infty \mu\{s \geq 0 : f^*(s^{p/q})^q > t\} dt \right)^{1/q} \\
&= \left( \frac{p}{q} \right)^{1/q} \left( \int_0^\infty \mu\{s \geq 0 : f^*(s^{p/q}) > t^{1/q}\} dt \right)^{1/q} \\
&= p^{1/q} \left( \int_0^\infty \lambda^q \mu\{s \geq 0 : f^*(s^{p/q}) > \lambda\} \frac{d\lambda}{\lambda} \right)^{1/q} \\
&= p^{1/q} \left( \int_0^\infty \lambda^q \mu\{s \geq 0 : f^*(s) > \lambda\}^{q/p} \frac{d\lambda}{\lambda} \right)^{1/q} \\
(2) \quad &= p^{1/q} \left( \int_0^\infty \lambda^q \|\chi_{\{x \in \mathbb{R}^n : |f(x)| > \lambda\}}\|_{L_p(\mathbb{R}^n)}^q \frac{d\lambda}{\lambda} \right)^{1/q}.
\end{aligned}$$

Here,  $\chi_{\{x \in \mathbb{R}^n : |f(x)| > \lambda\}}$  stands for the characteristic function of the set  $\{x \in \mathbb{R}^n : |f(x)| > \lambda\}$ . If no confusion is possible, this will also be denoted by  $\chi_{\{|f| > \lambda\}}$ .

The equation (2) can be discretized and we derive

$$(3) \quad \|f\|_{L_{p,q}(\mathbb{R}^n)} \sim p^{1/q} \left( \sum_{k=-\infty}^{\infty} 2^{kq} \|\chi_{\{x \in \mathbb{R}^n : |f(x)| > 2^k\}}\|_{L_p(\mathbb{R}^n)}^q \right)^{1/q}.$$

**2.3. Lorentz spaces with variable exponents.** The expression (2) for the norm can be generalized quite easily to variable exponent  $p(\cdot)$  with  $q$  constant. Surprisingly enough, even  $q$  can be considered variable when we use the spaces  $\ell_{q(\cdot)}(L_{p(\cdot)})$  of Almeida and Hästö [1] and the discretized equation (3). Furthermore we do not destroy the local properties of the function  $f$ , since it gets not rearranged and the exponents map from  $\mathbb{R}^n$ .

**Definition 2.** Let  $p \in \mathcal{P}(\mathbb{R}^n)$  be a variable exponent with range  $0 < p^- \leq p^+ \leq \infty$  and let  $0 < q \leq \infty$ . Then  $L_{p(\cdot),q}(\mathbb{R}^n)$  is the collection of all measurable functions  $f : \mathbb{R}^n \rightarrow \mathbb{C}$  such that

$$(4) \quad \|f\|_{L_{p(\cdot),q}(\mathbb{R}^n)} = \begin{cases} \left( \int_0^\infty \lambda^q \|\chi_{\{x \in \mathbb{R}^n : |f(x)| > \lambda\}}\|_{L_{p(\cdot)}(\mathbb{R}^n)}^q \frac{d\lambda}{\lambda} \right)^{1/q}, & \text{if } q < \infty \\ \sup_{\lambda > 0} \lambda \|\chi_{\{x \in \mathbb{R}^n : |f(x)| > \lambda\}}\|_{L_{p(\cdot)}(\mathbb{R}^n)}, & \text{if } q = \infty \end{cases}$$

is finite.

Using the  $\ell_{q(\cdot)}(L_{p(\cdot)})$  spaces introduced recently in [1], we may even consider the situation, where also  $q$  is variable. Let us recall their approach. For a sequence  $(f_\nu)_{\nu \in \mathbb{Z}}$  we define the modular

$$\varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}((f_\nu)_\nu) = \sum_{\nu \in \mathbb{Z}} \inf \left\{ \lambda_\nu > 0 : \varrho_{p(\cdot)} \left( \frac{f_\nu}{\lambda_\nu^{1/q(\cdot)}} \right) \leq 1 \right\},$$

with the convention  $\lambda^{1/\infty} = 1$ . If  $q^+ < \infty$  or if  $q(\cdot) \leq p(\cdot)$  we can replace this by a simpler expression

$$(5) \quad \varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}((f_\nu)_\nu) = \sum_{\nu \in \mathbb{Z}} \left\| \varphi_{q(\cdot)}(|f_\nu(\cdot)|) \Big| L_{\frac{p(\cdot)}{q(\cdot)}}(\mathbb{R}^n) \right\|,$$

which is much more intuitive. Here  $\varphi_q(t)$  equals basically  $t^q$ , see (1). The norm in these spaces gets defined as usual as the Luxemburg norm

$$\| (f_\nu)_\nu | \ell_{q(\cdot)}(L_{p(\cdot)}) \| = \inf \{ \mu > 0 : \varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})} (f_\nu/\mu) \leq 1 \}.$$

Up to now, it is not completely clear under which conditions on  $p(\cdot)$  and  $q(\cdot)$  the expression above becomes a norm. It was shown in [1] that it always constitutes a quasi-norm if  $p^-, q^- > 0$ . Further it is known (see [20]) that it is a norm if either  $\frac{1}{p(\cdot)} + \frac{1}{q(\cdot)} \leq 1$  holds pointwise for all  $x \in \mathbb{R}^n$  or if  $1 \leq q(\cdot) \leq p(\cdot) \leq \infty$  holds pointwise. Also in this work there is given an example where  $\min(p(\cdot), q(\cdot)) \geq 1$  but the triangle inequality does not hold. Let us mention that it is an open question if there exists an equivalent norm on  $\ell_{q(\cdot)}(L_{p(\cdot)})$  whenever  $\min(p(\cdot), q(\cdot)) \geq 1$ . Nevertheless, since our exponents are between  $(0, \infty]$  we generally work with quasi-norms and there are no obstacles with that.

We use now the modular  $\varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}$  and (3) to define the variable Lorentz spaces  $L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$ .

**Definition 3.** Let  $p, q \in \mathcal{P}(\mathbb{R}^n)$  be two variable exponents with range  $0 < p^- \leq p^+ \leq \infty$  and  $0 < q^- \leq q^+ \leq \infty$ . Then  $L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$  is the collection of all measurable functions  $f : \mathbb{R}^n \rightarrow \mathbb{C}$  such that

$$(6) \quad \| f | L_{p(\cdot), q(\cdot)}(\mathbb{R}^n) \| = \inf \left\{ \lambda > 0 : \varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})} \left( 2^k \chi_{\{x \in \mathbb{R}^n : |f(x)/\lambda| > 2^k\}} \right) \leq 1 \right\} < \infty.$$

Before we discuss the properties of these new function spaces we derive an equivalent expression for  $\| f | L_{p(\cdot), q(\cdot)}(\mathbb{R}^n) \|$ .

**Lemma 4.** Let  $p, q \in \mathcal{P}(\mathbb{R}^n)$  be two variable exponents with range  $0 < p^- \leq p^+ \leq \infty$  and  $0 < q^- \leq q^+ \leq \infty$ . Then

$$(7) \quad \| f | L_{p(\cdot), q(\cdot)}(\mathbb{R}^n) \| \approx \left\| \left( 2^k \chi_{\{x \in \mathbb{R}^n : |f(x)| > 2^k\}} \right)_{k=-\infty}^\infty \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\|.$$

*Proof.* If  $\lambda = 2^j$  for some  $j \in \mathbb{Z}$ , we obtain

$$\varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})} \left( 2^k \chi_{\{x \in \mathbb{R}^n : |f(x)/\lambda| > 2^k\}} \right) = \varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})} \left( \frac{2^k \chi_{\{x \in \mathbb{R}^n : |f(x)| > 2^k\}}}{\lambda} \right).$$

The rest of the proof then follows by simple monotonicity arguments.  $\square$

*Remark 1.* The somehow more complicated definition of the quasi-norm in Definition 3 was necessary. Only the expression in (6) is homogeneous; i.e.

$$\| \lambda f | L_{p(\cdot), q(\cdot)}(\mathbb{R}^n) \| = |\lambda| \cdot \| f | L_{p(\cdot), q(\cdot)}(\mathbb{R}^n) \| \quad \text{for all } \lambda \in \mathbb{R}.$$

Easy examples show that the right-hand side of (7) fails to have this property. Nevertheless due to Lemma 4, both expressions are equivalent and therefore define the same spaces  $L_{p(\cdot),q(\cdot)}(\mathbb{R}^n)$ . In majority of our considerations, we shall work with the somehow simpler expression (7).

If  $q(\cdot) = q$  is a constant function, then the Proposition 3.3 in [1] shows that  $\ell_{q(\cdot)}(L_{p(\cdot)})$  is really an iterated space, i.e.

$$\|(f_\nu)_{\nu \in \mathbb{Z}}|_{\ell_q(L_{p(\cdot)})}\| = \left( \sum_{\nu \in \mathbb{Z}} \|f_\nu|_{L_{p(\cdot)}(\mathbb{R}^n)}\|^q \right)^{1/q}$$

(with an appropriate modification if  $q = \infty$ ). By (3) and (7), we obtain that Definitions 2 and 3 are equivalent.

Moreover, we observe by (2) and (3) that for constant functions  $p(x) = p$  and  $q(x) = q$  we get an equivalent norm for the usual Lorentz spaces  $L_{p,q}(\mathbb{R}^n)$ , whenever  $p, q < \infty$ . A similar calculation justifies this fact also if  $p < q = \infty$ . If  $p = \infty$ , then the usual Lorentz spaces  $L_{\infty,q}(\mathbb{R}^n)$  defined by Definition 1 consist only of the zero function whenever  $0 < q < \infty$ , see Section 1.4.2 in [15]. It is easy to see, that Definition 2 applied to  $p(\cdot) = \infty$  and  $q < \infty$  gives  $L_{\infty,q}(\mathbb{R}^n) = L_\infty(\mathbb{R}^n)$ . Nevertheless, we will show that  $L_{p(\cdot),p(\cdot)}(\mathbb{R}^n) = L_{p(\cdot)}(\mathbb{R}^n)$  and therefore the case  $p = q = \infty$  is also included for  $p = \infty$ .

Summarizing, our spaces  $L_{p(\cdot),q(\cdot)}(\mathbb{R}^n)$  are equivalent to the usual Lorentz spaces  $L_{p,q}(\mathbb{R}^n)$  if  $p(\cdot) = p$  and  $q(\cdot) = q$  are constant functions. The only exception is the case if  $p = \infty$  and  $0 < q < \infty$ , see Theorem 7.

*Remark 2.* Another approach to generalize this definition to variable exponents was given in [14], with forerunners [19] and [18]. They introduced the spaces  $\mathcal{L}^{p(\cdot),q(\cdot)}(\Omega)$  by the corresponding quasi-norm

$$\|f|_{\mathcal{L}^{p(\cdot),q(\cdot)}(\Omega)}\| = \|t^{\frac{1}{p(\cdot)} - \frac{1}{q(\cdot)}} f^*(t)|_{L_{q(\cdot)}(0,|\Omega|)}\|,$$

where  $\Omega \subset \mathbb{R}^n$  is a measurable set,  $|\Omega|$  is its Lebesgue measure and  $p, q : (0, |\Omega|) \rightarrow (0, \infty)$  are variable exponents. The spaces  $\mathcal{L}^{p(\cdot),q(\cdot)}(\Omega)$  coincide with usual Lorentz spaces  $L_{p,q}(\Omega)$  if  $p(\cdot) = p$  and  $q(\cdot) = q$  are constant. On the other hand, in this scale there is no hope for the identity  $\mathcal{L}^{p(\cdot),p(\cdot)}(\Omega) = L_{p(\cdot)}(\Omega)$  to hold, since in the definition of the Lebesgue spaces the variable exponent  $p(\cdot)$  is defined on whole  $\Omega$ .

### 3. BASIC PROPERTIES

In this section, we prove several basic properties of the new scale of function spaces.

**Theorem 5.** *Let  $p, q \in \mathcal{P}(\mathbb{R}^n)$ . Then  $L_{p(\cdot),q(\cdot)}(\mathbb{R}^n)$  are quasi-Banach spaces.*

*Proof.* To prove that (6) defines a quasi-norm, we only have to show the quasi-triangle inequality. We use the estimate

$$\begin{aligned} \{x \in \mathbb{R}^n : |f(x) + g(x)| > 2^k\} &\subset \{x \in \mathbb{R}^n : |f(x)| + |g(x)| > 2^k\} \\ &\subset \{x \in \mathbb{R}^n : |f(x)| > 2^{k-1}\} \cup \{x \in \mathbb{R}^n : |g(x)| > 2^{k-1}\} \end{aligned}$$

to obtain

$$\chi_{\{x \in \mathbb{R}^n : |f(x) + g(x)| > 2^k\}} \leq \chi_{\{x \in \mathbb{R}^n : |f(x)| > 2^{k-1}\}} + \chi_{\{x \in \mathbb{R}^n : |g(x)| > 2^{k-1}\}}.$$

This in turn implies

$$\begin{aligned}
\|f + g\|_{L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)} &\approx \left\| \left( 2^k \chi_{\{x \in \mathbb{R}^n: |f(x) + g(x)| > 2^k\}} \right)_{k=-\infty}^{\infty} \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\| \\
&\leq \left\| \left( 2^k \chi_{\{x \in \mathbb{R}^n: |f(x)| > 2^{k-1}\}} \right)_{k=-\infty}^{\infty} + \left( 2^k \chi_{\{x \in \mathbb{R}^n: |g(x)| > 2^{k-1}\}} \right)_{k=-\infty}^{\infty} \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\| \\
&\leq c \left\{ \left\| \left( 2^k \chi_{\{x \in \mathbb{R}^n: |f(x)| > 2^{k-1}\}} \right)_{k=-\infty}^{\infty} \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\| + \left\| \left( 2^k \chi_{\{x \in \mathbb{R}^n: |g(x)| > 2^{k-1}\}} \right)_{k=-\infty}^{\infty} \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\| \right\} \\
&\lesssim \|f\|_{L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)} + \|g\|_{L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)},
\end{aligned}$$

where  $c$  is the constant from the quasi-triangle inequality of  $\ell_{q(\cdot)}(L_{p(\cdot)})$ .

To show that the spaces  $L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$  are complete we take a Cauchy sequence  $(f_l)_{l \in \mathbb{N}} \subset L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$ . We chose a subsequence (which we denote by  $(f_l)_{l \in \mathbb{N}}$  again) with

$$\|f_{l+1} - f_l\|_{L_{p(\cdot), q(\cdot)}} \leq \frac{1}{2^{2l}}, \quad l \in \mathbb{N}.$$

For notational reasons, we put  $f_0 = 0$ . We consider the function

$$g(t) := \sum_{l=0}^{\infty} |f_{l+1}(t) - f_l(t)|.$$

As  $\chi_{\{g > \lambda\}}(x) \leq \sum_{l=0}^{\infty} \chi_{\{|f_{l+1} - f_l| > \lambda/2^{l+1}\}}(x)$ , we obtain

$$\begin{aligned}
\|\chi_{\{g > \lambda\}}\|_{L_{p(\cdot)}(\mathbb{R}^n)}^r &\leq \sum_{l=0}^{\infty} \|\chi_{\{|f_{l+1} - f_l| > \lambda/2^{l+1}\}}\|_{L_{p(\cdot)}(\mathbb{R}^n)}^r \\
&\leq \sum_{l=0}^{\infty} \frac{2^{(l+1)r}}{\lambda^r} \cdot \|f_{l+1} - f_l\|_{L_{p(\cdot), \infty}}^r \lesssim \sum_{l=0}^{\infty} \frac{2^{(l+1)r}}{\lambda^r} 2^{-2lr}
\end{aligned}$$

where  $r = \min(p^-, 1)$  and we have used the embedding  $L_{p(\cdot), q(\cdot)}(\mathbb{R}^n) \hookrightarrow L_{p(\cdot), \infty}(\mathbb{R}^n)$ , see Theorem 7. As the last sum converges, we get  $\|\chi_{\{g > \lambda\}}\|_{L_{p(\cdot)}(\mathbb{R}^n)} \rightarrow 0$  for  $\lambda \rightarrow \infty$  and  $g$  is finite almost everywhere. Therefore, the series

$$f(x) = \sum_{l=0}^{\infty} f_{l+1}(x) - f_l(x) \quad \text{and} \quad \tilde{f}(x) = \sum_{l=1}^{\infty} f_{l+1}(x) - f_l(x) = f(x) - f_1(x), \quad x \in \mathbb{R}^n$$

converge also almost everywhere.

It remains to show that  $f \in L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$  and  $f_l \rightarrow f$  in  $L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$ .

The estimate  $2^k \chi_{\{|\tilde{f}| > 2^k\}} \leq \sum_{l=1}^{\infty} 2^k \chi_{\{|f_{l+1} - f_l| > 2^{k-l}\}}$  implies that

$$\begin{aligned}
\|2^k \chi_{\{|\tilde{f}| > 2^k\}}\|_{\ell_{q(\cdot)}(L_{p(\cdot)})}^{\varrho} &\lesssim \sum_{l=1}^{\infty} \|2^k \chi_{\{|f_{l+1} - f_l| > 2^{k-l}\}}\|_{\ell_{q(\cdot)}(L_{p(\cdot)})}^{\varrho} \\
&= \sum_{l=1}^{\infty} 2^{l\varrho} \|2^{k-l} \chi_{\{|f_{l+1} - f_l| > 2^{k-l}\}}\|_{\ell_{q(\cdot)}(L_{p(\cdot)})}^{\varrho} \lesssim \sum_{l=1}^{\infty} 2^{l\varrho} \cdot 2^{-2l\varrho} < \infty,
\end{aligned}$$

where  $\varrho > 0$  is chosen small enough, cf. [1, Theorem 3.8].

Therefore,  $\tilde{f} \in L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$  and  $f = \tilde{f} + f_1 \in L_{p(\cdot), q(\cdot)}(\mathbb{R}^n)$ .

Finally, for  $l \in \mathbb{N}$  fixed, we consider

$$f - f_l = \sum_{m=l}^{\infty} (f_{m+1} - f_m).$$

The estimate  $\chi_{\{|f-f_l|>2^k\}} \leq \sum_{m=l}^{\infty} \chi_{\{|f_{m+1}-f_m|>2^{k-(m-l+1)}\}}$  implies the convergence  $\|f-f_l\|_{L_{p(\cdot),q(\cdot)}(\mathbb{R}^n)} \rightarrow 0$  for  $l \rightarrow \infty$  in a similar manner as above.  $\square$

We continue with a theorem showing that the scale of variable Lorentz spaces includes the scale of Lebesgue spaces with variable exponent. We would like to emphasize, that the identity  $L_{p(\cdot),p(\cdot)}(\mathbb{R}^n) = L_{p(\cdot)}(\mathbb{R}^n)$  does hold without any restrictions on  $p(\cdot)$  with  $0 < p^- \leq p^+ \leq \infty$ .

**Theorem 6.** *If  $p \in \mathcal{P}(\mathbb{R}^n)$ , then it holds  $L_{p(\cdot),p(\cdot)}(\mathbb{R}^n) = L_{p(\cdot)}(\mathbb{R}^n)$ .*

*Proof.* We want to show

$$(8) \quad \varrho_{p(\cdot)}(f/2) \leq \varrho_{\varrho_{p(\cdot)}(L_{p(\cdot)})} \left( (2^k \chi_{\{x \in \mathbb{R}^n: |f(x)| > 2^k\}})_{k=-\infty}^{\infty} \right) \leq \varrho_{p(\cdot)}(cf),$$

where  $c = (1 - 2^{-p^-})^{-1/p^-}$ . From the inequalities above we conclude easily

$$\frac{1}{2} \|f\|_{L_{p(\cdot)}(\mathbb{R}^n)} \lesssim \|f\|_{L_{p(\cdot),p(\cdot)}(\mathbb{R}^n)} \lesssim c \|f\|_{L_{p(\cdot)}(\mathbb{R}^n)},$$

which proves the theorem. First this proof only treats the case  $|\mathbb{R}_{\infty}^n| = |\{x \in \mathbb{R}^n : p(x) = \infty\}| = 0$ . At the end we comment on the case  $|\mathbb{R}_{\infty}^n| > 0$ . Since  $p(\cdot) = q(\cdot)$  we can use the easy expression (5) for the modular and then the first inequality in (8) follows from

$$\begin{aligned} \varrho_{\varrho_{p(\cdot)}(L_{p(\cdot)})} \left( (2^k \chi_{\{|f|>2^k\}})_{k=-\infty}^{\infty} \right) &= \sum_{k=-\infty}^{\infty} \left\| |2^k \chi_{\{|f|>2^k\}}|^{p(x)} \right\|_{L_1(\mathbb{R}^n)} \\ &= \int_{\mathbb{R}^n} \sum_{\{k \in \mathbb{Z}: 2^k < |f(x)|\}} 2^{kp(x)} dx. \end{aligned}$$

For fixed  $x \in \mathbb{R}^n$  with  $|f(x)| > 0$  we choose the unique  $k_x \in \mathbb{Z}$  with  $2^{k_x p(x)} < |f(x)|^{p(x)} \leq 2^{(k_x+1)p(x)}$  and obtain

$$(9) \quad \sum_{\{k \in \mathbb{Z}: 2^k < |f(x)|\}} 2^{kp(x)} = \sum_{k=-\infty}^{k_x} \left(2^{-p(x)}\right)^{-k} = 2^{k_x p(x)} \frac{1}{1 - 2^{-p(x)}}.$$

Using  $1 \leq \frac{1}{1 - 2^{-p(x)}}$  we get from (9)

$$\begin{aligned} \varrho_{\varrho_{p(\cdot)}(L_{p(\cdot)})} \left( (2^k \chi_{\{|f(x)|>2^k\}})_{k=-\infty}^{\infty} \right) &= \int_{\mathbb{R}^n} \sum_{\{k \in \mathbb{Z}: 2^k < |f(x)|\}} 2^{kp(x)} dx \\ &\geq \int_{\mathbb{R}^n} 2^{k_x p(x)} dx = \int_{\mathbb{R}^n} 2^{(k_x+1)p(x)} 2^{-p(x)} dx \\ &\geq \int_{\mathbb{R}^n} \frac{1}{2} |f(x)|^{p(x)} dx = \varrho_{p(\cdot)}(f/2). \end{aligned}$$



The converse inequality uses again (9) with  $\frac{1}{1-2^{-p(x)}} \leq \frac{1}{1-2^{-p^-}}$  and follows in a similar way

$$\begin{aligned} \varrho_{\ell_{p(\cdot)}(L_{p(\cdot)})} \left( (2^k \chi_{\{|f(x)| > 2^k\}})_{k=-\infty}^{\infty} \right) &= \int_{\mathbb{R}^n} \sum_{\{k \in \mathbb{Z}: 2^k < |f(x)|\}} 2^{kp(x)} dx \\ &= \int_{\mathbb{R}^n} 2^{k_x p(x)} \frac{1}{1-2^{-p(x)}} dx \leq \int_{\mathbb{R}^n} 2^{k_x p(x)} \frac{1}{1-2^{-p^-}} dx \\ &\leq \int_{\mathbb{R}^n} |f(x)|^{p(x)} \left( \frac{1}{1-2^{-p^-}} \right)^{\frac{p(x)}{p^-}} dx = \varrho_{p(\cdot)}(cf), \end{aligned}$$

with  $c = (1-2^{-p^-})^{-1/p^-}$ .

Now, we come back to the case, where  $|\mathbb{R}_{\infty}^n| > 0$ . First, we split our function  $f = f_0 + f_{\infty} := f \cdot \chi_{\mathbb{R}^n_0} + f \cdot \chi_{\mathbb{R}^n_{\infty}}$ . Then we use the considerations above and

$$\|f_0|L_{p(\cdot)}(\mathbb{R}^n)\| + \|f_{\infty}|L_{p(\cdot)}(\mathbb{R}^n)\| \leq 2 \|f_0 + f_{\infty}|L_{p(\cdot)}(\mathbb{R}^n)\|,$$

see [13, Remark 3.2.3], and

$$\|f_0|L_{p(\cdot),p(\cdot)}(\mathbb{R}^n)\| + \|f_{\infty}|L_{p(\cdot),p(\cdot)}(\mathbb{R}^n)\| \leq 2 \|f_0 + f_{\infty}|L_{p(\cdot),p(\cdot)}(\mathbb{R}^n)\|,$$

which is implied by

$$\begin{aligned} \varrho_{\ell_{p(\cdot)}(L_{p(\cdot)})} \left( (2^k \chi_{\{|f_0+f_{\infty}| > 2^k\}})_{k=-\infty}^{\infty} \right) &= \varrho_{\ell_{p(\cdot)}(L_{p(\cdot)})} \left( (2^k \chi_{\{|f_0| > 2^k\}})_{k=-\infty}^{\infty} \right) \\ &\quad + \varrho_{\ell_{p(\cdot)}(L_{p(\cdot)})} \left( (2^k \chi_{\{|f_{\infty}| > 2^k\}})_{k=-\infty}^{\infty} \right). \end{aligned}$$

□

Next we show that the new scale of function spaces satisfies some elementary embeddings, which are very well know in the case of constant exponents.

- Theorem 7.** (i) Let  $p, q_0, q_1 \in \mathcal{P}(\mathbb{R}^n)$  with  $q_0(\cdot) \leq q_1(\cdot)$  pointwise. Then  $L_{p(\cdot),q_0(\cdot)}(\mathbb{R}^n) \hookrightarrow L_{p(\cdot),q_1(\cdot)}(\mathbb{R}^n)$ .  
(ii) Let  $q \in \mathcal{P}(\mathbb{R}^n)$ . Then  $L_{\infty,q(\cdot)}(\mathbb{R}^n) = L_{\infty}(\mathbb{R}^n)$ .  
(iii) Let  $p_0, p_1, q_0, q_1 \in \mathcal{P}(\mathbb{R}^n)$  with  $p_0^+ < \infty$  and  $\alpha := (p_1/p_0)^- > 1$ . Then the inequality

$$(10) \quad \|f|L_{p_0(\cdot),q_0(\cdot)}\| \leq c \|f|L_{p_1(\cdot),q_1(\cdot)}\|$$

holds for all measurable  $f$  with  $\text{supp } f \subset [0, 1]^n$  with  $c$  independent of  $f$ .

- (iv) Let  $p_0, p_1, q \in \mathcal{P}(\mathbb{R}^n)$  with  $p_0(\cdot) \leq p_1(\cdot)$  pointwise. Then the inequality

$$(11) \quad \|f|L_{p_0(\cdot),q(\cdot)}\| \leq c \|f|L_{p_1(\cdot),q(\cdot)}\|$$

holds for all measurable  $f$  with  $\text{supp } f \subset [0, 1]^n$  with  $c$  independent of  $f$ .

*Proof.* The first statement follows from Lemma 4 and the embedding  $\ell_{q_0(\cdot)}(L_{p(\cdot)}) \hookrightarrow \ell_{q_1(\cdot)}(L_{p(\cdot)})$  which has been proven in [1].

To prove the second part of the theorem, it is enough to use the above embedding in the form

$$L_{\infty,q^-}(\mathbb{R}^n) \hookrightarrow L_{\infty,q(\cdot)}(\mathbb{R}^n) \hookrightarrow L_{\infty,\infty}(\mathbb{R}^n) = L_{\infty}(\mathbb{R}^n)$$

and the simple embedding  $L_{\infty}(\mathbb{R}^n) \hookrightarrow L_{\infty,q^-}(\mathbb{R}^n)$ , which follows directly from Definition 3 and Lemma 4.

The proof of the third statement is based on the following simple fact. For every  $A \subset \mathbb{R}^n$  with  $\mu(A) \leq 1$  the following inequality holds

$$(12) \quad \|\chi_A|L_{p_0(\cdot)}\| \leq \|\chi_A|L_{p_1(\cdot)}\|^\alpha,$$

where again  $\alpha = (p_1/p_0)^- > 1$ .

To show (10), it is enough to assume that  $q_1(\cdot) = \infty$  and

$$\|f|L_{p_1(\cdot),\infty}(\mathbb{R}^n)\| \approx \sup_{k \in \mathbb{Z}} 2^k \|\chi_{\{|f|>2^k\}}|L_{p_1(\cdot)}(\mathbb{R}^n)\| = 1.$$

Using (12), we obtain

$$\begin{aligned} \|f|L_{p_0(\cdot),q_0(\cdot)}\|^{q_0^-} &\lesssim \|f|L_{p_0(\cdot),q_0^-}\|^{q_0^-} \leq \sum_{k=-\infty}^{\infty} 2^{kq_0^-} \|\chi_{\{|f|>2^k\}}|L_{p_0(\cdot)}(\mathbb{R}^n)\|^{q_0^-} \\ &\leq \sum_{k=-\infty}^0 2^{kq_0^-} + \sum_{k=1}^{\infty} 2^{kq_0^-} \|\chi_{\{|f|>2^k\}}|L_{p_1(\cdot)}(\mathbb{R}^n)\|^{q_0^-} \\ &\leq c + \sum_{k=1}^{\infty} 2^{kq_0^-} (2^{-k} \|f|L_{p_1(\cdot),\infty}(\mathbb{R}^n)\|)^{q_0^-} \leq c' \end{aligned}$$

with an obvious modification if  $q_0^- = \infty$ . This justifies (10).

The proof of the fourth statement follows a similar pattern. We start with  $f$  such that

$$\|f|L_{p_1(\cdot),q(\cdot)}(\mathbb{R}^n)\| \approx \left\| (2^k \chi_{\{x \in \mathbb{R}^n: |f(x)|>2^k\}})_{k=-\infty}^{\infty} \Big| \ell_{q(\cdot)}(L_{p_1(\cdot)}) \right\| = 1.$$

We use again the splitting into two parts, namely with  $k \leq 0$  and  $k \geq 1$ , respectively. In the first case, we use the bounded support of  $f$  to obtain.

$$\begin{aligned} \|(2^k \chi_{\{|f|>2^k\}})_{k=-\infty}^0 | \ell_{q(\cdot)}(L_{p_0(\cdot)})\| &\leq \|(2^k \chi_{[0,1]^n})_{k=-\infty}^0 | \ell_{q(\cdot)}(L_{p_0(\cdot)})\| \\ &\lesssim \|(2^k \chi_{[0,1]^n})_{k=-\infty}^0 | \ell_{q^-}(L_{p_0(\cdot)})\| \lesssim 1 \end{aligned}$$

The second part with  $k \in \mathbb{N}$  may be estimated directly as

$$\|(2^k \chi_{\{|f|>2^k\}})_{k=1}^{\infty} | \ell_{q(\cdot)}(L_{p_0(\cdot)})\| \leq \|(2^k \chi_{\{|f|>2^k\}})_{k=1}^{\infty} | \ell_{q(\cdot)}(L_{p_1(\cdot)})\| \lesssim 1,$$

which finishes the proof of (11).  $\square$

*Remark 3.* The second part of this theorem is in contrast to [15, Section 1.4.2], where  $L_{\infty,q}(\mathbb{R}^n) = \{0\}$  is stated. But this is also not surprising since we did not take the extra factor  $p^{1/q}$  appearing in (2) and (3) into our Definitions 2 and 3 of our variable Lorentz spaces.

#### 4. INTERPOLATION

We stated already in the introduction that the main importance of Lorentz spaces lies in their connection with (real) interpolation theory. In this section, we shall explore the interpolation properties of Lorentz spaces with variable exponents. But before we come to this, we recall some basics of the interpolation theory, as they may be found for example in the classical monographs [5] and [32].

We shall touch only the two most important interpolation methods - the real interpolation and the complex interpolation. Complex interpolation of variable

exponent spaces has already been treated in [11]. It turned out that the expected result

$$[L_{p_0(\cdot)}(\mathbb{R}^n), L_{p_1(\cdot)}(\mathbb{R}^n)]_\theta = L_{p_\theta(\cdot)}(\mathbb{R}^n)$$

does hold for all  $0 < \theta < 1$  and all  $\frac{1}{p_\theta(\cdot)} = \frac{1-\theta}{p_0(\cdot)} + \frac{\theta}{p_1(\cdot)}$  with  $p_0^-, p_1^- \geq 1$ .

This complex interpolation result has been complemented in [21] by showing

$$[L_{p(\cdot)}(\mathbb{R}^n), BMO(\mathbb{R}^n)]_\theta = L_{p_\theta(\cdot)}(\mathbb{R}^n) \quad \text{and} \quad [L_{p(\cdot)}(\mathbb{R}^n), H_1(\mathbb{R}^n)]_\theta = L_{p_\theta(\cdot)}(\mathbb{R}^n)$$

under some regularity conditions on  $p(\cdot)$ .

We shall therefore concentrate on the real interpolation method (the so-called *K-method*). Let  $X_0$  and  $X_1$  be two (quasi-)Banach spaces, which are both embedded into a topological vector space  $Y$ . Then the spaces  $X_0 + X_1$  is defined as the set of all  $x \in Y$ , which may be written as  $x = x_0 + x_1$  with  $x_0 \in X_0$  and  $x_1 \in X_1$ .

For any  $x \in X_0 + X_1$  and any  $0 < t < \infty$ , the so-called *Peetre K-functional* is defined by

$$(13) \quad K(x, t, X_0, X_1) = \inf\{\|x_0\|_{X_0} + t\|x_1\|_{X_1} : x = x_0 + x_1, x_0 \in X_0, x_1 \in X_1\}.$$

If the spaces  $X_0$  and  $X_1$  are fixed and no confusion is possible, then we abbreviate this to  $K(x, t)$ . If  $0 < \theta < 1$  and  $0 < q \leq \infty$ , then the *real interpolation space*  $(X_0, X_1)_{\theta, q}$  is defined as the set of all  $x \in X_0 + X_1$ , such that

$$\|x\|_{(X_0, X_1)_{\theta, q}} = \begin{cases} \left( \int_0^\infty t^{-\theta q} K(x, t)^q \frac{dt}{t} \right)^{1/q}, & \text{if } q < \infty, \\ \sup_{t>0} t^{-\theta} K(x, t), & \text{if } q = \infty \end{cases}$$

is finite.

**Theorem 8.** *Let  $p \in \mathcal{P}(\mathbb{R}^n)$  with  $p^+ < \infty$ . Let  $0 < q \leq \infty$  and  $0 < \theta < 1$  and put*

$$\frac{1}{\tilde{p}(\cdot)} = \frac{1-\theta}{p(\cdot)}.$$

*Then*

$$(14) \quad (L_{p(\cdot)}(\mathbb{R}^n), L_\infty(\mathbb{R}^n))_{\theta, q} = L_{\tilde{p}(\cdot), q}(\mathbb{R}^n)$$

*in the sense of equivalent quasi-norms.*

*Proof. Step 1.* We shall show that

$$(15) \quad (L_{p(\cdot), \infty}(\mathbb{R}^n), L_\infty(\mathbb{R}^n))_{\theta, q} \hookrightarrow L_{\tilde{p}(\cdot), q}(\mathbb{R}^n),$$

i.e. that

$$(16) \quad \int_0^\infty \lambda^q \|\chi_{\{x \in \mathbb{R}^n : |f(x)| > \lambda\}}\|_{L_{\tilde{p}(\cdot)}(\mathbb{R}^n)}^q \frac{d\lambda}{\lambda} \lesssim \int_0^\infty t^{-\theta q} K(f, t)^q \frac{dt}{t}.$$

We shall use that

$$\begin{aligned} K(f, t) &= \inf\{\|f^0\|_{p(\cdot), \infty} + t\|f^1\|_\infty : f = f^0 + f^1\} \\ &= \inf_{\mu > 0} \{ \|(|f(x)| - \mu)_+\|_{p(\cdot), \infty} + t\|\min(|f(x)|, \mu)\|_\infty \} \\ &\geq \inf_{\mu > 0} \{ \mu \|\chi_{\{x \in \mathbb{R}^n : |f(x)| \geq 2\mu\}}\|_{p(\cdot)} + t\|\min(f(x), \mu)\|_\infty \} \end{aligned}$$

for every fixed  $t > 0$ .

We denote  $h(\lambda) = \|\chi_{\{x \in \mathbb{R}^n : |f(x)| \geq \lambda\}}\|_{p(\cdot)}$  for  $\lambda > 0$  and  $f_*(t) = \sup\{\lambda > 0 : h(\lambda) \geq t\}$  its generalized inverse function. Using the assumption  $p^+ < \infty$ , we

obtain that  $h(f_*(t)) \geq t$  for all  $t > 0$ . Then we choose  $\mu$  by  $\mu = f_*(t)/2$ . This leads to  $K(f, t) \geq f_*(t)h(f_*(t))/2 \geq tf_*(t)/2$ . The proof is then a consequence of the following two estimates on the left and right hand side of (16)

$$\begin{aligned} LHS(16) &= \int_0^\infty \lambda^q h(\lambda)^{(1-\theta)q} \frac{d\lambda}{\lambda} \approx \sum_{k=-\infty}^\infty \int_{\lambda: 2^k < h(\lambda) \leq 2^{k+1}} \lambda^q h(\lambda)^{(1-\theta)q} \frac{d\lambda}{\lambda} \\ &\lesssim \sum_{k=-\infty}^\infty 2^{k(1-\theta)q} \int_{\lambda: 2^k \leq h(\lambda)} \lambda^q \frac{d\lambda}{\lambda} \leq \sum_{k=-\infty}^\infty 2^{k(1-\theta)q} \int_0^{f_*(2^k)} \lambda^q \frac{d\lambda}{\lambda} \lesssim \sum_{k=-\infty}^\infty 2^{k(1-\theta)q} f_*(2^k)^q \end{aligned}$$

and

$$\begin{aligned} RHS(16) &\geq \sum_{k=-\infty}^\infty \int_{2^k}^{2^{k+1}} t^{(1-\theta)q} f_*(t)^q \frac{dt}{t} \gtrsim \sum_{k=-\infty}^\infty 2^{k(1-\theta)q} \int_{2^k}^{2^{k+1}} f_*(t)^q \frac{dt}{t} \\ &\gtrsim \sum_{k=-\infty}^\infty 2^{(k+1)(1-\theta)q} f_*(2^{k+1})^q = \sum_{k=-\infty}^\infty 2^{k(1-\theta)q} f_*(2^k)^q. \end{aligned}$$

If  $q = \infty$ , only notational modifications are necessary.

*Step 2.* Next, we prove that

$$(17) \quad L_{\bar{p}(\cdot), q}(\mathbb{R}^n) \hookrightarrow (L_{p(\cdot)}(\mathbb{R}^n), L_\infty(\mathbb{R}^n))_{\theta, q},$$

i.e.

$$(18) \quad \int_0^\infty t^{-\theta q} K(f, t)^q \frac{dt}{t} \lesssim \int_0^\infty \lambda^q \|\chi_{\{x \in \mathbb{R}^n: |f(x)| > \lambda\}}\|_{L_{\bar{p}(\cdot)}(\mathbb{R}^n)}^q \frac{d\lambda}{\lambda}.$$

We start again with a reformulation of  $K(f, t)$ .

$$\begin{aligned} K(f, t) &= \inf\{\|f^0\|_{p(\cdot)} + t\|f^1\|_\infty : f = f^0 + f^1\} \\ &= \inf_{\mu > 0} \{(\|f(x)\| - \mu)_+ \|_{p(\cdot)} + t\| \min(|f(x)|, \mu) \|_\infty\} \\ &\leq \inf_{\mu > 0} \{\|f \chi_{\{x \in \mathbb{R}^n: |f(x)| > \mu\}}\|_{p(\cdot)} + t\mu\} \\ &\lesssim \inf_{\mu > 0} \left\{ \left\| \sum_{j=0}^\infty 2^j \mu \chi_{\{x \in \mathbb{R}^n: |f(x)| > 2^j \mu\}} \right\|_{p(\cdot)} + t\mu \right\} \\ &\leq \inf_{\mu > 0} \left\{ \sum_{j=0}^\infty 2^j \mu \|\chi_{\{x \in \mathbb{R}^n: |f(x)| > 2^j \mu\}}\|_{p(\cdot)} + t\mu \right\} \\ &= \inf_{\mu > 0} \left\{ \sum_{j=0}^\infty 2^j \mu h(2^j \mu) + t\mu \right\}, \end{aligned}$$

where we denoted  $h(\lambda) := \|\chi_{\{x \in \mathbb{R}^n: |f(x)| > \lambda\}}\|_{p(\cdot)}$ . Let us remark, that we have assumed  $p^- \geq 1$  in the calculation above to be able to use the triangle inequality without additional powers. The modification in the case  $p^- < 1$  is straightforward and left to the reader.

For fixed  $t > 0$ , we choose  $\mu = \mu(t)$  by

$$\mu(t) := \inf\{\mu > 0 : \sum_{j=0}^\infty 2^j h(2^j \mu) \leq t\}.$$

As the function  $h$  is right-continuous, we obtain immediately  $\sum_{j=0}^{\infty} 2^j h(2^j \mu(t)) \leq t$ . We first estimate the right-hand side of (18) as

$$RHS(18) \gtrsim \sum_{k=-\infty}^{\infty} 2^{kq} h(2^k)^{(1-\theta)q}.$$

Furthermore, we discretize the left-hand side of (18) as

$$\begin{aligned} LHS(18) &\leq \int_0^{\infty} t^{-\theta q} t^q \mu(t)^q \frac{dt}{t} = \sum_{k=-\infty}^{\infty} 2^{kq} \int_{t:2^k < \mu(t) \leq 2^{k+1}} t^{(1-\theta)q} \frac{dt}{t} \\ &\leq \sum_{k=-\infty}^{\infty} 2^{kq} \int_{t:2^k < \mu(t)} t^{(1-\theta)q} \frac{dt}{t}. \end{aligned}$$

If  $\mu(t) > 2^k$ , we obtain

$$t \leq \sum_{j=0}^{\infty} 2^j h(2^{j+k}).$$

Therefore, we continue

$$LHS(18) \lesssim \sum_{k=-\infty}^{\infty} 2^{kq} \int_0^{\sum_{j=0}^{\infty} 2^j h(2^{j+k})} t^{(1-\theta)q} \frac{dt}{t} \lesssim \sum_{k=-\infty}^{\infty} 2^{kq} \left( \sum_{j=0}^{\infty} 2^j h(2^{j+k}) \right)^{(1-\theta)q}.$$

If  $(1-\theta)q \leq 1$ , we may write ( $l = j + k$ )

$$\begin{aligned} LHS(18) &\lesssim \sum_{k=-\infty}^{\infty} 2^{kq} \sum_{j=0}^{\infty} 2^{j(1-\theta)q} h(2^{j+k})^{(1-\theta)q} \\ &= \sum_{l=-\infty}^{\infty} \sum_{j=0}^{\infty} 2^{(l-j)q} 2^{j(1-\theta)q} h(2^l)^{(1-\theta)q} \\ &= \sum_{l=-\infty}^{\infty} 2^{lq} h(2^l)^{(1-\theta)q} \sum_{j=0}^{\infty} 2^{-j\theta q} \lesssim \sum_{l=-\infty}^{\infty} 2^{lq} h(2^l)^{(1-\theta)q}. \end{aligned}$$

If  $(1-\theta)q > 1$ , we use a similar approach combined with Hölder's inequality

$$\begin{aligned} LHS(18) &\lesssim \sum_{k=-\infty}^{\infty} 2^{kq} \sum_{j=0}^{\infty} 2^{j(1+\varepsilon)(1-\theta)q} h(2^{j+k})^{(1-\theta)q} \\ &= \sum_{l=-\infty}^{\infty} \sum_{j=0}^{\infty} 2^{(l-j)q} 2^{j(1+\varepsilon)(1-\theta)q} h(2^l)^{(1-\theta)q} \\ &= \sum_{l=-\infty}^{\infty} 2^{lq} h(2^l)^{(1-\theta)q} \sum_{j=0}^{\infty} 2^{jq(-1+(1+\varepsilon)(1-\theta))} \lesssim \sum_{l=-\infty}^{\infty} 2^{lq} h(2^l)^{(1-\theta)q}, \end{aligned}$$

where we have assumed that  $\varepsilon > 0$  was small enough to obtain  $-1 + (1+\varepsilon)(1-\theta) = -\theta + \varepsilon(1-\theta) < 0$ . This finishes the proof.  $\square$

*Remark 4.* (i) Let us point out that the assumption  $p^+ < \infty$  is forced mainly be the technique of generalized inverse functions used in the proof of Theorem 8. We leave it as an open problem if this assumption might be removed.

- (ii) As the proof of Theorem 8 works only with norms of characteristic functions, one can easily generalize its statement to

$$(L_{p(\cdot),q_0}(\mathbb{R}^n), L_\infty(\mathbb{R}^n))_{\theta,q} = L_{\bar{p}(\cdot),q}(\mathbb{R}^n)$$

for any  $0 < q_0 \leq \infty$ . And due to monotonicity, the result holds for  $q_0 \in \mathcal{P}(\mathbb{R}^n)$  as well.

## 5. MARCINKIEWICZ INTERPOLATION THEOREM

Let  $T$  be an operator defined on measurable functions on  $\mathbb{R}^n$ . We say, that  $T$  is sublinear, if

$$|T(f+g)(x)| \leq |Tf(x)| + |Tg(x)|$$

holds for (almost) every  $x \in \mathbb{R}^n$ . One of the most important tools in analysis of (sub-)linear operators is the Marcinkiewicz interpolation theorem. Let us recall its statement as it may be found for example in [15, Corollary 1.4.21].

**Theorem 9.** *Let  $\Omega \subset \mathbb{R}^n$  be a measurable set and let  $T$  be a sublinear operator, which maps  $L_{p_0}(\Omega)$  to  $L_{q_0,\infty}(\Omega)$  and  $L_{p_1}(\Omega)$  to  $L_{q_1,\infty}(\Omega)$ , where  $0 < p_0 \neq p_1 \leq \infty$  and  $0 < q_0 \neq q_1 \leq \infty$ . Let  $0 < \theta < 1$  and put*

$$\frac{1}{p} := \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \quad \frac{1}{q} := \frac{1-\theta}{q_0} + \frac{\theta}{q_1}.$$

If

$$(19) \quad p \leq q,$$

then  $T$  maps boundedly  $L_p(\Omega)$  into  $L_q(\Omega)$ .

One of the prominent applications of Marcinkiewicz interpolation theorem was given by Stein in his classical book [30]. The Hardy-Littlewood maximal operator is defined for every locally-integrable function  $f$  on  $\mathbb{R}^n$  by

$$Mf(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(x)| dx, \quad x \in \mathbb{R}^n,$$

where the supremum is taken over all balls in  $\mathbb{R}^n$  containing  $x$ . It is easy to see that  $M$  acts boundedly from  $L_\infty(\mathbb{R}^n)$  into  $L_\infty(\mathbb{R}^n)$ . Furthermore, one shows that  $M$  maps  $L_1(\mathbb{R}^n)$  into  $L_{1,\infty}(\mathbb{R}^n)$ . This two facts, combined with Theorem 9, lead immediately to the boundedness of  $M$  on  $L_p(\mathbb{R}^n)$  for every  $1 < p \leq \infty$ .

The study of the maximal operator in the frame of Lebesgue spaces with variable exponents attracted a lot of attention with the most important breakthroughs being achieved in [10, 8, 26, 12, 7]. It turned out, cf. [13, Theorem 4.3.8], that  $M$  is bounded on  $L_{p(\cdot)}(\mathbb{R}^n)$  if  $p^- > 1$  with the function  $1/p(\cdot)$  satisfying the so-called *log-Hölder continuity* conditions. Under the same regularity conditions on  $p$  it was proven in [7], that if  $p^- \geq 1$  the maximal operator maps  $L_{p(\cdot)}(\mathbb{R}^n)$  into  $L_{p(\cdot),\infty}(\mathbb{R}^n)$ . Quite naturally, this raises the question if the boundedness of  $M$  on Lebesgue spaces of variable integrability could be deduced from this weak-type estimate and some version of the Marcinkiewicz interpolation theorem.

In its abstract setting, the same question was already posed as an open problem in [11]. We recall their notation first. We say, that the sublinear operator  $T$  is of weak-type  $(\pi_0(\cdot), \pi_1(\cdot))$ , if

$$\lambda \|\chi_{\{x \in \mathbb{R}^n: |Tf(x)| > \lambda\}}\|_{\pi_1(\cdot)} \leq c \|f\|_{\pi_0(\cdot)}$$

holds for some  $c > 0$  and all  $f \in L_{\pi_0(\cdot)}(\mathbb{R}^n)$  and all  $\lambda > 0$ .

Let  $\pi_0(\cdot)$  and  $\pi_1(\cdot)$  be two variable exponents and let  $0 < \theta < 1$  be a real number. Then we put

$$\frac{1}{\pi_\theta(x)} := \frac{1-\theta}{\pi_0(x)} + \frac{\theta}{\pi_1(x)}.$$

The Question 2.8 from [11] sounds

**Question 2.8**([11], Marcinkiewicz Interpolation) Let  $T$  be a sublinear operator that is of weak type  $(\pi_0(\cdot), \pi_0(\cdot))$  and  $(\pi_1(\cdot), \pi_1(\cdot))$ . Is  $T$  then bounded from  $L_{\pi_\theta(\cdot)}(\mathbb{R}^n)$  to  $L_{\pi_\theta(\cdot)}(\mathbb{R}^n)$ ?

We shall prove that the answer to Question 2.8 is negative.

The basic idea of our construction is based on the observation that the condition (19) is necessary for the Marcinkiewicz interpolation theorem on usual  $L_p(\mathbb{R}^n)$  with constant exponent, cf. [16]. It means that there are  $0 < p_0 \neq p_1 \leq \infty$  and  $0 < q_0 \neq q_1 \leq \infty$  and  $0 < \theta < 1$  such that  $T$  is of weak type  $(p_0, q_0)$  and  $(p_1, q_1)$ ,  $p > q$  and  $T$  is not bounded from  $L_p[0, 1]$  to  $L_q[0, 1]$ .

Then we set

$$\tilde{T}f(x) := \begin{cases} T(\chi_{[0,1]}f)(x-1), & x \in [1, 2], \\ 0, & x \in [0, 1]. \end{cases}$$

We put

$$\pi_0(x) := \begin{cases} p_0, & x \in [0, 1], \\ q_0, & x \in [1, 2] \end{cases} \quad \text{and} \quad \pi_1(x) := \begin{cases} p_1, & x \in [0, 1], \\ q_1, & x \in [1, 2]. \end{cases}$$

We obtain that  $\tilde{T}$  is of weak type  $(\pi_0(\cdot), \pi_0(\cdot))$  and of weak type  $(\pi_1(\cdot), \pi_1(\cdot))$  but not of strong type  $(\pi_\theta(\cdot), \pi_\theta(\cdot))$ . Furthermore, a simple modification of this argument allows to construct a counterexample even for smooth parameters  $\pi_0(\cdot)$  and  $\pi_1(\cdot)$ . To make the presentation self-contained, we provide a simple construction of  $T$  with the properties mentioned above.

**5.1. Specific counterexample for  $T$ .** In this section, we shall provide more details on the above given construction. Especially, we shall construct an operator  $T$  which satisfies the assumptions from our counterexample. Following the work of Hunt [16, 17], we define for  $\alpha > 0$  the following Hardy type operator

$$(T_\alpha f)(x) = x^{-\alpha-1} \int_0^x f(t)dt, \quad 0 < x < 1.$$

We observe first that  $T_\alpha$  is linear and defined on all  $L_1(0, 1)$ . Using the estimate  $|\int_0^x f(t)dt| \leq \int_0^x f^*(t)dt$  and Hölder's inequality

$$\|T_{1/2}f\|_{1,\infty} \leq \sup_{0 < x < 1} x \cdot x^{-3/2} \int_0^x f^*(t)dt \leq \sup_{0 < x < 1} x^{-1/2} \left( \int_0^x (f^*(t))^2 dt \right)^{1/2} \cdot x^{1/2} = \|f\|_2,$$

we obtain also the weak-type estimate  $T_{1/2} : L_2(0, 1) \rightarrow L_{1,\infty}(0, 1)$ . Furthermore, the boundedness of  $T_{1/2}$  from  $L_\infty(0, 1)$  into  $L_{2,\infty}(0, 1)$  follows by

$$\|T_{1/2}f\|_{2,\infty} \leq \sup_{0 < x < 1} x^{1/2} \cdot x^{-3/2} \int_0^x f^*(t)dt = \sup_{0 < x < 1} x^{-1} \cdot \int_0^x f^*(t)dt \leq \|f\|_\infty.$$

On the other hand, it is easy to see, that  $T_{1/2}$  is not bounded from  $L_4(0, 1)$  into  $L_{4/3}(0, 1)$ . Just take  $f(t) = t^{-1/4} |\ln(t)|^{-1/4-\varepsilon} \chi_{[0,1/2]}(t) \in L_4(0, 1)$  for  $\varepsilon > 0$  and

calculate

$$\begin{aligned}
\|T_{1/2}f\|_{4/3} &= \left( \int_0^1 x^{-2} \left( \int_0^x f(t) dt \right)^{4/3} dx \right)^{3/4} \\
&\geq \left( \int_0^{1/2} x^{-2} \left( \int_0^x t^{-1/4} |\ln(t)|^{-1/4-\varepsilon} dt \right)^{4/3} dx \right)^{3/4} \\
&\approx \left( \int_0^{1/2} x^{-2} \left( x^{3/4} |\ln(x)|^{-1/4-\varepsilon} \right)^{4/3} dx \right)^{3/4} \\
&= \left( \int_0^{1/2} x^{-1} \cdot |\ln(x)|^{-1/3-\varepsilon \cdot 4/3} dx \right)^{3/4} = \infty
\end{aligned}$$

for  $0 < \varepsilon \leq 1/2$ .

## 6. OPEN PROBLEMS

Although we presented some basic properties of the scale of Lorentz spaces with variable exponents, many questions remained opened for further investigations.

The first obvious generalization is to treat Lorentz spaces  $L_{p(\cdot),q(\cdot)}(\Omega, \mu)$  on arbitrary measure spaces  $(\Omega, \mu)$  as it is usually done for Lorentz spaces with constant exponents, cf. [15]. We believe that the considerations above are also true in this case and we have studied mainly spaces on  $\mathbb{R}^n$  to simplify the notation.

It would be also highly desirable to obtain further results on real interpolation in this scale, complementing Theorem 8. Especially, it would be useful to have two variable exponent spaces as interpolation couple, i.e. to characterize  $(L_{p_0(\cdot)}(\mathbb{R}^n), L_{p_1(\cdot)}(\mathbb{R}^n))_{\theta, q}$ .

The use of  $\ell_{q(\cdot)}(L_{p(\cdot)})$  spaces suggests yet another interesting idea, namely to allow for interpolation with variable parameter  $q(\cdot)$ . This option was already noticed in the introduction of [1]. Unfortunately it turns out, and it is also mentioned in [2], that the real interpolation spaces with variable parameter  $q(\cdot)$  lack in general the interpolation property. On the other hand, it is possible, that under suitable conditions on the endpoint spaces, the real interpolation method with variable  $q(\cdot)$  works well and the interpolation property is restored again.

The last open problem is the starting point of this paper. If the Marcinkiewicz interpolation theorem would work in the scale of variable exponent spaces, we would have found a very elegant way to prove the boundedness of the Hardy-Littlewood maximal operator on  $L_{p(\cdot)}(\mathbb{R}^n)$ . Combining the weak estimate from [7]

$$M : L_{p_0(\cdot)}(\mathbb{R}^n) \rightarrow L_{p_0(\cdot), \infty}(\mathbb{R}^n)$$

with the trivial boundedness

$$M : L_\infty(\mathbb{R}^n) \rightarrow L_\infty(\mathbb{R}^n)$$

we could get the boundedness of  $M$  on  $L_{p(\cdot)}(\mathbb{R}^n)$ , with  $\frac{1}{p(\cdot)} = \frac{1-\theta}{p_0(\cdot)}$ . Unfortunately the previous counterexample tells us, that Marcinkiewicz does not work on variable  $L_{p(\cdot)}(\mathbb{R}^n)$  spaces. On the other hand, using Theorem 8, we may easily show that

$$M : L_{p(\cdot), q}(\mathbb{R}^n) \rightarrow L_{p(\cdot), q}(\mathbb{R}^n)$$

under the regularity assumptions on  $p(\cdot)$  as used in [7]. We would be therefore very much interested in additional conditions on the sublinear operator  $T$ , which would ensure the validity of Marcinkiewicz interpolation.



## REFERENCES

- [1] A. Almeida, P. Hästö, *Besov spaces with variable smoothness and integrability*, J. Funct. Anal. **258** no. 5 (2010), 1628–1655.
- [2] A. Almeida, P. Hästö, *Interpolation in variable exponent spaces*, preprint.
- [3] M. A. Ariño, B. Muckenhoupt, *Maximal functions on classical Lorentz spaces and Hardy's inequality with weights for nonincreasing functions*, Trans. Amer. Math. Soc. **320** no. 2 (1990), 727–735.
- [4] C. Bennett, R. Sharpley, *Interpolation of operators*, Academic Press, San Diego, 1988.
- [5] J. Bergh, J. Löfström, *Interpolation spaces. An introduction*, Springer, Berlin, 1976.
- [6] M. Carro, L. Pick, J. Soria, V. D. Stepanov, *On embeddings between classical Lorentz spaces*, Math. Inequal. Appl. **4** no. 3 (2001), 397–428.
- [7] D. Cruz-Uribe, L. Diening, A. Fiorenza, *A new proof of the boundedness of maximal operators on variable Lebesgue spaces*, Boll. Unione Mat. Ital. (9) **2** no. 1 (2009), 151–173.
- [8] D. Cruz-Uribe, A. Fiorenza, J. Martell, C. Pérez, *The boundedness of classical operators in variable  $L^p$  spaces*, Ann. Acad. Sci. Fenn. Math. **31** (2006), 239–264.
- [9] D. Cruz-Uribe, A. Fiorenza, C. J. Neugebauer, *The maximal function on variable  $L^p$  spaces*, Ann. Acad. Sci. Fenn. Math. **28** (2003), 223–238.
- [10] L. Diening, *Maximal function on generalized Lebesgue spaces  $L^{p(\cdot)}$* , Math. Inequal. Appl. **7** (2004), 245–253.
- [11] L. Diening, P. Hästö, A. Nekvinda, *Open problems in variable exponent Lebesgue and Sobolev spaces*, Proceedings FSDONA 2004, Academy of Sciences, Prague, 38–52.
- [12] L. Diening, P. Harjulehto, P. Hästö, Y. Mizuta, T. Shimomura, *Maximal functions in variable exponent spaces: limiting cases of the exponent*, Ann. Acad. Sci. Fenn. Math. **34** (2009), 503–522.
- [13] L. Diening, P. Harjulehto, P. Hästö, M. Růžička, *Lebesgue and Sobolev Spaces with Variable Exponents*, Lecture Notes in Mathematics **2017**, Springer, Berlin, 2011.
- [14] L. Ephremidze, V. Kokilashvili, S. Samko, *Fractional, maximal and singular operators in variable exponent Lorentz spaces*, Fract. Calc. Appl. Anal. **11** no. 4 (2008), 407–420.
- [15] L. Grafakos, *Classical Fourier analysis*, Second edition, Grad. Texts Math. **249**, Springer, Berlin, 2008.
- [16] R. A. Hunt, *An extension of the Marcinkiewicz interpolation theorem to Lorentz spaces*, Bull. Amer. Math. Soc. **70**, (1964) 803–807.
- [17] R. A. Hunt and G. Weiss, *The Marcinkiewicz interpolation theorem*, Proc. Amer. Math. Soc. **15**, (1964) 996–998.
- [18] D.M. Israfilov, V. Kokilashvili, N.P. Tuzkaya, *The classical integral operators in weighted Lorentz spaces with variable exponent*, IBSU Scientific Journal **1** issue 1, (2006), 171–178.
- [19] V. Kokilashvili, S. Samko, *Singular integrals and potentials in some Banach spaces with variable exponent*, J. Funct. Spaces and Appl. **1** (1), (2003), 45–59.
- [20] H. Kempka, J. Vybíral, *A note on the spaces of variable integrability and summability of Almeida and Hästö*, to appear in Proc. AMS.
- [21] T. Kopaliani, *Interpolation theorems for variable exponent Lebesgue spaces*, J. Funct. Anal. **257** (2009), 3541–3551.
- [22] O. Kováčik, J. Rákosník, *On spaces  $L^{p(x)}$  and  $W^{1,p(x)}$* , Czechoslovak Math. J. **41** (116) (1991), 592–618.
- [23] G. G. Lorentz, *Some new functional spaces*, Ann. of Math. (2) **51** (1950), 37–55.
- [24] G. G. Lorentz, *On the theory of spaces  $\Lambda$* , Pacific J. Math. **1** (1951), 411–429.
- [25] J. Musielak, *Orlicz spaces and modular spaces*, Lecture Notes in Mathematics **1034**, Springer, Berlin, 1983.
- [26] A. Nekvinda, *Hardy-Littlewood maximal operator on  $L^{p(x)}(\mathbb{R}^n)$* , Math. Inequal. Appl. **7** (2004), 255–266.
- [27] W. Orlicz, *Über konjugierte Exponentenfolgen*, Studia Math. **3** (1931), 200–212.
- [28] M. Růžička, *Electrorheological fluids: modeling and mathematical theory*, Lecture Notes in Mathematics **1748**, Springer, Berlin, 2000.
- [29] E. Sawyer, *Boundedness of classical operators on classical Lorentz spaces*, Studia Math. **96** no. 2 (1990), 145–158.
- [30] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30 Princeton University Press, Princeton, N.J. 1970.

- [31] E. M. Stein, G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton Mathematical Series, No. 32. Princeton University Press, Princeton, N.J., 1971.
- [32] H. Triebel, *Interpolation theory, function spaces, differential operators*, Verlag der Wissenschaften, Berlin, 1978.

MATHEMATICAL INSTITUTE, FRIEDRICH-SCHILLER-UNIVERSITY JENA, D-07737 JENA, GERMANY

*Current address:* Institute of Applied Analysis, Technical University Bergakademie Freiberg, D-09596 Freiberg, Germany

*E-mail address:* [henning.kempka@uni-jena.de](mailto:henning.kempka@uni-jena.de)

DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY BERLIN, STREET OF 17. JUNE 136, D-10623 BERLIN, GERMANY

*E-mail address:* [vybiral@math.tu-berlin.de](mailto:vybiral@math.tu-berlin.de)