

Spaces of variable smoothness and integrability: Characterizations by local means and ball means of differences

Henning Kempka*, Jan Vybíral†

March 28, 2011

Abstract

We study the spaces $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ of Besov and Triebel-Lizorkin type as introduced recently in [1] and [17]. Both scales cover many classical spaces with fixed exponents as well as function spaces of variable smoothness and function spaces of variable integrability.

The spaces $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ have been introduced in [1] and [17] by Fourier analytical tools, as the decomposition of unity. Surprisingly, our main result states that these spaces also allow a characterization in the time-domain with the help of classical ball means of differences.

To that end, we first prove a local means characterization for $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ with the help of the so-called Peetre maximal functions. Our results do also hold for 2-microlocal function spaces $B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ which are a slight generalization of generalized smoothness spaces and spaces of variable smoothness.

Key words: Besov spaces, Triebel-Lizorkin spaces, variable smoothness, variable integrability, ball means of differences, Peetre maximal operator, 2-microlocal spaces.
2010 MSC: 46E35, 46E30, 42B25.

1 Introduction

Function spaces of variable integrability appeared in a work by Orlicz [41] already in 1931, but the recent interest in these spaces is based on the paper of Kováčik and Rákosník [32] together with the applications to the modeling of electrorheological fluids [45]. A fundamental breakthrough in spaces of variable integrability was the observation that, under certain regularity assumptions on $p(\cdot)$, the Hardy-Littlewood maximal operator is also bounded on $L_{p(\cdot)}(\mathbb{R}^n)$, see [13]. This result has been generalized to wider classes of exponents $p(\cdot)$ in [11], [40] and [15].

*Mathematical Institute, Friedrich-Schiller-University Jena, D-07737 Jena, Germany, email: henning.kempka@uni-jena.de

†Johann Radon Institute for Computational and Applied Mathematics, Austrian Academy of Sciences, Altenbergerstrasse 69, A-4040 Linz, Austria, email: jan.vybiral@oeaw.ac.at.

Besides electrorheological fluids, the spaces $L_{p(\cdot)}(\mathbb{R}^n)$ possess interesting applications in the theory of PDE's, variational calculus, financial mathematics and image processing. Recent overview of this vastly growing field is given in [16].

Sobolev and Besov spaces with variable smoothness but fixed integrability have been introduced in the late 60's and early 70's in the works of Unterberger [57], Višik and Eskin [58], Unterberger and Bokobza [56] and in the work of Beuzamy [7]. Leopold studied in [33] Besov spaces where the smoothness is determined by a symbol $a(x, \xi)$ of a certain class of hypoelliptic pseudodifferential operators. In the special case $a(x, \xi) = (1 + |\xi|^2)^{\sigma(x)/2}$ these spaces coincide with spaces of variable smoothness $B_{p,p}^{\sigma(x)}(\mathbb{R}^n)$.

A more general approach to spaces of variable smoothness are the so-called 2-microlocal function spaces $B_{p,q}^{\mathbf{w}}(\mathbb{R}^n)$ and $F_{p,q}^{\mathbf{w}}(\mathbb{R}^n)$. Here the smoothness in this spaces gets measured by a weight sequence $\mathbf{w} = (w_j)_{j=0}^{\infty}$. Besov spaces with such weight sequences appeared first in the works of Peetre [42] and Bony [9]. Establishing a wavelet characterization for 2-microlocal Hölder-Zygmund spaces in [23] it turned out that 2-microlocal spaces are well adapted in connection to regularity properties of functions ([24],[37],[35]). Spaces of variable smoothness are a special case of 2-microlocal function spaces and in [34] and [8] characterizations by differences have been given for certain classes of them.

The theories of function spaces with fixed smoothness and variable integrability and function spaces with variable smoothness and fixed integrability finally crossed each other in [17], where the authors introduced the function spaces of Triebel-Lizorkin type with variable smoothness and simultaneously with variable integrability. It turned out that many of the spaces mentioned above are really included in this new structure, see [17] and references therein. The key point to merge both lines of investigation has been the study of traces. From Theorem 3.13 in [17]

$$tr_{\mathbb{R}^{n-1}} F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) = F_{p(\cdot),p(\cdot)}^{s(\cdot)-1/p(\cdot)}(\mathbb{R}^{n-1})$$

one immediately recognizes why it is necessary to take all exponents variable if only $p(\cdot)$ or $s(\cdot)$ had been variable. So the trace embeddings may be described in a natural way in the context of these spaces. Furthermore, this was complemented in [59] by showing, that the classical Sobolev embedding theorem

$$F_{p_0(\cdot),q(\cdot)}^{s_0(\cdot)}(\mathbb{R}^n) \hookrightarrow F_{p_1(\cdot),q(\cdot)}^{s_1(\cdot)}(\mathbb{R}^n)$$

holds also in this scale of function spaces if the usual condition is replaced by its point-wise analogue

$$s_0(x) - n/p_0(x) = s_1(x) - n/p_1(x), \quad x \in \mathbb{R}^n.$$

Finally, Almeida and Hästö managed in [1] to adapt the definition of Besov spaces to the setting of variable smoothness and integrability and proved the Sobolev and other usual embeddings in this scale.

The properties of Besov and Triebel-Lizorkin spaces of variable smoothness and integrability known so far give a reasonable hope that these new scales of function

spaces enjoy sufficiently many properties to allow a local description of many effects, which up to now could only be described in a global way. Subsequently, for the spaces $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ there was a characterization by local means in [29] given. This characterization still works with Fourier analytical tools but the analyzing functions $k_0, k \in \mathcal{S}(\mathbb{R}^n)$ are compactly supported in the time-domain and we only need local values of f around $x \in \mathbb{R}^n$ to calculate the building blocks $k(2^{-j}, f)(x)$. This is in sharp contrast to the definition of the spaces by the decomposition of unity, cf. Definitions 1 and 3. For the spaces $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ we will prove a local means assertion of this type in Section 3 which will be helpful later on.

The main aim of this paper is to present another essential property of the function spaces from [17] and [1]. We prove the surprising result that these spaces $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ with variable smoothness and integrability do also allow a characterization purely in the time-domain by classical ball means of differences.

The paper is organized as follows. First of all we provide all necessary notation in Section 2. Since the proofs for spaces of variable smoothness and 2-microlocal function spaces work very similar (see Remark 2) we present our results for both scales. The proof for the local means characterization will be given in Section 3 in terms of 2-microlocal function spaces and we present the version for spaces of variable smoothness in Section 3.2. In Section 4 we prove the characterization by ball means of differences for $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and the version for 2-microlocal function spaces will be given in Section 4.5.

2 Notation

In this section we collect all the necessary definitions. We start with the variable Lebesgue spaces $L_{p(\cdot)}(\mathbb{R}^n)$. A measurable function $p : \mathbb{R}^n \rightarrow (0, \infty]$ is called a *variable exponent function* if it is bounded away from zero, i.e. if $p^- = \text{ess-inf}_{x \in \mathbb{R}^n} p(x) > 0$. We denote the set of all variable exponent functions by $\mathcal{P}(\mathbb{R}^n)$. We put also $p^+ = \text{ess-sup}_{x \in \mathbb{R}^n} p(x)$.

The variable exponent Lebesgue space $L_{p(\cdot)}(\mathbb{R}^n)$ consists of all measurable functions f such that there exist an $\lambda > 0$ such that the modular

$$\varrho_{L_{p(\cdot)}(\mathbb{R}^n)}(f/\lambda) = \int_{\mathbb{R}^n} \varphi_{p(x)} \left(\frac{|f(x)|}{\lambda} \right) dx$$

is finite, where

$$\varphi_p(t) = \begin{cases} t^p & \text{if } p \in (0, \infty), \\ 0 & \text{if } p = \infty \text{ and } t \leq 1, \\ \infty & \text{if } p = \infty \text{ and } t > 1. \end{cases}$$

If we define $\mathbb{R}_\infty^n = \{x \in \mathbb{R}^n : p(x) = \infty\}$ and $\mathbb{R}_0^n = \mathbb{R}^n \setminus \mathbb{R}_\infty^n$, then the Luxemburg

norm of a function $f \in L_{p(\cdot)}(\mathbb{R}^n)$ is given by

$$\begin{aligned} \|f\|_{L_{p(\cdot)}(\mathbb{R}^n)} &= \inf\{\lambda > 0 : \varrho_{L_{p(\cdot)}(\mathbb{R}^n)}(f/\lambda) \leq 1\} \\ &= \inf\left\{\lambda > 0 : \int_{\mathbb{R}_0^n} \left(\frac{f(x)}{\lambda}\right)^{p(x)} dx < 1 \text{ and } |f(x)| < \lambda \text{ for a.e. } x \in \mathbb{R}_\infty^n\right\}. \end{aligned}$$

If $p(\cdot) \geq 1$, then it is a norm otherwise it is always a quasi-norm.

To define the mixed spaces $\ell_{q(\cdot)}(L_{p(\cdot)})$ we have to define another modular. For $p, q \in \mathcal{P}(\mathbb{R}^n)$ and a sequence $(f_\nu)_{\nu \in \mathbb{N}_0}$ of $L_{p(\cdot)}(\mathbb{R}^n)$ functions we define

$$\varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}(f_\nu) = \sum_{\nu=0}^{\infty} \inf\left\{\lambda_\nu > 0 : \varrho_{p(\cdot)}\left(\frac{f_\nu}{\lambda_\nu^{1/q(\cdot)}}\right) \leq 1\right\}. \quad (1)$$

If $q^+ < \infty$, then we can replace (1) by the simpler expression

$$\varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}(f_\nu) = \sum_{\nu} \left\| |f_\nu|^{q(\cdot)} \right\|_{L_{\frac{p(\cdot)}{q(\cdot)}}}. \quad (2)$$

The (quasi-)norm in the $\ell_{q(\cdot)}(L_{p(\cdot)})$ spaces is defined as usual by

$$\|f_\nu\|_{\ell_{q(\cdot)}(L_{p(\cdot)})} = \inf\{\mu > 0 : \varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}(f_\nu/\mu) \leq 1\}. \quad (3)$$

It is known, cf. [1, 31], that $\ell_{q(\cdot)}(L_{p(\cdot)})$ is a norm if $q(\cdot) \geq 1$ is constant almost everywhere (a.e.) on \mathbb{R}^n and $p(\cdot) \geq 1$, or if $1/p(x) + 1/q(x) \leq 1$ a.e. on \mathbb{R}^n , or if $1 \leq q(x) \leq p(x) \leq \infty$ a.e. on \mathbb{R}^n . Surprisingly enough, it turned out in [31] that the condition $\min(p(x), q(x)) \geq 1$ a.e. on \mathbb{R}^n is not sufficient for $\ell_{q(\cdot)}(L_{p(\cdot)})$ to be a norm. Nevertheless, it was proven in [1] that it is a quasi-norm for every $p, q \in \mathcal{P}(\mathbb{R}^n)$.

For the sake of completeness, we state also the definition of the space $L_{p(\cdot)}(\ell_{q(\cdot)})$, which is much more intuitive than the definition of $\ell_{q(\cdot)}(L_{p(\cdot)})$. One just takes the $\ell_{q(x)}$ norm of $(f_\nu(x))_{\nu \in \mathbb{N}_0}$ for every $x \in \mathbb{R}^n$ and then the $L_{p(\cdot)}$ -norm with respect to $x \in \mathbb{R}^n$, i.e.

$$\|f_\nu\|_{L_{p(\cdot)}(\ell_{q(\cdot)})} = \left\| \left\| |f_\nu(x)| \right\|_{\ell_{q(x)}} \right\|_{L_{p(\cdot)}}.$$

It is easy to show ([17]) that $L_{p(\cdot)}(\ell_{q(\cdot)})$ is always a quasi-normed space and it is a normed space, if $\min(p(x), q(x)) \geq 1$ holds pointwise.

The summation in the definition of the norms of $\ell_{q(\cdot)}(L_{p(\cdot)})$ and $L_{p(\cdot)}(\ell_{q(\cdot)})$ can also be taken for $\nu \in \mathbb{Z}$. It always comes out of the context over which interval the summation is taken. Occasionally, we may indicate it by $\|(f_\nu)_{\nu=-\infty}^{\infty}\|_{\ell_{q(\cdot)}(L_{p(\cdot)})}$.

By $\hat{f} = \mathcal{F}f$ and $f^\vee = \mathcal{F}^{-1}f$ we denote the usual Fourier transform and its inverse on $\mathcal{S}(\mathbb{R}^n)$, the Schwartz space of smooth and rapidly decreasing functions, and on $\mathcal{S}'(\mathbb{R}^n)$, the dual of the Schwartz space.

2.1 Spaces $B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$

The definition of Besov and Triebel-Lizorkin spaces of variable smoothness and integrability is based on the technique of *decomposition of unity* exactly in the same manner as in the case of constant exponents.

Definition 1. Let $\varphi_0 \in \mathcal{S}(\mathbb{R}^n)$ with $\varphi_0(x) = 1$ for $|x| \leq 1$ and $\text{supp } \varphi_0 \subseteq \{x \in \mathbb{R}^n : |x| \leq 2\}$. For $j \geq 1$ we define

$$\varphi_j(x) = \varphi_0(2^{-j}x) - \varphi_0(2^{-j+1}x).$$

One may verify easily that

$$\sum_{j=0}^{\infty} \varphi_j(x) = 1 \quad \text{for all } x \in \mathbb{R}^n.$$

The following regularity classes for the exponents are necessary to make the definition of the spaces independent on the chosen decomposition of unity.

Definition 2. Let $g \in C(\mathbb{R}^n)$.

- (i) We say that g is *locally log-Hölder continuous*, abbreviated $g \in C_{loc}^{\log}(\mathbb{R}^n)$, if there exists $c_{\log}(g) > 0$ such that

$$|g(x) - g(y)| \leq \frac{c_{\log}(g)}{\log(e + 1/|x - y|)} \quad (4)$$

holds for all $x, y \in \mathbb{R}^n$.

- (ii) We say that g is *globally log-Hölder continuous*, abbreviated $g \in C^{\log}(\mathbb{R}^n)$, if g is locally log-Hölder continuous and there exists $g_{\infty} \in \mathbb{R}$ such that

$$|g(x) - g_{\infty}| \leq \frac{c_{\log}}{\log(e + |x|)} \quad (5)$$

holds for all $x \in \mathbb{R}^n$.

Remark 1. With (4) we obtain

$$|g(x)| \leq c_{\log}(g) + |g(0)|, \quad \text{for all } x \in \mathbb{R}^n.$$

This implies that all functions $g \in C_{loc}^{\log}(\mathbb{R}^n)$ always belong to $L_{\infty}(\mathbb{R}^n)$

If an exponent $p \in \mathcal{P}(\mathbb{R}^n)$ satisfies $1/p \in C^{\log}(\mathbb{R}^n)$, then we say it belongs to the class $\mathcal{P}^{\log}(\mathbb{R}^n)$. We recall the definition of the spaces $B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$, as given in [17] and [1].

Definition 3. (i) Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < p^- \leq p^+ < \infty$, $0 < q^- \leq q^+ < \infty$ and let $s \in C_{loc}^{\log}(\mathbb{R}^n)$. Then

$$F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : \left\| f \Big| F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \right\|_{\varphi} < \infty \right\}, \text{ where}$$

$$\left\| f \Big| F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \right\|_{\varphi} = \left\| 2^{js(\cdot)}(\varphi_j \hat{f})^{\vee} \Big| L_{p(\cdot)}(\ell_{q(\cdot)}) \right\|.$$

(ii) Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and let $s \in C_{loc}^{\log}(\mathbb{R}^n)$. Then

$$B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : \left\| f \Big| B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \right\|_{\varphi} < \infty \right\}, \text{ where}$$

$$\left\| f \Big| B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \right\|_{\varphi} = \left\| 2^{js(\cdot)}(\varphi_j \hat{f})^{\vee} \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\|.$$

The subscript φ at the norm symbolizes that the definition formally does depend on the resolution of unity. From [29] and [1] we have that the definition of the spaces $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ is independent of the chosen resolution of unity if $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and $s \in C_{loc}^{\log}(\mathbb{R}^n)$. That means that different start functions φ_0 and $\tilde{\varphi}_0$ from Definition 1 induce equivalent norms in the above definition. So we will suppress the subscript φ in the notation of the norms.

Let us comment on the conditions on $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ for the Triebel-Lizorkin spaces. The condition $0 < p^- \leq p^+ < \infty$ is quite natural since there exists also the restriction $p < \infty$ in the case of constant exponents, see [51] and [63]. The second one, $0 < q^- \leq q^+ < \infty$, is a bit unnatural and comes from the use of the convolution Lemma 21 ([17, Theorem 3.2]). There is some hope that this convolution lemma can be generalized and the case $q^+ = \infty$ can be incorporated in the definition of the F -spaces.

The Triebel-Lizorkin spaces with variable smoothness have first been introduced in [17] under much more restrictive conditions on $s(\cdot)$. These conditions have been relaxed in [29] in the context of 2-microlocal function spaces (see the next subsection).

Besov spaces with variable $p(\cdot), q(\cdot)$ and $s(\cdot)$ have been introduced in [1].

Both scales contain as special cases a lot of well known function spaces. If s, p and q are constants, then we derive the well known Besov and Triebel-Lizorkin spaces with usual Hölder and Sobolev spaces included, see [51] and [52]. If the smoothness $s \in \mathbb{R}$ is a constant and $p \in \mathcal{P}^{\log}(\mathbb{R}^n)$, then $F_{p(\cdot),2}^s(\mathbb{R}^n) = \mathcal{L}_{p(\cdot)}^s(\mathbb{R}^n)$ are the variable Bessel potential spaces from [2] and [22] with its special cases $F_{p(\cdot),2}^0(\mathbb{R}^n) = L_{p(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),2}^k(\mathbb{R}^n) = W_{p(\cdot)}^k(\mathbb{R}^n)$ for $k \in \mathbb{N}_0$, see [17].

Taking $s \in \mathbb{R}$ and $q \in (0, \infty]$ as constants we derive the spaces $F_{p(\cdot),q}^s(\mathbb{R}^n)$ and $B_{p(\cdot),q}^s(\mathbb{R}^n)$ studied by Xu in [61] and [62].

Furthermore it holds $F_{p(\cdot),p(\cdot)}^{s(\cdot)}(\mathbb{R}^n) = B_{p(\cdot),p(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $B_{\infty,\infty}^{s(\cdot)}(\mathbb{R}^n)$ equals the variable Hölder-Zygmund space $\mathcal{C}^{s(\cdot)}(\mathbb{R}^n)$ introduced in [3], [4] and [44] with $0 < s^- \leq s^+ \leq 1$, see [1].

2.2 2-microlocal spaces

The definition of Besov and Triebel-Lizorkin spaces of variable smoothness and integrability is a special case of the so-called 2-microlocal spaces of variable integrability. As some of the results presented here get proved in this more general scale, we present also the definition of 2-microlocal spaces. It is based on the dyadic decomposition of unity as presented above combined with the concept of admissible weight sequences.

Definition 4. Let $\alpha \geq 0$ and let $\alpha_1, \alpha_2 \in \mathbb{R}$ with $\alpha_1 \leq \alpha_2$. A sequence of non-negative measurable functions $\mathbf{w} = (w_j)_{j=0}^{\infty}$ belongs to the class $\mathcal{W}_{\alpha_1, \alpha_2}^{\alpha}$ if and only if

- (i) there exists a constant $C > 0$ such that

$$0 < w_j(x) \leq C w_j(y) (1 + 2^j |x - y|)^{\alpha} \quad \text{for all } j \in \mathbb{N}_0 \text{ and all } x, y \in \mathbb{R}^n$$

(ii) and for all $j \in \mathbb{N}_0$ and all $x \in \mathbb{R}^n$ we have

$$2^{\alpha_1} w_j(x) \leq w_{j+1}(x) \leq 2^{\alpha_2} w_j(x).$$

Such a system $(w_j)_{j=0}^\infty \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha$ is called an *admissible weight sequence*.

Finally, here is the definition of the spaces under consideration.

Definition 5. Let $\mathbf{w} = (w_j)_{j \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha$. Further, let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ (with $p^+, q^+ < \infty$ in the F -case), then we define

$$B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : \left\| f \Big| B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \right\|_\varphi < \infty \right\}, \text{ where}$$

$$\left\| f \Big| B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \right\|_\varphi = \left\| w_j (\varphi_j \hat{f})^\vee \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\|$$

and

$$F_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : \left\| f \Big| F_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \right\|_\varphi < \infty \right\}, \text{ where}$$

$$\left\| f \Big| F_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \right\|_\varphi = \left\| w_j (\varphi_j \hat{f})^\vee \Big| L_{p(\cdot)}(\ell_{q(\cdot)}) \right\|.$$

The independence of the decomposition of unity for the 2-microlocal spaces from Definition 5 follows from the local means characterization (see [29] for the Triebel-Lizorkin and Section 3 for the Besov spaces).

The 2-microlocal spaces with the special weight sequence

$$w_j(x) = 2^{js} (1 + 2^j |x - x_0|)^{s'} \quad \text{with } s, s' \in \mathbb{R} \text{ and } x_0 \in \mathbb{R}^n \quad (6)$$

have first been introduced by Peetre in [42] and by Bony in [9]. Later on, Jaffard and Meyer in [23] and [24] gave a characterization with wavelets of the spaces $C_{x_0}^{s, s'} = B_{\infty, \infty}^{\mathbf{w}}(\mathbb{R}^n)$ and $H_{x_0}^{s, s'} = B_{2, 2}^{\mathbf{w}}(\mathbb{R}^n)$ with the weight sequence (6). It turned out that spaces of this type are very useful to study regularity properties of functions. Subsequently, Lévy-Véhel and Seuret developed in [35] the 2-microlocal formalism and studied the behavior of cusps, chirps and fractal functions with respect to the spaces $C_{x_0}^{s, s'}$.

A first step to a more general weight sequence \mathbf{w} has been taken by Moritoh and Yamada in [38] and wider ranges of function spaces have been studied by Xu in [60] and by Andersson in [5].

The above definition for 2-microlocal weight sequences was presented by Besov in [8] and also in [29] by Kempka.

A different line of study to spaces of variable smoothness are the spaces of generalized smoothness introduced using different methods by Goldman and Kalyabin in [19], [20], [25] and [26]. A systematic treatment of these spaces based on differences has been given by Goldman in [21], see also the survey [27] and references therein. Later on, spaces of generalized smoothness appeared in interpolation theory and have been investigated in [36], [10] and [39]. For further information on these spaces see the survey paper [18] where also a characterization by atoms and local means

for that spaces is given.

From the definition of admissible weight sequences, $d_1\sigma_j \leq \sigma_{j+1} \leq d_2\sigma_j$, it follows directly that the spaces of generalized smoothness $B_{p,q}^{(\sigma_j)}(\mathbb{R}^n)$ and $F_{p,q}^{(\sigma_j)}(\mathbb{R}^n)$ of Farkas and Leopold [18] and $B_{p,q}^{(s,\Psi)}(\mathbb{R}^n)$ and $F_{p,q}^{(s,\Psi)}(\mathbb{R}^n)$ from Moura [39] are a special subclass of 2-microlocal function spaces with $\alpha_1 = d_1$, $\alpha_2 = d_2$ and $\alpha = 0$.

In a different approach Schneider in [50] studied spaces of varying smoothness. Here the smoothness at a point gets determined by a global smoothness $s_0 \in \mathbb{R}$ and a local smoothness function $s(\cdot)$. These spaces can not be incorporated into the scale of 2-microlocal function spaces, but there exist some embeddings.

Remark 2. Surprisingly, these 2-microlocal weight sequences are directly connected to variable smoothness functions $s : \mathbb{R}^n \rightarrow \mathbb{R}$ if we set

$$w_j(x) = 2^{js(x)}. \quad (7)$$

If $s \in C_{loc}^{\log}(\mathbb{R}^n)$ (which is the standard condition on $s(\cdot)$), then $\mathbf{w} = (w_j(x))_{j \in \mathbb{N}_0} = (2^{js(x)})_{j \in \mathbb{N}_0}$ belongs to $\mathcal{W}_{\alpha_1, \alpha_2}^\alpha$ with $\alpha_1 = s^-$ and $\alpha_2 = s^+$. For the third index α we use Lemma 19 with $m = 0$ and obtain $\alpha = c_{\log}(s)$, where $c_{\log}(s)$ is the constant for $s(\cdot)$ from (4). That means that spaces of variable smoothness from Definition 3 are a special case of 2-microlocal function spaces from Definition 5. Both types of function spaces are very closely connected and the properties used in the proofs are either

$$2^{k|s(x)-s(y)|} \leq c \quad \text{or} \quad \frac{w_k(x)}{w_k(y)} \leq c \quad (8)$$

for $|x - y| \leq c2^{-k}$. This property follows directly either from the definition of $s \in C_{loc}^{\log}(\mathbb{R}^n)$ or from Definition 4.

Nevertheless there exist examples of admissible weight sequences which can not be expressed in terms of variable smoothness functions. For example the important and well studied case of the weight sequence \mathbf{w} from (6) can not be expressed via (7) if $s' \neq 0$. Another example are the spaces of generalized smoothness which can not be identified as spaces of variable smoothness.

Since spaces of variable smoothness are included in the scale of 2-microlocal function spaces all special cases of the previous subsection can be identified in the definition of 2-microlocal spaces.

Although the 2-microlocal spaces include the scales of spaces of variable smoothness, we will give some of our proofs in the notation of variable smoothness, since this notation is more common. We will then reformulate the results in terms of 2-microlocal spaces, the proof works than very similar; we just have to use (8).

3 Local means characterization

The main result of this section is the local means characterization of the spaces $B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$. For the spaces $F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ there already exists a local means characterization ([29, Corollary 4.7]). We shall first give the full proof for the 2-microlocal

spaces and later on (in Section 3.2) we restate the result also for spaces $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$.

The crucial tool will be the Peetre maximal operator, as defined by Peetre in [42]. The operator assigns to each system $(\Psi_k)_{k \in \mathbb{N}_0} \subset \mathcal{S}(\mathbb{R}^n)$, to each distribution $f \in \mathcal{S}'(\mathbb{R}^n)$ and to each number $a > 0$ the following quantities

$$(\Psi_k^* f)_a(x) := \sup_{y \in \mathbb{R}^n} \frac{|(\Psi_k * f)(y)|}{1 + |2^k(y-x)|^a}, \quad x \in \mathbb{R}^n \text{ and } k \in \mathbb{N}_0. \quad (9)$$

We start with two given functions $\psi_0, \psi_1 \in \mathcal{S}(\mathbb{R}^n)$. We define

$$\psi_j(x) = \psi_1(2^{-j+1}x), \quad \text{for } x \in \mathbb{R}^n \text{ and } j \in \mathbb{N}. \quad (10)$$

Furthermore, for all $j \in \mathbb{N}_0$ we write $\Psi_j = \hat{\psi}_j$. The main theorem of this section reads as follows.

Theorem 6. *Let $w = (w_k)_{k \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha$, $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and let $a > 0$, $R \in \mathbb{N}_0$ with $R > \alpha_2$. Further, let ψ_0, ψ_1 belong to $\mathcal{S}(\mathbb{R}^n)$ with*

$$D^\beta \psi_1(0) = 0, \quad \text{for } 0 \leq |\beta| < R, \quad (11)$$

and

$$|\psi_0(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : |x| < \varepsilon\} \quad (12)$$

$$|\psi_1(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : \varepsilon/2 < |x| < 2\varepsilon\} \quad (13)$$

for some $\varepsilon > 0$. For $a > \frac{n+c_{\log}(1/q)}{p^-} + \alpha$ and all $f \in \mathcal{S}'(\mathbb{R}^n)$ we have

$$\left\| f | B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n) \right\| \sim \left\| (\Psi_k * f) w_k | \ell_{q(\cdot)}(L_{p(\cdot)}) \right\| \sim \left\| (\Psi_k^* f)_a w_k | \ell_{q(\cdot)}(L_{p(\cdot)}) \right\|.$$

Remark 3. (i) The proof relies on [46] and will be shifted to the next section. Moreover, Theorem 6 shows that the definition of the 2-microlocal spaces of variable integrability is independent of the resolution of unity used in the Definition 5.

(ii) The conditions (11) are usually called *moment conditions* while (12) and (13) are the so called *Tauberian conditions*.

(iii) If $R = 0$, then there are no moment conditions (11) on ψ_1 .

(iv) The notation $c_{\log}(1/q)$ stays for the constant from (4) with $1/q(\cdot)$.

Next we reformulate the abstract Theorem 6 in the sense of classical local means (see Sections 2.4.6 and 2.5.3 in [52]). Since the proof is the same as the one from Theorem 2.4 in [29] we just state the result.

Corollary 1. *There exist functions $k_0, k \in \mathcal{S}(\mathbb{R}^n)$ with $\text{supp } k_0, \text{supp } k \subset \{x \in \mathbb{R}^n : |x| < 1\}$ and $D^\beta k(0) = 0$ for all $0 \leq |\beta| < \alpha_2$ such that for all $f \in \mathcal{S}'(\mathbb{R}^n)$*

$$\left\| k_0(1, f) w_0 | L_{p(\cdot)}(\mathbb{R}^n) \right\| + \left\| k(2^{-j}, f) w_j | \ell_{q(\cdot)}(L_{p(\cdot)}) \right\|$$

is an equivalent norm on $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$.
The building blocks get calculated by

$$k(t, f)(x) = \int_{\mathbb{R}^n} k(y)f(x + ty)dy = t^{-n} \int_{\mathbb{R}^n} k\left(\frac{y-x}{t}\right) f(y)dy$$

and similarly for $k_0(1, f)(x)$.

A similar characterization for $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and details how these functions $k_0, k \in \mathcal{S}(\mathbb{R}^n)$ can be constructed can be found in [29].

3.1 Proof of local means

The proof of Theorem 6 is divided into three parts. The next section is devoted to some technical lemmas needed later. Section 3.1.2 is devoted to the proof of Theorem 12, which gives an inequality between different Peetre maximal operators. Finally, Section 3.1.3 proves the boundedness of the Peetre maximal operator in Theorem 13. These two theorems combined give immediately the proof of Theorem 6.

3.1.1 Helpful lemmas

Before proving the local means characterization we recall some technical lemmas, which appeared in the paper of Rychkov [46]. For some of them we need adapted versions to our situation.

The first lemma describes the use of the so called moment conditions.

Lemma 7 ([46], Lemma 1). *Let $g, h \in \mathcal{S}(\mathbb{R}^n)$ and let $M \in \mathbb{N}_0$. Suppose that*

$$(D^{|\beta|}\hat{g})(0) = 0 \quad \text{for } 0 \leq |\beta| < M. \quad (14)$$

Then for each $N \in \mathbb{N}_0$ there is a constant C_N such that

$$\sup_{z \in \mathbb{R}^n} |(g_t * h)(z)|(1 + |z|^N) \leq C_N t^M, \quad \text{for } 0 < t < 1, \quad (15)$$

where $g_t(x) = t^{-n}g(x/t)$.

The next lemma is a discrete convolution inequality. We formulate it in a rather abstract notation and point out later on the conclusions we need.

Lemma 8. *Let $X \subset \{(f_k)_{k \in \mathbb{Z}} : f_k : \mathbb{R}^n \rightarrow [-\infty, \infty] \text{ measurable}\}$ be a quasi-Banach space of sequences of measurable functions. Further we assume that its quasi-norm is shift-invariant, i.e. it satisfies*

$$\|(f_{k+l})_{k \in \mathbb{Z}}\|_X = \|(f_k)_{k \in \mathbb{Z}}\|_X \quad \text{for every } l \in \mathbb{Z} \text{ and } (f_k)_{k \in \mathbb{Z}} \in X. \quad (16)$$

For a sequence of non-negative functions $(g_k)_{k \in \mathbb{Z}} \in X$ and $\delta > 0$ we denote

$$G_\nu(x) = \sum_{k=-\infty}^{\infty} 2^{-|\nu-k|\delta} g_k(x), \quad x \in \mathbb{R}^n, \nu \in \mathbb{Z}. \quad (17)$$

Then there exists a constant $c > 0$ depending only on δ and X such that for every sequence $(g_k)_{k \in \mathbb{Z}}$

$$\|(G_\nu)_\nu | X\| \leq c \|(g_k)_k | X\|.$$

Proof. Since X is a quasi-Banach space, there exists a $r > 0$ such that $\|\cdot\|_X$ is equivalent to some r -norm, cf. [6, 43]. We have then the following

$$\begin{aligned} \|(G_\nu)_\nu | X\|^r &= \left\| \left(\sum_{k=-\infty}^{\infty} 2^{-|\nu-k|\delta} g_k \right)_\nu \right\|_X^r = \left\| \left(\sum_{l \in \mathbb{Z}} 2^{-|l|\delta} g_{\nu+l} \right)_\nu \right\|_X^r \\ &\lesssim \sum_{l \in \mathbb{Z}} 2^{-|l|r\delta} \|(g_{\nu+l})_\nu | X\|^r \leq c \|(g_\nu)_\nu | X\|^r. \end{aligned}$$

Now taking the power $1/r$ yields the desired estimate. \square

The spaces $L_{p(\cdot)}(\ell_{q(\cdot)})$ and $\ell_{q(\cdot)}(L_{p(\cdot)})$ are quasi-Banach spaces which fulfill the conditions of Lemma 8. Therefore, we obtain the following

Lemma 9. *Let $p, q \in \mathcal{P}(\mathbb{R}^n)$ and $\delta > 0$. Let $(g_k)_{k \in \mathbb{Z}}$ be a sequence of non-negative measurable functions on \mathbb{R}^n and denote*

$$G_\nu(x) = \sum_{k \in \mathbb{Z}} 2^{-|\nu-k|\delta} g_k(x), \quad x \in \mathbb{R}^n, \nu \in \mathbb{Z}. \quad (18)$$

Then there exist constants $C_1, C_2 > 0$, depending on $p(\cdot), q(\cdot)$ and δ , such that

$$\begin{aligned} \|G_\nu | \ell_{q(\cdot)}(L_{p(\cdot)})\| &\leq C_1 \|g_k | \ell_{q(\cdot)}(L_{p(\cdot)})\| \quad \text{and} \\ \|G_\nu | L_{p(\cdot)}(\ell_{q(\cdot)})\| &\leq C_2 \|g_k | L_{p(\cdot)}(\ell_{q(\cdot)})\|. \end{aligned}$$

Remark 4. Of course, Lemma 9 holds true also if the indices k and ν run only over natural numbers.

Since the maximal operator is in general not bounded on $\ell_{q(\cdot)}(L_{p(\cdot)})$ (see [1, Example 4.1]) we need an replacement for that. It turned out that a convolution with radial decreasing functions fits very well into the scheme. A careful evaluation of the proof in [1, Lemma 4.7] together with Lemma 19 gives us the following convolution inequality.

Lemma 10. *Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $p(\cdot) \geq 1$ and let $\eta_{\nu, m}(x) = 2^{\nu m} (1 + 2^\nu |x|)^{-m}$. For all $m > n + c_{\log}(1/q)$ there exists a constant $c > 0$ such that for all sequences $(f_j)_{j \in \mathbb{N}_0} \in \ell_{q(\cdot)}(L_{p(\cdot)})$ it holds*

$$\|(\eta_{\nu, m} * f_\nu)_{\nu \in \mathbb{N}_0} | \ell_{q(\cdot)}(L_{p(\cdot)})\| \leq c \|(f_j)_{j \in \mathbb{N}_0} | \ell_{q(\cdot)}(L_{p(\cdot)})\|.$$

The last technical lemma is overtaken literally from [46].

Lemma 11 ([46], Lemma 3). *Let $0 < r \leq 1$ and let $(\gamma_\nu)_{\nu \in \mathbb{N}_0}, (\beta_\nu)_{\nu \in \mathbb{N}_0}$ be two sequences taking values in $(0, \infty)$. Assume that for some $N^0 \in \mathbb{N}_0$,*

$$\limsup_{\nu \rightarrow \infty} \frac{\gamma_\nu}{2^{\nu n_0}} < \infty. \quad (19)$$

Furthermore, we assume that for any $N \in \mathbb{N}$

$$\gamma_\nu \leq C_N \sum_{k=0}^{\infty} 2^{-kN} \beta_{k+\nu} \gamma_{k+\nu}^{1-r}, \quad \nu \in \mathbb{N}_0, \quad C_N < \infty$$

holds, then for any $N \in \mathbb{N}$

$$\gamma_\nu^r \leq C_N \sum_{k=0}^{\infty} 2^{-kNr} \beta_{k+\nu}, \quad \nu \in \mathbb{N}_0 \quad (20)$$

holds with the same constants C_N .

3.1.2 Comparison of different Peetre maximal operators

In this subsection we present an inequality between different Peetre maximal operators. Let us recall the notation given before Theorem 6. For two given functions $\psi_0, \psi_1 \in \mathcal{S}(\mathbb{R}^n)$ we define

$$\psi_j(x) = \psi_1(2^{-j+1}x), \quad \text{for } x \in \mathbb{R}^n \text{ and } j \in \mathbb{N}. \quad (21)$$

Furthermore, for all $j \in \mathbb{N}_0$ we write $\Psi_j = \hat{\psi}_j$ and in an analogous manner we define Φ_j from two starting functions $\phi_0, \phi_1 \in \mathcal{S}(\mathbb{R}^n)$. Using this notation we are ready to formulate the theorem.

Theorem 12. *Let $\mathbf{w} = (w_j)_{j \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha$, $p, q \in \mathcal{P}(\mathbb{R}^n)$ and $a > 0$. Moreover, let $R \in \mathbb{N}_0$ with $R > \alpha_2$,*

$$D^\beta \psi_1(0) = 0, \quad 0 \leq |\beta| < R \quad (22)$$

and for some $\varepsilon > 0$

$$|\phi_0(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : |x| < \varepsilon\} \quad (23)$$

$$|\phi_1(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : \varepsilon/2 < |x| < 2\varepsilon\}, \quad (24)$$

then

$$\|(\Psi_k^* f)_a w_k|_{\ell_{q(\cdot)}(L_{p(\cdot)})}\| \leq c \|(\Phi_k^* f)_a w_k|_{\ell_{q(\cdot)}(L_{p(\cdot)})}\| \quad (25)$$

holds for every $f \in \mathcal{S}'(\mathbb{R}^n)$.

Remark 5. Observe that there are no restrictions on $a > 0$ and $p, q \in \mathcal{P}(\mathbb{R}^n)$ in the theorem above.

Proof. We have the fixed resolution of unity from Definition 1 and define the sequence of functions $(\lambda_j)_{j \in \mathbb{N}_0}$ by

$$\lambda_j(x) = \frac{\varphi_j\left(\frac{2x}{\varepsilon}\right)}{\phi_j(x)}.$$

It follows from the Tauberian conditions (23) and (24) that they satisfy

$$\sum_{j=0}^{\infty} \lambda_j(x) \phi_j(x) = 1, \quad x \in \mathbb{R}^n, \quad (26)$$

$$\lambda_j(x) = \lambda_1(2^{-j+1}x), \quad x \in \mathbb{R}^n, \quad j \in \mathbb{N}, \quad (27)$$

and

$$\text{supp } \lambda_0 \subset \{x \in \mathbb{R}^n : |x| \leq \varepsilon\} \quad \text{and} \quad \text{supp } \lambda_1 \subset \{x \in \mathbb{R}^n : \varepsilon/2 \leq |x| \leq 2\varepsilon\}. \quad (28)$$

Furthermore, we denote $\Lambda_k = \hat{\lambda}_k$ for $k \in \mathbb{N}_0$ and obtain together with (26) the following identities (convergence in $\mathcal{S}'(\mathbb{R}^n)$)

$$f = \sum_{k=0}^{\infty} \Lambda_k * \Phi_k * f, \quad \Psi_\nu * f = \sum_{k=0}^{\infty} \Psi_\nu * \Lambda_k * \Phi_k * f. \quad (29)$$

We have

$$\begin{aligned} |(\Psi_\nu * \Lambda_k * \Phi_k * f)(y)| &\leq \int_{\mathbb{R}^n} |(\Psi_\nu * \Lambda_k)(z)| |(\Phi_k * f)(y-z)| dz \\ &\leq (\Phi_k^* f)_a(y) \int_{\mathbb{R}^n} |(\Psi_\nu * \Lambda_k)(z)| (1 + |2^k z|^a) dz \\ &=: (\Phi_k^* f)_a(y) I_{\nu,k}, \end{aligned} \quad (30)$$

where

$$I_{\nu,k} := \int_{\mathbb{R}^n} |(\Psi_\nu * \Lambda_k)(z)| (1 + |2^k z|^a) dz.$$

According to Lemma 7 we get

$$I_{\nu,k} \leq c \begin{cases} 2^{(k-\nu)R}, & k \leq \nu, \\ 2^{(\nu-k)(a+1+|\alpha_1|)}, & \nu \leq k. \end{cases} \quad (31)$$

Namely, we have for $1 \leq k < \nu$ with the change of variables $2^k z \mapsto z$

$$\begin{aligned} I_{\nu,k} &= 2^{-n} \int_{\mathbb{R}^n} |(\Psi_{\nu-k} * \Lambda_1(\cdot/2))(z)| (1 + |z|^a) dz \\ &\leq c \sup_{z \in \mathbb{R}^n} |(\Psi_{\nu-k} * \Lambda_1(\cdot/2))(z)| (1 + |z|)^{a+n+1} \leq c 2^{(k-\nu)R}. \end{aligned}$$

Similarly, we get for $1 \leq \nu < k$ with the substitution $2^\nu z \mapsto z$

$$\begin{aligned} I_{\nu,k} &= 2^{-n} \int_{\mathbb{R}^n} |(\Psi_1(\cdot/2) * \Lambda_{k-\nu})(z)| (1 + |2^{k-\nu} z|^a) dz \\ &\leq c 2^{(\nu-k)(M-a)}. \end{aligned}$$

M can be taken arbitrarily large because Λ_1 has infinitely many vanishing moments. Taking $M = 2a + |\alpha_1| + 1$ we derive (31) for the cases $k, \nu \geq 1$ with $k \neq \nu$. The

missing cases can be treated separately in an analogous manner. The needed moment conditions are always satisfied by (22) and (28). The case $k = \nu = 0$ is covered by the constant c in (31).

Furthermore, we have

$$\begin{aligned} (\Phi_k^* f)_a(y) &\leq (\Phi_k^* f)_a(x)(1 + |2^k(x - y)|^a) \\ &\leq (\Phi_k^* f)_a(x)(1 + |2^\nu(x - y)|^a) \max(1, 2^{(k-\nu)a}). \end{aligned}$$

We put this into (30) and get

$$\sup_{y \in \mathbb{R}^n} \frac{|(\Psi_\nu * \Lambda_k * \Phi_k^* f)(y)|}{1 + |2^\nu(x - y)|^a} \leq c(\Phi_k^* f)_a(x) \begin{cases} 2^{(k-\nu)R}, & k \leq \nu, \\ 2^{(\nu-k)(1+|\alpha_1|)}, & k \geq \nu. \end{cases}$$

Multiplying both sides with $w_\nu(x)$ and using

$$w_\nu(x) \leq w_k(x) \begin{cases} 2^{(k-\nu)(-\alpha_2)}, & k \leq \nu, \\ 2^{(\nu-k)\alpha_1}, & k \geq \nu, \end{cases} \quad (32)$$

leads us to

$$\sup_{y \in \mathbb{R}^n} \frac{|(\Psi_\nu * \Lambda_k * \Phi_k^* f)(y)|}{1 + |2^\nu(x - y)|^a} w_\nu(x) \leq c(\Phi_k^* f)_a(x) w_k(x) \begin{cases} 2^{(k-\nu)(R-\alpha_2)}, & k \leq \nu, \\ 2^{(\nu-k)}, & k \geq \nu. \end{cases}$$

This inequality together with (29) gives for $\delta := \min(1, R - \alpha_2) > 0$

$$(\Psi_\nu^* f)_a(x) w_\nu(x) \leq c \sum_{k=0}^{\infty} 2^{-|k-\nu|\delta} (\Phi_k^* f)_a(x) w_k(x), \quad x \in \mathbb{R}^n. \quad (33)$$

Taking the $\ell_{q(\cdot)}(L_{p(\cdot)})$ norm and using Lemma 9 yields immediately the desired result. \square

3.1.3 Boundedness of the Peetre maximal operator

We will present a theorem which describes the boundedness of the Peetre maximal operator. We use the same notation introduced at the beginning of the last subsection. Especially, we have the functions $\psi_k \in \mathcal{S}(\mathbb{R}^n)$ and $\Psi_k = \hat{\psi}_k \in \mathcal{S}(\mathbb{R}^n)$ for all $k \in \mathbb{N}_0$.

Theorem 13. *Let $(w_k)_{k \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha$, $a > 0$ and $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$. For some $\varepsilon > 0$ we assume $\psi_0, \psi_1 \in \mathcal{S}(\mathbb{R}^n)$ with*

$$|\psi_0| > 0 \quad \text{on} \quad \{x \in \mathbb{R}^n : |x| < \varepsilon\}, \quad (34)$$

$$|\psi_1| > 0 \quad \text{on} \quad \{x \in \mathbb{R}^n : \varepsilon/2 < |x| < 2\varepsilon\}. \quad (35)$$

For $a > \frac{n+c_{\log}(1/q)}{p^-} + \alpha$

$$\|(\Psi_k^* f)_a w_k\|_{\ell_{q(\cdot)}(L_{p(\cdot)})} \leq c \|(\Psi_k * f) w_k\|_{\ell_{q(\cdot)}(L_{p(\cdot)})} \quad (36)$$

holds for all $f \in \mathcal{S}'(\mathbb{R}^n)$.

Remark 6. Observe that in the theorem above no moment conditions on ψ_1 are stated but this time there are restrictions on a and $p(\cdot), q(\cdot)$.

Proof. As in the last proof we find the functions $(\lambda_j)_{j \in \mathbb{N}_0}$ with the properties (27), (28) and

$$\sum_{k=0}^{\infty} \lambda_k(2^{-\nu}x) \psi_k(2^{-\nu}x) = 1 \quad \text{for all } \nu \in \mathbb{N}_0. \quad (37)$$

Instead of (29) we get the identity

$$\Psi_\nu * f = \sum_{k=0}^{\infty} \Lambda_{k,\nu} * \Psi_{k,\nu} * \Psi_\nu * f, \quad (38)$$

where

$$\Lambda_{k,\nu}(\xi) = [\lambda_k(2^{-\nu}\cdot)]^\wedge(\xi) = 2^{\nu n} \Lambda_k(2^\nu \xi) \quad \text{for all } \nu, k \in \mathbb{N}_0.$$

The $\Psi_{k,\nu}$ are defined similarly. For $k \geq 1$ and $\nu \in \mathbb{N}_0$ we have $\Psi_{k,\nu} = \Psi_{k+\nu}$ and with the notation

$$\sigma_{k,\nu}(x) = \begin{cases} \psi_0(2^{-\nu}x) & \text{if } k = 0, \\ \psi_\nu(x) & \text{otherwise} \end{cases}$$

we get $\psi_k(2^{-\nu}x) \psi_\nu(x) = \sigma_{k,\nu}(x) \psi_{k+\nu}(x)$. Hence, we can rewrite (38) as

$$\Psi_\nu * f = \sum_{k=0}^{\infty} \Lambda_{k,\nu} * \hat{\sigma}_{k,\nu} * \Psi_{k+\nu} * f. \quad (39)$$

For $k \geq 1$ we get from Lemma 7

$$|(\Lambda_{k,\nu} * \hat{\sigma}_{k,\nu})(z)| = 2^{\nu n} |(\Lambda_k * \Psi)(2^\nu z)| \leq C_M 2^{\nu n} \frac{2^{-kM}}{(1 + |2^\nu z|^a)} \quad (40)$$

for all $k, \nu \in \mathbb{N}_0$ and arbitrary large $M \in \mathbb{N}$. For $k = 0$ we get the estimate (40) by using Lemma 7 with $M = 0$. This together with (39) gives us

$$|(\Psi_\nu * f)(y)| \leq C_M 2^{\nu n} \sum_{k=0}^{\infty} \int_{\mathbb{R}^n} \frac{2^{-kM}}{(1 + |2^\nu(y-z)|^a)} |(\Psi_{k+\nu} * f)(z)| dz. \quad (41)$$

For fixed $r \in (0, 1]$ we divide both sides of (41) by $(1 + |2^\nu(x-y)|^a)$ and we take the supremum with respect to $y \in \mathbb{R}^n$. Using the inequalities

$$(1 + |2^\nu(y-z)|^a)(1 + |2^\nu(x-y)|^a) \geq c(1 + |2^\nu(x-z)|^a),$$

$$|(\Psi_{k+\nu} * f)(z)| \leq |(\Psi_{k+\nu} * f)(z)|^r (\Psi_{k+\nu}^* f)_a(x)^{1-r} (1 + |2^{k+\nu}(x-z)|^a)^{1-r}$$

and

$$\frac{(1 + |2^{k+\nu}(x-z)|^a)^{1-r}}{(1 + |2^\nu(x-z)|^a)} \leq \frac{2^{ka}}{(1 + |2^{k+\nu}(x-z)|^a)^r},$$

we get

$$(\Psi_\nu^* f)_a(x) \leq C_M \sum_{k=0}^{\infty} 2^{-k(M+n-a)} (\Psi_{k+\nu}^* f)_a(x)^{1-r} \int_{\mathbb{R}^n} \frac{2^{(k+\nu)n} |(\Psi_{k+\nu} * f)(z)|^r}{(1 + |2^{k+\nu}(x-z)|^a)^r} dz. \quad (42)$$

Now, we apply Lemma 11 with

$$\gamma_\nu = (\Psi_\nu^* f)_a(x), \quad \beta_\nu = \int_{\mathbb{R}^n} \frac{2^{\nu n} |(\Psi_\nu * f)(z)|^r}{(1 + |2^\nu(x-z)|^a)^r} dz, \quad \nu \in \mathbb{N}_0$$

$N = M + n - a$, $C_N = C_M + n - a$ and N^0 in (19) equals the order of the distribution $f \in \mathcal{S}'(\mathbb{R}^n)$.

By Lemma 11 we obtain for every $N \in \mathbb{N}$, $x \in \mathbb{R}^n$ and $\nu \in \mathbb{N}_0$

$$(\Psi_\nu^* f)_a(x)^r \leq C_N \sum_{k=0}^{\infty} 2^{-kNr} \int_{\mathbb{R}^n} \frac{2^{(k+\nu)n} |(\Psi_{k+\nu} * f)(z)|^r}{(1 + |2^{k+\nu}(x-z)|^a)^r} dz \quad (43)$$

provided that $(\Psi_\nu^* f)_a(x) < \infty$.

Since $f \in \mathcal{S}'(\mathbb{R}^n)$, we see that $(\Psi_\nu^* f)_a(x) < \infty$ for all $x \in \mathbb{R}^n$ and all $\nu \in \mathbb{N}_0$ at least if $a > N^0$, where N^0 is the order of the distribution. Thus we have (43) with C_N independent of $f \in \mathcal{S}'(\mathbb{R}^n)$ for $a \geq N^0$ and therefore with $C_N = C_{N,f}$ for all $a > 0$ (the right side of (43) decreases as a increases). One can easily check that (43) with $C_N = C_{N,f}$ implies that if for some $a > 0$ the right side of (43) is finite, then $(\Psi_\nu^* f)_a(x) < \infty$. Now, repeating the above argument resurrects the independence of C_N . If the right side of (43) is infinite, there is nothing to prove. More exhaustive arguments of this type have been used in [54] and [47].

We point out that (43) holds also for $r > 1$, where the proof is much simpler. We only have to take (41) with $a + n$ instead of a , divide both sides by $(1 + |2^\nu(x-y)|^a)$ and apply Hölder's inequality with respect to k and then z .

Multiplying (43) by $w_\nu(x)^r$ we derive with the properties of our weight sequence

$$(\Psi_\nu^* f)_a(x)^r w_\nu(x)^r \leq C'_N \sum_{k=0}^{\infty} 2^{-k(N+\alpha_1)r} \int_{\mathbb{R}^n} \frac{2^{(k+\nu)n} |(\Psi_{k+\nu} * f)(z)|^r w_{k+\nu}(z)^r}{(1 + |2^{k+\nu}(x-z)|^{a-\alpha})^r} dz, \quad (44)$$

for all $x \in \mathbb{R}^n$, $\nu \in \mathbb{N}_0$ and all $N \in \mathbb{N}$.

Now, we choose $r = p^-$ and we have $r(a - \alpha) > n + c_{\log}(1/q)$. We denote $g_{k+\nu}^r(z) = |(\Psi_{k+\nu} * f)(z)|^r w_{k+\nu}(z)^r$ then we can rewrite (44) by

$$(\Psi_\nu^* f)_a(x)^r w_\nu(x)^r \leq C'_N \sum_{l=\nu}^{\infty} 2^{-(l-\nu)(N+\alpha_1)r} (g_l^r * \eta_{l,r(a-\alpha)})(x). \quad (45)$$

For fixed $N > 0$ with $\delta = N + \alpha_1 > 0$ we apply the $\ell_{\frac{q(\cdot)}{r}}(L_{\frac{p(\cdot)}{r}})$ norm and derive from (45)

$$\left\| (\Psi_\nu^* f)_a(x)^r w_\nu(x)^r \right\|_{\ell_{\frac{q(\cdot)}{r}}(L_{\frac{p(\cdot)}{r}})} \leq C_N \left\| \sum_{l=\nu}^{\infty} 2^{-(l-\nu)\delta} (g_l^r * \eta_{l,r(a-\alpha)}) \right\|_{\ell_{\frac{q(\cdot)}{r}}(L_{\frac{p(\cdot)}{r}})}.$$

Now application of Lemma 9 and Lemma 10 ($r(a-\alpha) > n + c_{\log}(1/q)$) on the formula above give us

$$\left\| (\Psi_k^* f)_a^r(\cdot) w_k^r(\cdot) \Big| \ell_{q(\cdot)/r}(L_{p(\cdot)/r}) \right\| \leq C'_N \left\| |(\Psi_\nu * f)(\cdot)|^r w_\nu(\cdot)^r \Big| \ell_{q(\cdot)/r}(L_{p(\cdot)/r}) \right\|$$

which proves the theorem. \square

3.2 Local means characterization of $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$

In this section we reformulate the local means characterization for $B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ from above and for $F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ from Corollary 4.7 in [29] in terms of variable smoothness. If we have a variable smoothness function $s \in C_{loc}^{\log}(\mathbb{R}^n)$ given, then $w_j(x) = 2^{js(x)}$ defines an admissible weight sequence $\mathbf{w} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha$ with $\alpha_1 = s^-$, $\alpha_2 = s^+$ and $\alpha = c_{\log}(s)$, cf. Remark 2. Here, we denote by $c_{\log}(s)$ the constant in (4) for $s(\cdot)$.

Theorem 14. *Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ ($p^+, q^+ < \infty$ in the F -case) and $s \in C_{loc}^{\log}(\mathbb{R}^n)$. Further let $a > 0$, $R \in \mathbb{N}_0$ with $R > s^+$ and let ψ_0, ψ_1 belong to $\mathcal{S}(\mathbb{R}^n)$ with*

$$D^\beta \psi_1(0) = 0, \quad \text{for } 0 \leq |\beta| < R, \quad (46)$$

and

$$|\psi_0(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : |x| < \varepsilon\} \quad (47)$$

$$|\psi_1(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : \varepsilon/2 < |x| < 2\varepsilon\} \quad (48)$$

for some $\varepsilon > 0$.

1. For $a > \frac{n+c_{\log}(1/q)}{p^-} + c_{\log}(s)$ and all $f \in \mathcal{S}'(\mathbb{R}^n)$ we have

$$\left\| f \Big| B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \right\| \sim \left\| 2^{ks(\cdot)}(\Psi_k * f) \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\| \sim \left\| 2^{ks(\cdot)}(\Psi_k^* f)_a \Big| \ell_{q(\cdot)}(L_{p(\cdot)}) \right\|.$$

2. For $a > \frac{n}{\min(p^-, q^-)} + c_{\log}(s)$ and all $f \in \mathcal{S}'(\mathbb{R}^n)$ we have

$$\left\| f \Big| F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \right\| \sim \left\| 2^{ks(\cdot)}(\Psi_k * f) \Big| L_{p(\cdot)}(\ell_{q(\cdot)}) \right\| \sim \left\| 2^{ks(\cdot)}(\Psi_k^* f)_a \Big| L_{p(\cdot)}(\ell_{q(\cdot)}) \right\|.$$

4 Ball means of differences

This section is devoted to the characterization of Besov and Triebel-Lizorkin spaces $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ by ball means of differences. In the case of constant indices p, q and s , this is a classical part of the theory of function spaces. We refer especially to [51, Section 2.5] and references given there. It turns out that, under the restriction

$$s > \sigma_p = n \left(\frac{1}{\min(p, 1)} - 1 \right) \quad (49)$$

in the B -case and

$$s > \sigma_{p,q} = n \left(\frac{1}{\min(p, q, 1)} - 1 \right) \quad (50)$$

in the F -case, Besov and Triebel-Lizorkin spaces with constant indices may be characterized by expressions involving only the differences of the function values without any use of Fourier analysis. This was complemented in [48] and [49] by showing that these conditions are also indispensable. Of course, we are limited by (49) and (50) also in the case of variable exponents.

The characterization by (local means of) differences for 2-microlocal spaces with constant $p, q > 1$ was given by Besov [8] and a similar characterization for Besov spaces with $p = q = \infty$ and the special weight sequence from (6) was given by Seuret and Levy Véhél in [34]. We refer to [18] and [28] for the treatment of spaces of generalized smoothness.

Our approach follows essentially [51] with some modifications described in [53]. The main obstacle on this way is the unboundedness of the maximal operator in the frame of $L_{p(\cdot)}(\ell_{q(\cdot)})$ and $\ell_{q(\cdot)}(L_{p(\cdot)})$ spaces, cf. [17, Section 5] and [1, Example 4.1]. This is circumvented by the use of convolution with radial functions in the sense of [17] and [1] together with a certain bootstrapping argument, which shall be described in detail below.

The plan of this part of the work is as follows. First we give in Section 4.1 the necessary notation. We state the main assertions of this part in Section 4.2. Then we prove in Section 4.3 a certain preliminary version of these assertions. In Section 4.4 we prove a characterization by ball means of differences for spaces with $q \in (0, \infty]$ constant (where the maximal operator is bounded) and use this together with our preliminary characterization from Section 4.3 to conclude the proof. Finally, in Section 4.5 we will present the ball means of differences characterization also for the 2-microlocal function spaces $B_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ and $F_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ and in Section 4.6 we present separately some useful Lemmas, not to disturb the main proofs of this part.

4.1 Notation

Let f be a function on \mathbb{R}^n and let $h \in \mathbb{R}^n$. Then we define

$$\Delta_h^1 f(x) = f(x+h) - f(x), \quad x \in \mathbb{R}^n.$$

The higher order differences are defined inductively by

$$\Delta_h^M f(x) = \Delta_h^1(\Delta_h^{M-1} f)(x), \quad M = 2, 3, \dots$$

This definition also allows a direct formula

$$\Delta_h^M f(x) := \sum_{j=0}^M (-1)^j \binom{M}{j} f(x + (M-j)h). \quad (51)$$

By *ball means of differences* we mean the quantity

$$d_t^M f(x) = t^{-n} \int_{|h| \leq t} |\Delta_h^M f(x)| dh = \int_B |\Delta_{th}^M f(x)| dh,$$

where $B = \{y \in \mathbb{R}^n : |y| < 1\}$ is the unit ball of \mathbb{R}^n , $t > 0$ is a real number and M is a natural number.

Let us now introduce the (quasi-)norms, which shall be the main subject of our study. We define

$$\begin{aligned} \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^* &:= \|f|L_{p(\cdot)}(\mathbb{R}^n)\| \\ &+ \left\| \left(\int_0^\infty t^{-s(x)q(x)} (d_t^M f(x))^{q(x)} \frac{dt}{t} \right)^{1/q(x)} \Big|_{L_{p(\cdot)}(\mathbb{R}^n)} \right\| \end{aligned} \quad (52)$$

and its partially discretized counterpart

$$\begin{aligned} \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**} &:= \|f|L_{p(\cdot)}(\mathbb{R}^n)\| \\ &+ \left\| \left(\sum_{k=-\infty}^\infty 2^{ks(x)q(x)} (d_{2^{-k}}^M f(x))^{q(x)} \right)^{1/q(x)} \Big|_{L_{p(\cdot)}(\mathbb{R}^n)} \right\| \\ &= \|f|L_{p(\cdot)}(\mathbb{R}^n)\| + \left\| \left(2^{ks(x)} d_{2^{-k}}^M f(x) \right)_{k=-\infty}^\infty \Big|_{L_{p(\cdot)}(\ell_{q(\cdot)})} \right\|. \end{aligned} \quad (53)$$

The norm $\|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**}$ admits a direct counterpart also for Besov spaces, namely

$$\|f|B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**} := \|f|L_{p(\cdot)}(\mathbb{R}^n)\| + \left\| \left(2^{ks(x)} d_{2^{-k}}^M f(x) \right)_{k=-\infty}^\infty \Big|_{\ell_{q(\cdot)}(L_{p(\cdot)})} \right\|. \quad (54)$$

Finally, we shall use as a technical tool also the analogues of (52)–(54) with the integration over t restricted to $0 < t < 1$. This leads to the following expressions.

$$\begin{aligned} \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|_1^* &:= \|f|L_{p(\cdot)}(\mathbb{R}^n)\| \\ &+ \left\| \left(\int_0^1 t^{-s(x)q(x)} (d_t^M f(x))^{q(x)} \frac{dt}{t} \right)^{1/q(x)} \Big|_{L_{p(\cdot)}(\mathbb{R}^n)} \right\|, \end{aligned} \quad (55)$$

$$\begin{aligned} \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|_1^{**} &:= \|f|L_{p(\cdot)}(\mathbb{R}^n)\| \\ &+ \left\| \left(\sum_{k=0}^\infty 2^{ks(x)q(x)} (d_{2^{-k}}^M f(x))^{q(x)} \right)^{1/q(x)} \Big|_{L_{p(\cdot)}(\mathbb{R}^n)} \right\| \\ &= \|f|L_{p(\cdot)}(\mathbb{R}^n)\| + \left\| \left(2^{ks(x)} d_{2^{-k}}^M f(x) \right)_{k=0}^\infty \Big|_{L_{p(\cdot)}(\ell_{q(\cdot)})} \right\|, \end{aligned} \quad (56)$$

$$\|f|B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|_1^{**} := \|f|L_{p(\cdot)}(\mathbb{R}^n)\| + \left\| \left(2^{ks(x)} d_{2^{-k}}^M f(x) \right)_{k=0}^\infty \Big|_{\ell_{q(\cdot)}(L_{p(\cdot)})} \right\|, \quad (57)$$

where now $k \in \mathbb{N}_0$.

4.2 Main Theorem

Using the notation introduced above, we may now state the main result of this section.

Theorem 15. (i) Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $p^+, q^+ < \infty$ and $s \in C_{loc}^{\log}(\mathbb{R}^n)$. Let $M \in \mathbb{N}$ with $M > s^+$ and let

$$s^- > \sigma_{p^-, q^-} \cdot \left[1 + \frac{c_{\log}(s)}{n} \cdot \min(p^-, q^-) \right]. \quad (58)$$

Then

$$F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) = \{f \in L_{p(\cdot)}(\mathbb{R}^n) \cap \mathcal{S}'(\mathbb{R}^n) : \|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^* < \infty\}$$

and $\|\cdot|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|$ and $\|\cdot|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^*$ are equivalent on $F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$. The same holds for $\|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**}$.

(ii) Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and $s \in C_{loc}^{\log}(\mathbb{R}^n)$. Let $M \in \mathbb{N}$ with $M > s^+$ and let

$$s^- > \sigma_{p^-} \cdot \left[1 + \frac{c_{\log}(1/q)}{n} + \frac{c_{\log}(s)}{n} \cdot p^- \right]. \quad (59)$$

Then

$$B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) = \{f \in L_{p(\cdot)}(\mathbb{R}^n) \cap \mathcal{S}'(\mathbb{R}^n) : \|f|B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**} < \infty\}$$

and $\|\cdot|B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|$ and $\|\cdot|B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**}$ are equivalent on $B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$.

Remark 7. Let us comment on the rather technical conditions (58) and (59).

- If $\min(p^-, q^-) \geq 1$, then (58) becomes just $s^- > 0$. Furthermore, if p, q and s are constant functions, then (58) coincides with (50).
- If $p^- \geq 1$, then (59) reduces also to $s^- > 0$ and in the case of constant exponents we again recover (49).

As indicated already above, the proof is divided into several parts.

4.3 Preliminary version of Theorem 15

This subsection contains a preliminary version of Theorem 15 (Lemma 16). Its proof represents the heart of the proof of Theorem 15. For better lucidity, it is again divided into more parts.

Lemma 16. *Under the conditions of Theorem 15, the following estimates hold for all $f \in L_{p(\cdot)}(\mathbb{R}^n) \cap \mathcal{S}'(\mathbb{R}^n)$:*

$$\|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^* \approx \|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**}, \quad (60)$$

$$\|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|_1^* \approx \|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|_1^{**}, \quad (61)$$

$$\|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|_1^{**} \lesssim \|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\| \lesssim \|f|F_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**}, \quad (62)$$

$$\|f|B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|_1^{**} \lesssim \|f|B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\| \lesssim \|f|B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**}. \quad (63)$$

Proof. Part I. First we prove (60) and (61). We discretize the inner part of $\|\cdot\|_*$ and obtain

$$\begin{aligned}
& \left[\int_0^\infty t^{-s(x)q(x)} \left(\int_B |\Delta_{th}^M f(x)| dh \right)^{q(x)} \frac{dt}{t} \right]^{1/q(x)} \\
&= \left[\int_0^\infty t^{-s(x)q(x)} \left(t^{-n} \int_{tB} |\Delta_{\mathcal{X}}^M f(x)| d\mathcal{X} \right)^{q(x)} \frac{dt}{t} \right]^{1/q(x)} \\
&= \left[\sum_{k=-\infty}^\infty \int_{2^{-k-1}B}^{2^{-k}B} t^{-s(x)q(x)} \left(t^{-n} \int_{tB} |\Delta_{\mathcal{X}}^M f(x)| d\mathcal{X} \right)^{q(x)} \frac{dt}{t} \right]^{1/q(x)}.
\end{aligned} \tag{64}$$

If $2^{-k-1} \leq t \leq 2^{-k}$, then $2^{ks(x)q(x)} \leq t^{-s(x)q(x)} \leq 2^{(k+1)s(x)q(x)}$ and

$$2^{kn} \int_{2^{-(k+1)}B} |\Delta_{\mathcal{X}}^M f(x)| d\mathcal{X} \lesssim t^{-n} \int_{tB} |\Delta_{\mathcal{X}}^M f(x)| d\mathcal{X} \lesssim 2^{(k+1)n} \int_{2^{-k}B} |\Delta_{\mathcal{X}}^M f(x)| d\mathcal{X}.$$

Plugging these estimates into (64), we may further estimate

$$\begin{aligned}
& \left[\int_0^\infty t^{-s(x)q(x)} \left(\int_B |\Delta_{th}^M f(x)| dh \right)^{q(x)} \frac{dt}{t} \right]^{1/q(x)} \\
& \lesssim \left[\sum_{k=-\infty}^\infty 2^{(k+1)s(x)q(x)} \left(2^{kn} \int_{2^{-k}B} |\Delta_{\mathcal{X}}^M f(x)| d\mathcal{X} \right)^{q(x)} \right]^{1/q(x)} \\
& \lesssim \left[\sum_{k=-\infty}^\infty 2^{ks(x)q(x)} \left(\int_B |\Delta_{2^{-k}\mathcal{X}}^M f(x)| d\mathcal{X} \right)^{q(x)} \right]^{1/q(x)}.
\end{aligned}$$

The estimate from below follows in the same manner. Finally, the proof of (61) is almost the same.

Part II. This part is devoted to the proof of the left hand side of (62). It is divided into several steps to make the presentation more clear.

Step 1. First, we point out that the estimate

$$\|f\|_{L_{p(\cdot)}(\mathbb{R}^n)} \lesssim \|f\|_{B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)}$$

follows from the characterization of $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ in terms of Nikol'skij representations (cf. Theorem 8.1 of [1]). We refer also to Remark 2.5.3/1 in [51]. The extension to F -spaces is then given by simple embedding

$$\|f\|_{L_{p(\cdot)}(\mathbb{R}^n)} \lesssim \|f\|_{B_{p(\cdot),p(\cdot)}^{s(\cdot)-\varepsilon}(\mathbb{R}^n)} \lesssim \|f\|_{F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)}$$

with $\varepsilon > 0$ chosen small enough.

Step 2. Let $(\varphi_j)_{j \in \mathbb{N}_0}$ be the functions used in Definition 3. We use the decomposition

$$f = \sum_{l=-\infty}^\infty f_{(k+l)}, \quad k \in \mathbb{Z},$$

where $f_{(k+l)} = (\varphi_{k+l}\hat{f})^\vee$, or $= 0$ if $k+l < 0$ and get

$$\begin{aligned} (\clubsuit) &:= \sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(\int_B |\Delta_{2^{-k}h}^M f(x)| dh \right)^{q(x)} \\ &= \sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(\int_B |\Delta_{2^{-k}h}^M \left(\sum_{l=-\infty}^{\infty} f_{(k+l)} \right)(x)| dh \right)^{q(x)}. \end{aligned}$$

If $q(x) \leq 1$ then we proceed further

$$\begin{aligned} (\clubsuit) &\leq \sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(\int_B \sum_{l=-\infty}^{\infty} |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \right)^{q(x)} \\ &\leq \sum_{k=0}^{\infty} \sum_{l=-\infty}^{\infty} 2^{ks(x)q(x)} \left(\int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \right)^{q(x)}. \end{aligned}$$

If $q(x) > 1$, we use Minkowski's inequality

$$\begin{aligned} (\clubsuit)^{1/q(x)} &\leq \left(\sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(\int_B \sum_{l=-\infty}^{\infty} |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \right)^{q(x)} \right)^{1/q(x)} \\ &\leq \sum_{l=-\infty}^{\infty} \left(\sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(\int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \right)^{q(x)} \right)^{1/q(x)}. \end{aligned}$$

We split in both cases

$$\sum_{l=-\infty}^{\infty} \dots = I + II = \sum_{l=-\infty}^0 \dots + \sum_{l=1}^{\infty} \dots \quad (65)$$

Step 3. We estimate the first summand with $l \leq 0$.

We use Lemma 22 in the form

$$|\Delta_h^M f_{(k+l)}(x)| \leq C \max(1, |bh|^a) \cdot \min(1, |bh|^M) P_{b,a} f_{(k+l)}(x),$$

where $a > 0$ is arbitrary, $b = 2^{k+l}$ and

$$P_{b,a} f(x) = \sup_{z \in \mathbb{R}^n} \frac{|f(x-z)|}{1 + |bz|^a}.$$

Furthermore, we use this estimate with $2^{-k}h$ instead of h . We obtain

$$\begin{aligned} \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh &\lesssim \int_B \max(1, |b2^{-k}h|^a) \cdot \min(1, |b2^{-k}h|^M) P_{b,a} f_{(k+l)}(x) dh \\ &\lesssim 2^{lM} P_{2^{k+l},a} f_{(k+l)}(x). \end{aligned} \quad (66)$$

The last inequality follows from $\max(1, |b2^{-k}h|^a) \leq 1$ (recall that $l \leq 0$ and $|h| \leq 1$) and $\min(1, |b2^{-k}h|^M) \leq 2^{lM}$.

If $q(x) \leq 1$, we estimate the first sum in (65)

$$\begin{aligned}
I &\leq \sum_{l=-\infty}^0 \sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(\int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \right)^{q(x)} \\
&\lesssim \sum_{l=-\infty}^0 \sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(2^{lM} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{q(x)} \\
&= \sum_{l=-\infty}^0 2^{l(M-s(x))q(x)} \sum_{k=0}^{\infty} 2^{(k+l)s(x)q(x)} P_{2^{k+l},a}^{q(x)} f_{(k+l)}(x) \\
&\approx \sum_{k=0}^{\infty} 2^{ks(x)q(x)} P_{2^k,a}^{q(x)} f_{(k)}(x),
\end{aligned}$$

where the last estimate makes use of $M > s^+$, $q^- > 0$ and the fact that $f_{(k+l)} = 0$ for $k+l < 0$.

If $q(x) > 1$, we proceed in a similar way to obtain

$$\begin{aligned}
I^{1/q(x)} &\leq \sum_{l=-\infty}^0 \left(\sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(\int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \right)^{q(x)} \right)^{1/q(x)} \\
&\lesssim \sum_{l=-\infty}^0 \left(\sum_{k=0}^{\infty} 2^{ks(x)q(x)} \left(2^{lM} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{q(x)} \right)^{1/q(x)} \\
&= \sum_{l=-\infty}^0 2^{l(M-s(x))} \left(\sum_{k=0}^{\infty} 2^{(k+l)s(x)q(x)} P_{2^{k+l},a}^{q(x)} f_{(k+l)}(x) \right)^{1/q(x)} \\
&\lesssim \left(\sum_{k=0}^{\infty} 2^{ks(x)q(x)} P_{2^k,a}^{q(x)} f_{(k)}(x) \right)^{1/q(x)}.
\end{aligned}$$

We have used in the last estimate again $M > s^+$ and the definition of $f_{(k+l)}$.

Hence,

$$I^{1/q(x)} \lesssim \left(\sum_{k=0}^{\infty} 2^{ks(x)q(x)} P_{2^k,a}^{q(x)} f_{(k)}(x) \right)^{1/q(x)}$$

holds for all $x \in \mathbb{R}^n$.

Finally, we obtain

$$\begin{aligned}
\|I^{1/q(\cdot)}|_{L_{p(\cdot)}(\mathbb{R}^n)}\| &\lesssim \left\| \left(\sum_{k=0}^{\infty} 2^{ks(x)q(x)} P_{2^k,a}^{q(x)} f_{(k)}(x) \right)^{1/q(x)} \right\|_{L_{p(\cdot)}(\mathbb{R}^n)} \\
&= \left\| \left(2^{ks(x)} P_{2^k,a} f_{(k)}(x) \right)_{k=0}^{\infty} \right\|_{L_{p(\cdot)}(\ell_{q(\cdot)})} \\
&\lesssim \left\| \left(2^{ks(\cdot)} f_{(k)} \right)_{k=0}^{\infty} \right\|_{L_{p(\cdot)}(\ell_{q(\cdot)})} = \|f\|_{F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)},
\end{aligned} \tag{67}$$

where we used the boundedness of Peetre maximal operator as described in Theorem 14 for $a > 0$ large enough.

Step 4. We estimate the second summand in (65) with $l > 0$. If $\min(p^-, q^-) > 1$, then we put $\lambda = 1$. Otherwise we choose real parameters $0 < \lambda < \min(p^-, q^-)$ and $a > 0$ such that

$$a > \frac{n}{\min(p^-, q^-)} + c_{\log}(s)$$

and $a(1 - \lambda) < s^-$. Due to (58), this is always possible.

We start again with estimates of the ball means of differences. We use Lemma 22 and (51) and obtain

$$\begin{aligned} \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh &= \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)|^\lambda \cdot |\Delta_{2^{-k}h}^M f_{(k+l)}(x)|^{1-\lambda} dh \\ &\lesssim \int_B \left(\max(1, |2^{k+l}2^{-k}h|^a) \min(1, |2^{k+l}2^{-k}h|^M) P_{2^{k+l}, a} f_{(k+l)}(x) \right)^{1-\lambda} \\ &\quad \cdot |\Delta_{2^{-k}h}^M f_{(k+l)}(x)|^\lambda dh \\ &\leq \left(2^{la} P_{2^{k+l}, a} f_{(k+l)}(x) \right)^{1-\lambda} \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)|^\lambda dh \\ &\leq \left(2^{la} P_{2^{k+l}, a} f_{(k+l)}(x) \right)^{1-\lambda} \sum_{j=0}^M c_{j, M} \int_B |f_{(k+l)}(x + j2^{-k}h)|^\lambda dh, \end{aligned} \tag{68}$$

where the constants $c_{j, M}$ are given by (51).

We shall deal in detail only with term with $j = 1$. The term with $j = 0$ is much simpler to handle (as there the integration over $h \in B$ immediately disappears) and this case reduces essentially to Hölder's inequality and boundedness of the Peetre maximal operator. The terms with $2 \leq j \leq M$ may be handled in the same way as the one with $j = 1$.

We use Lemma 20 with $r = \lambda$ in the form

$$|f_{(k+l)}(y)|^\lambda \lesssim (\eta_{k+l, 2m} * |f_{(k+l)}|^\lambda)(y),$$

with $m > \max(n, c_{\log}(s))$, Lemma 23 and Lemma 19 to get

$$\begin{aligned} 2^{ks(x)\lambda} \int_B |f_{(k+l)}(x + 2^{-k}h)|^\lambda dh &\lesssim 2^{ks(x)\lambda} \int_B (\eta_{k+l, 2m} * |f_{(k+l)}|^\lambda)(x + 2^{-k}h) dh \\ &= 2^{ks(x)\lambda} ([2^{kn} \chi_{2^{-k}B}] * \eta_{k+l, 2m} * |f_{(k+l)}|^\lambda)(x) \\ &\lesssim 2^{ks(x)\lambda} (\eta_{k, 2m} * |f_{(k+l)}|^\lambda)(x) \\ &\lesssim (\eta_{k, m} * |2^{ks(\cdot)} f_{(k+l)}|^\lambda)(x) \\ &\leq 2^{-ls^- \lambda} (\eta_{k, m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^\lambda)(x). \end{aligned} \tag{69}$$

We insert (69) into (68) and arrive at

$$\begin{aligned} 2^{ks(x)} \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh &\lesssim 2^{la(1-\lambda) - ls^-} \left(2^{(k+l)s(x)} P_{2^{k+l}, a} f_{(k+l)}(x) \right)^{1-\lambda} (\eta_{k, m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^\lambda)(x). \end{aligned} \tag{70}$$

If $q(x) > 1$, we proceed further with the use of Hölder's inequality

$$\begin{aligned}
II^{1/q(x)} &\lesssim \sum_{l=1}^{\infty} 2^{la(1-\lambda)-ls^-} \\
&\quad \left(\sum_{k=0}^{\infty} \left(2^{(k+l)s(x)} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{(1-\lambda)q(x)} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^{\lambda})^{q(x)}(x) \right)^{1/q(x)} \\
&\leq \sum_{l=1}^{\infty} 2^{la(1-\lambda)-ls^-} \left(\sum_{k=0}^{\infty} \left(2^{(k+l)s(x)} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{q(x)} \right)^{(1-\lambda)/q(x)} \\
&\quad \cdot \left(\sum_{k=0}^{\infty} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^{\lambda})^{q(x)/\lambda}(x) \right)^{\lambda/q(x)} \\
&= \left(\sum_{k=0}^{\infty} \left(2^{ks(x)} P_{2^k,a} f_{(k)}(x) \right)^{q(x)} \right)^{(1-\lambda)/q(x)} \\
&\quad \cdot \sum_{l=1}^{\infty} 2^{la(1-\lambda)-ls^-} \left(\sum_{k=0}^{\infty} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^{\lambda})^{q(x)/\lambda}(x) \right)^{\lambda/q(x)}.
\end{aligned}$$

If $q(x) \leq 1$, we obtain in a similar way

$$\begin{aligned}
II &\lesssim \sum_{l=1}^{\infty} 2^{(la(1-\lambda)-ls^-)q(x)} \\
&\quad \sum_{k=0}^{\infty} \left(2^{(k+l)s(x)} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{(1-\lambda)q(x)} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^{\lambda})^{q(x)}(x) \\
&\leq \sum_{l=1}^{\infty} 2^{(la(1-\lambda)-ls^-)q(x)} \left(\sum_{k=0}^{\infty} \left(2^{(k+l)s(x)} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{q(x)} \right)^{1-\lambda} \\
&\quad \cdot \left(\sum_{k=0}^{\infty} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^{\lambda})^{q(x)/\lambda}(x) \right)^{\lambda} \\
&= \left(\sum_{k=0}^{\infty} \left(2^{ks(x)} P_{2^k,a} f_{(k)}(x) \right)^{q(x)} \right)^{1-\lambda} \\
&\quad \cdot \sum_{l=1}^{\infty} 2^{(la(1-\lambda)-ls^-)q(x)} \left(\sum_{k=0}^{\infty} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^{\lambda})^{q(x)/\lambda}(x) \right)^{\lambda}
\end{aligned}$$

and further (with use of Lemma 24)

$$\begin{aligned}
II^{1/q(x)} &\lesssim \left(\sum_{k=0}^{\infty} \left(2^{ks(x)} P_{2^k, a} f_{(k)}(x) \right)^{q(x)} \right)^{(1-\lambda)/q(x)} \\
&\quad \cdot \left(\sum_{l=1}^{\infty} 2^{(la(1-\lambda)-ls^-)q(x)} \left(\sum_{k=0}^{\infty} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^\lambda)^{q(x)/\lambda}(x) \right)^\lambda \right)^{1/q(x)} \\
&\lesssim \left(\sum_{k=0}^{\infty} \left(2^{ks(x)} P_{2^k, a} f_{(k)}(x) \right)^{q(x)} \right)^{(1-\lambda)/q(x)} \\
&\quad \cdot \sum_{l=1}^{\infty} 2^{1/2 \cdot (la(1-\lambda)-ls^-)} \left(\sum_{k=0}^{\infty} (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}|^\lambda)^{q(x)/\lambda}(x) \right)^{\lambda/q(x)}.
\end{aligned}$$

If we denote

$$F(x) := \left(\sum_{k=0}^{\infty} \left(2^{ks(x)} P_{2^k, a} f_{(k)}(x) \right)^{q(x)} \right)^{1/q(x)}, \quad x \in \mathbb{R}^n$$

and

$$B_{k+l}(x) := |2^{(k+l)s(x)} f_{(k+l)}(x)|, \quad x \in \mathbb{R}^n$$

we get for $\delta := -1/2 \cdot (a(1-\lambda) - s^-) > 0$

$$II^{1/q(x)} \lesssim F(x)^{1-\lambda} \cdot \sum_{l=1}^{\infty} 2^{-l\delta} \left(\sum_{k=0}^{\infty} (\eta_{k,m} * B_{k+l}^\lambda)^{q(x)/\lambda}(x) \right)^{\lambda/q(x)}. \quad (71)$$

We use $\|F_1^{1-\lambda} F_2^\lambda\|_{p(\cdot)} \leq 2 \|F_1\|_{p(\cdot)}^{1-\lambda} \|F_2\|_{p(\cdot)}^\lambda$, cf. [16, Lemma 3.2.20], and suppose that the $L_{p(\cdot)}$ - (quasi-)norm is equivalent to an r -norm with $0 < r \leq 1$. Together with Lemma 21 we arrive to

$$\begin{aligned}
\|II^{1/q(x)}\|_{p(\cdot)}^r &\lesssim \|F(x)\|_{p(\cdot)}^{(1-\lambda)r} \cdot \left\| \sum_{l=1}^{\infty} 2^{-l\delta} \left(\sum_{k=0}^{\infty} (\eta_{k,m} * B_{k+l}^\lambda)^{q(x)/\lambda}(x) \right)^{1/q(x)} \right\|_{p(\cdot)}^{\lambda r} \\
&\lesssim \|F(x)\|_{p(\cdot)}^{(1-\lambda)r} \cdot \sum_{l=1}^{\infty} 2^{-l\delta r} \left\| \left(\sum_{k=0}^{\infty} (\eta_{k,m} * B_{k+l}^\lambda)^{q(x)/\lambda}(x) \right)^{1/q(x)} \right\|_{p(\cdot)}^{\lambda r} \\
&\lesssim \|f\|_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)^{(1-\lambda)r} \cdot \sum_{l=1}^{\infty} 2^{-l\delta r} \left\| (\eta_{k,m} * B_{k+l}^\lambda(x))_{k=0}^\infty \right\|_{L_{p(\cdot)/\lambda}(\ell_{q(\cdot)/\lambda})}^r \\
&\lesssim \|f\|_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)^{(1-\lambda)r} \cdot \sum_{l=1}^{\infty} 2^{-l\delta r} \left\| (B_{k+l}^\lambda(x))_{k=0}^\infty \right\|_{L_{p(\cdot)/\lambda}(\ell_{q(\cdot)/\lambda})}^r \quad (72) \\
&\lesssim \|f\|_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)^{(1-\lambda)r} \cdot \sum_{l=1}^{\infty} 2^{-l\delta r} \left\| (B_k^\lambda(x))_{k=0}^\infty \right\|_{L_{p(\cdot)/\lambda}(\ell_{q(\cdot)/\lambda})}^r
\end{aligned}$$

$$\begin{aligned}
&\lesssim \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{(1-\lambda)r} \cdot \|B_k(x)\|_{L_{p(\cdot)}(\ell_{q(\cdot)})}^{\lambda r} \\
&\lesssim \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^r,
\end{aligned} \tag{73}$$

which finishes the proof.

Part III.: We prove the right hand side of (62). We follow again essentially [51, Section 2.5.9] with some modifications as presented in [53]. Roughly speaking, compared to the case of constant exponents, only minor modifications are necessary.

Let $\psi \in C_0^\infty(\mathbb{R}^n)$ with $\psi(x) = 1, |x| \leq 1$ and $\psi(x) = 0, |x| > 3/2$. We define

$$\varphi_0(x) = (-1)^{M+1} \sum_{\mu=0}^{M-1} (-1)^\mu \binom{M}{\mu} \psi((M-\mu)x).$$

It follows that $\varphi_0 \in C_0^\infty(\mathbb{R}^n)$ with $\varphi_0(x) = 0, |x| > 3/2$ and $\varphi_0(x) = 1, |x| < 1/M$. We also put $\varphi_j(x) = \varphi_0(2^{-j}x) - \varphi_0(2^{-j+1}x)$ for $j \geq 1$. This is the decomposition of unity we used in the definition of $\|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|$, cf. Definition 3. Recall that due to [17] and [29], this (quasi-)norm of $\|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|$ does not depend on the choice of the decomposition of unity.

We observe that

$$\varphi_0(x) = (-1)^{M+1} (\Delta_x^M \psi(0) - (-1)^M),$$

and

$$(\mathcal{F}^{-1} \varphi_j \mathcal{F} f)(x) = \begin{cases} (\mathcal{F}^{-1} \Delta_\xi^M \psi(0) \mathcal{F} f)(x) + (-1)^{M+1} f(x), & j = 0, \\ (\mathcal{F}^{-1} (\Delta_{2^{-j}\xi}^M \psi(0) - \Delta_{2^{-j+1}\xi}^M \psi(0)) \mathcal{F} f)(x), & j \geq 1. \end{cases} \tag{74}$$

Furthermore, a straightforward calculation shows that

$$\begin{aligned}
|(\mathcal{F}^{-1} (\Delta_{2^{-j}\xi}^M \psi(0)) \mathcal{F} f)(x)| &= \left| \sum_{u=0}^M (-1)^u \mathcal{F}^{-1} [\psi((M-u)2^{-j}\cdot) \mathcal{F} f](x) \right| \\
&\approx \left| \sum_{u=0}^M (-1)^u \mathcal{F}^{-1} [\psi((M-u)2^{-j}\cdot)] * f(x) \right| \\
&\approx \left| \sum_{u=0}^M (-1)^u \int_{\mathbb{R}^n} \mathcal{F}^{-1} \psi(h) f(x - (M-u)2^{-j}h) dh \right| \\
&= \left| \int_{\mathbb{R}^n} \hat{\psi}(h) \Delta_{2^{-j}h}^M f(x) dh \right| \leq \int_{\mathbb{R}^n} |\hat{\psi}(h)| \cdot |\Delta_{2^{-j}h}^M f(x)| dh
\end{aligned} \tag{75}$$

holds for every $j \in \mathbb{N}_0$. We denote $g = \hat{\psi} \in \mathcal{S}(\mathbb{R}^n)$ and obtain

$$\begin{aligned}
\|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\| &\approx \|2^{js(x)} (\mathcal{F}^{-1} \varphi_j \mathcal{F} f)(x)\|_{L_{p(\cdot)}(\ell_{q(\cdot)})} \\
&\lesssim \|f|L_{p(\cdot)}(\mathbb{R}^n)\| + \left\| 2^{js(x)} \int_{\mathbb{R}^n} |g(h)| \cdot |\Delta_{2^{-j}h}^M f(x)| dh \right\|_{L_{p(\cdot)}(\ell_{q(\cdot)})}.
\end{aligned} \tag{76}$$

The rest of this part consists essentially of using the property of $g \in \mathcal{S}(\mathbb{R}^n)$ to come from (76) to $\|\cdot\|^{**}$.

We denote

$$I_0 := B, \quad I_u := 2^u B \setminus 2^{u-1} B, \quad u \in \mathbb{N}$$

and use $|g(h)| \leq c2^{-ur}$, $h \in I_u$ with r taken large enough (recall that $g \in \mathcal{S}(\mathbb{R}^n)$) and estimate

$$\begin{aligned} \int_{\mathbb{R}^n} |g(h)| \cdot |\Delta_{2^{-j}h}^M f(x)| dh &= \sum_{u=0}^{\infty} \int_{I_u} |g(h)| \cdot |\Delta_{2^{-j}h}^M f(x)| dh \\ &\lesssim \sum_{u=0}^{\infty} 2^{-ur} 2^{jn} \int_{2^{u-j} B} |\Delta_h^M f(x)| dh \\ &= \sum_{u=0}^{\infty} 2^{u(n-r)} 2^{-(u-j)n} \int_{2^{u-j} B} |\Delta_h^M f(x)| dh. \end{aligned} \quad (77)$$

We put

$$G_j(x) := 2^{js(x)} |(\mathcal{F}^{-1}(\Delta_{2^{-j}\xi}^M \psi(0)) \mathcal{F} f)(x)|, \quad j \in \mathbb{N}_0$$

and

$$g_k(x) := 2^{ks(x)} 2^{kn} \int_{2^{-k} B} |\Delta_h^M f(x)| dh, \quad k \in \mathbb{Z}.$$

Using (74), (75) and (77), we obtain the estimate

$$\begin{aligned} G_j(x) &\lesssim 2^{js(x)} \sum_{u=0}^{\infty} 2^{u(n-r)} 2^{-(u-j)n} \int_{2^{u-j} B} |\Delta_h^M f(x)| dh \\ &= \sum_{k=-\infty}^j 2^{(j-k)s(x)} 2^{(j-k)(n-r)} 2^{ks(x)} 2^{kn} \int_{2^{-k} B} |\Delta_h^M f(x)| dh \\ &= \sum_{k=-\infty}^j 2^{(j-k)(s(x)+n-r)} g_k(x) \leq \sum_{k=-\infty}^{\infty} 2^{|j-k| \cdot (s(x)+n-r)} g_k(x). \end{aligned} \quad (78)$$

Choosing $r > s^+ + n$ and applying Lemma 9 then finishes the proof.

Part IV. The proof of the left hand side of (63) follows in the same manner as in Part II. We shall describe the necessary modifications. First, let us mention, that the condition $q^+ < \infty$ was used only in the application of Lemma 21. In the rest of the arguments also the case $q(x) = \infty$ may be incorporated with only slight change of notation.

Let us put

$$f^{(k)}(x) := 2^{ks(x)} \int_B |\Delta_{2^{-k}h}^M f(x)| dh, \quad x \in \mathbb{R}^n.$$

We obtain (in analogue to (65))

$$\begin{aligned} f^{(k)} &\leq f^{(k),I} + f^{(k),II} := \sum_{l=-\infty}^0 2^{ks(x)} \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \\ &\quad + \sum_{l=1}^{\infty} 2^{ks(x)} \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh. \end{aligned}$$

We estimate the first sum using (66) and get

$$f^{(k),I} \lesssim \sum_{l=-\infty}^0 2^{l(M-s^+)} g_{k+l}^1 = \sum_{u=0}^k 2^{(u-k)(M-s^+)} g_u^1 \leq \sum_{u=0}^{\infty} 2^{-|u-k|(M-s^+)} g_u^1,$$

where $g_u^1 := 2^{us(x)} P_{2^u, a} f_{(u)}(x)$. The application of Lemma 9 and Theorem 14 with $a > 0$ large enough gives

$$\| (f^{(k),I})_{k=0}^{\infty} | \ell_{q(\cdot)}(L_{p(\cdot)}) \| \lesssim \| (g_u)_{u=0}^{\infty} | \ell_{q(\cdot)}(L_{p(\cdot)}) \| \lesssim \| f | B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \|.$$

To estimate $f^{(k),II}$, we proceed as in the Step 4 of Part II. If $p^- > 1$, we choose again $\lambda = 1$, otherwise we take $0 < \lambda < p^-$ and

$$a > \frac{n + c_{\log}(1/q)}{p^-} + c_{\log}(s)$$

such that $a(1-\lambda) < s^-$. This is possible due to (59).

We use (69) with $m > \max(n + c_{\log}(1/q), c_{\log}(s))$ to get

$$\begin{aligned} f^{(k),II} &\lesssim \sum_{l=1}^{\infty} 2^{la(1-\lambda)-ls^-} \left(2^{(k+l)s(x)} P_{2^{k+l}, a} f_{(k+l)}(x) \right)^{1-\lambda} \cdot \\ &\quad \cdot (\eta_{k,m} * |2^{(k+l)s(\cdot)} f_{(k+l)}(\cdot)|^{\lambda})(x) \\ &= \sum_{l=1}^{\infty} 2^{la(1-\lambda)-ls^-} (g_{k+l}^1(x))^{1-\lambda} \cdot (\eta_{k,m} * (g_{k+l}^2)^{\lambda})(x), \end{aligned} \quad (79)$$

where $g_{k+l}^2(x) := |2^{(k+l)s(x)} f_{(k+l)}(x)|$. We take the $\ell_{q(\cdot)}(L_{p(\cdot)})$ (quasi-)norm of the last expression - and assume that it is equivalent to some r -norm. This gives for $\delta := s^- - a(1-\lambda) > 0$ the following estimate

$$\begin{aligned} \| f^{(k),II} | \ell_{q(\cdot)}(L_{p(\cdot)}) \| &\lesssim \sum_{l=1}^{\infty} 2^{-l\delta r} \| (g_{k+l}^1(x))^{1-\lambda} \cdot (\eta_{k,m} * (g_{k+l}^2)^{\lambda})(x) | \ell_{q(\cdot)}(L_{p(\cdot)}) \| & (80) \\ &\lesssim \sum_{l=1}^{\infty} 2^{-l\delta r} \| g_{k+l}^1 | \ell_{q(\cdot)}(L_{p(\cdot)}) \|^{(1-\lambda)r} \cdot \| [\eta_{k,m} * (g_{k+l}^2)^{\lambda}]^{1/\lambda} | \ell_{q(\cdot)}(L_{p(\cdot)}) \| & \\ &\lesssim \sum_{l=1}^{\infty} 2^{-l\delta r} \| g_{k+l}^1 | \ell_{q(\cdot)}(L_{p(\cdot)}) \|^{(1-\lambda)r} \cdot \| \eta_{k,m} * (g_{k+l}^2)^{\lambda} | \ell_{q(\cdot)/\lambda}(L_{p(\cdot)/\lambda}) \| & \\ &\lesssim \sum_{l=1}^{\infty} 2^{-l\delta r} \| g_{k+l}^1 | \ell_{q(\cdot)}(L_{p(\cdot)}) \|^{(1-\lambda)r} \cdot \| (g_{k+l}^2)^{\lambda} | \ell_{q(\cdot)/\lambda}(L_{p(\cdot)/\lambda}) \| & \\ &\lesssim \sum_{l=1}^{\infty} 2^{-l\delta r} \| g_{k+l}^1 | \ell_{q(\cdot)}(L_{p(\cdot)}) \|^{(1-\lambda)r} \cdot \| g_{k+l}^2 | \ell_{q(\cdot)}(L_{p(\cdot)}) \| & \\ &\lesssim \| f | B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) \|^r. \end{aligned}$$

We have used Lemma 10 and Lemma 25.

Part V. The right hand side inequality of (63) follows also along the same line as in Part III. We just combine (78) with the choice $r > s^+ + n$ and apply Lemma 9. \square

4.4 Proof of Theorem 15

This section is devoted to the proof of Theorem 15. We start with the case of constant q . In that case, the usual Hardy-Littlewood maximal operator

$$Mf(x) = \sup_{r>0} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy$$

is bounded on $\ell_q(L_{p(\cdot)})$ and $L_{p(\cdot)}(\ell_q)$. Indeed, the following lemma is a consequence of [12] and [16, Theorem 4.3.8].

Lemma 17. (i) Let $p \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $1 < p^- \leq p^+ < \infty$ and $1 < q < \infty$. Then

$$\|(Mf_j)_{j=-\infty}^{\infty} | L_{p(\cdot)}(\ell_q)\| \lesssim \|(f_j)_{j=-\infty}^{\infty} | L_{p(\cdot)}(\ell_q)\|$$

for all $(f_j)_{j=-\infty}^{\infty} \in L_{p(\cdot)}(\ell_q)$.

(ii) Let $p \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $p^- > 1$ and $0 < q \leq \infty$. Then

$$\|(Mf_j)_{j=-\infty}^{\infty} | \ell_q(L_{p(\cdot)})\| \lesssim \|(f_j)_{j=-\infty}^{\infty} | \ell_q(L_{p(\cdot)})\|$$

for all $(f_j)_{j=-\infty}^{\infty} \in \ell_q(L_{p(\cdot)})$.

Proof of Theorem 15. With the help of Lemma 17, we prove Theorem 15 for q constant. In view of Lemma 16, it is enough to prove

$$\|f | F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**} \lesssim \|f | F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\| \quad (81)$$

and a corresponding analogue for the B -spaces.

Part I. In this part we point out the necessary modifications in the proof of Lemma 16 to obtain a characterization by ball means of differences for $B_{p(\cdot),q}^{s(\cdot)}(\mathbb{R}^n)$ and $F_{p(\cdot),q}^{s(\cdot)}(\mathbb{R}^n)$. The proof follows the scheme of Part II of the proof of Lemma 16.

We start with $\sum_{k=-\infty}^{\infty}$ instead of $\sum_{k=0}^{\infty}$. With this modification the Steps 1-3 go through without any other changes and we obtain (67) again (just recall that $f_{(k)} = 0$ if $k < 0$).

Due to the boundedness of the maximal operator there is no need for the use of r -trick and convolution with $\eta_{\nu,m}$. The analogue of (68), (69) and (70) now reads as follows:

$$\begin{aligned} & 2^{ks(x)} \int_B |\Delta_{2^{-k}h}^M f_{(k+l)}(x)| dh \\ & \lesssim \left(2^{ks(x)} 2^{la} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{1-\lambda} \sum_{j=0}^M c_{j,M} \int_B 2^{ks(x+j2^{-k}h)\lambda} |f_{(k+l)}(x+j2^{-k}h)|^\lambda dh \\ & \leq 2^{la(1-\lambda)-ls^-} \left(2^{(k+l)s(x)} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{1-\lambda} \\ & \quad \cdot \sum_{j=0}^M c_{j,M} \int_B 2^{(k+l)s(x+j2^{-k}h)\lambda} |f_{(k+l)}(x+j2^{-k}h)|^\lambda dh \\ & \lesssim 2^{la(1-\lambda)-ls^-} \left(2^{(k+l)s(x)} P_{2^{k+l},a} f_{(k+l)}(x) \right)^{1-\lambda} \sum_{j=0}^M c_{j,M} M(|2^{(k+l)s(\cdot)} f_{(k+l)}(\cdot)|^\lambda)(x), \end{aligned}$$

where we used Hölder's regularity of $s(\cdot)$, see (8). As a consequence, we obtain

$$II^{1/q(x)} \lesssim F(x)^{1-\lambda} \cdot \sum_{l=1}^{\infty} 2^{-l\delta} \left(\sum_{k=-l}^{\infty} (MB_{k+l}^\lambda)^{q(x)/\lambda}(x) \right)^{\lambda/q(x)}.$$

instead of (71). The rest then follows in the same manner with the help of Lemma 17 and the proof of (81) is finished.

The proof of

$$\|f|B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|^{**} \lesssim \|f|B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\| \quad (82)$$

goes along the same lines. Especially, we get

$$f^{(k),II} \lesssim \sum_{l=1}^{\infty} 2^{la(1-\lambda)-ls^-} (g_{k+l}^1(x))^{1-\lambda} \cdot (M(g_{k+l}^2)^\lambda)(x)$$

instead of (79). The rest follows again by Lemma 17.

Part II. Finally, we present how the characterization for q constant can help us to improve on the case of variable exponent $q(\cdot)$.

In view of Lemma 16, it is enough to show that

$$\left\| \left(\sum_{k=-\infty}^0 2^{ks(x)q(x)} (d_{2^{-k}}^M f(x))^{q(x)} \right)^{1/q(x)} \Big|_{L_{p(\cdot)}(\mathbb{R}^n)} \right\| \lesssim \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|.$$

But this is a consequence of

$$\begin{aligned} & \left\| \left(\sum_{k=-\infty}^0 2^{ks(x)q(x)} (d_{2^{-k}}^M f(x))^{q(x)} \right)^{1/q(x)} \Big|_{L_{p(\cdot)}(\mathbb{R}^n)} \right\| \\ & \lesssim \left\| \left(\sum_{k=-\infty}^0 2^{k(s(x)-\varepsilon)q^-} (d_{2^{-k}}^M f(x))^{q^-} \right)^{1/q^-} \Big|_{L_{p(\cdot)}(\mathbb{R}^n)} \right\| \\ & \lesssim \|f|F_{p(\cdot),q^-}^{s(\cdot)-\varepsilon}(\mathbb{R}^n)\| \lesssim \|f|F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)\|, \end{aligned}$$

where $\varepsilon > 0$ is small enough and we used the differences characterization for fixed q and a trivial embedding theorem.

The same arguments apply for the Besov spaces and the proof is finished. \square

Remark 8. The somewhat complicated proof for Theorem 15 would be made more direct and simpler if we could use versions of Lemmas 10 and 21 in (72) and (80) where the $\ell_{q(\cdot)}$ summation runs over $\nu \in \mathbb{Z}$.

For Triebel-Lizorkin spaces there seems to exist such an extension [14], but for Besov spaces the proof of Lemma 10 in [1] seems to be to customized to the situation $\nu \in \mathbb{N}_0$.

4.5 Ball means of differences for 2-microlocal spaces

As already remarked in Section 2.2 all the proofs for spaces of variable smoothness do also serve for 2-microlocal spaces. One just has to use the definition of admissible weight sequences and the property (8), see Remark 2.

First of all we give the notation for the (quasi-)norms. For simplicity we just use the discrete versions, although it is also possible to give continuous versions of 2-microlocal weights, see [55, Definition 4.1]. In analogy to the spaces of variable smoothness we introduce the following norms

$$\left\| |f| B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \right\|^{**} = \left\| |f| L_{p(\cdot)}(\mathbb{R}^n) \right\| + \left\| (w_k(x) d_{2^{-k}}^M f(x))_{k=-\infty}^{\infty} \Big|_{\ell_{q(\cdot)}(L_{p(\cdot)})} \right\|$$

and

$$\left\| |f| F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \right\|^{**} = \left\| |f| L_{p(\cdot)}(\mathbb{R}^n) \right\| + \left\| (w_k(x) d_{2^{-k}}^M f(x))_{k=-\infty}^{\infty} \Big|_{L_{p(\cdot)}(\ell_{q(\cdot)})} \right\|.$$

Finally, the preceding calculations show that the following theorem is true.

Theorem 18. (i) Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $p^+, q^+ < \infty$ and $\mathbf{w} \in \mathcal{W}_{\alpha_1, \alpha_2}^{\alpha}$. Let $M > \alpha_2$ and

$$\alpha_1 > \sigma_{p^-, q^-} \cdot \left[1 + \frac{\alpha}{n} \cdot \min(p^-, q^-) \right]. \quad (83)$$

Then

$$F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) = \{f \in L_{p(\cdot)}(\mathbb{R}^n) : \| |f| F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \|^{**} < \infty\}$$

and $\| \cdot \|_{F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)}$ and $\| \cdot \|_{|F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)|}^{**}$ are equivalent on $F_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$.

(ii) Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and $\mathbf{w} \in \mathcal{W}_{\alpha_1, \alpha_2}^{\alpha}$. Let $M > \alpha_2$ and

$$\alpha_1 > \sigma_{p^-} \cdot \left[1 + \frac{c_{\log}(1/q)}{n} + \frac{\alpha}{n} \cdot p^- \right]. \quad (84)$$

Then

$$B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) = \{f \in L_{p(\cdot)}(\mathbb{R}^n) : \| |f| B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \|^{**} < \infty\}$$

and $\| \cdot \|_{B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)}$ and $\| \cdot \|_{|B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)|}^{**}$ are equivalent on $B_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$.

Remark 9. Again, if $\min(p^-, q^-) \geq 1$ in the F-case, or $p^- \geq 1$ in the B-case, then the conditions (83) and (84) simplify to $\alpha_1 > 0$. In the case of constant exponents p, q we obtain similar results to [8] and [34].

4.6 Lemmas

The following lemma is a variant of Lemma 6.1 from [17].

Lemma 19. Let $s \in C_{loc}^{\log}(\mathbb{R}^n)$ and let $R \geq c_{\log}(s)$, where $c_{\log}(s)$ is the constant from (4) for $s(\cdot)$. Then

$$2^{\nu s(x)} \eta_{\nu, m+R}(x-y) \leq c 2^{\nu s(y)} \eta_{\nu, m}(x-y)$$

holds for all $x, y \in \mathbb{R}^n$ and $m \in \mathbb{N}_0$.

Lemma 20. *Let $r > 0$, $\nu \geq 0$ and $m > n$. Then there exists $c > 0$, which depends only on m, n and r , such that for all $g \in S'(\mathbb{R}^n)$ with $\text{supp } \hat{g} \subset \{\xi \in \mathbb{R}^n : |\xi| \leq 2^{\nu+1}\}$, we have*

$$|g(x)| \leq c(\eta_{\nu, m} * |g|^r(x))^{1/r}, \quad x \in \mathbb{R}^n.$$

The following lemma is the counterpart to Lemma 10 for Triebel-Lizorkin spaces.

Lemma 21 ([17], Theorem 3.2). *Let $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $1 < p^- \leq p^+ < \infty$ and $1 < q^- \leq q^+ < \infty$. Then the inequality*

$$\|(\eta_{\nu, m} * f)_{\nu=0}^{\infty}|_{L_{p(\cdot)}(\ell_{q(\cdot)})}\| \leq c \| (f_{\nu})_{\nu=0}^{\infty}|_{L_{p(\cdot)}(\ell_{q(\cdot)})}\|$$

holds for every sequence $(f_{\nu})_{\nu \in \mathbb{N}_0}$ of $L_1^{\text{loc}}(\mathbb{R}^n)$ functions and $m > n$.

The following lemma is well known (cf. [51]). We sketch its proof for the sake of completeness.

Lemma 22. *Let $a, b > 0$, $M \in \mathbb{N}$ and $h \in \mathbb{R}^n$. Let $f \in S'(\mathbb{R}^n)$ with $\text{supp } \hat{f} \subset \{\xi \in \mathbb{R}^n : |\xi| \leq b\}$. Then there is a constant $C > 0$ independent of f, b and h , such that*

$$|\Delta_h^M f(x)| \leq C \max(1, |bh|^a) \cdot \min(1, |bh|^M) P_{b,a} f(x)$$

holds for every $x \in \mathbb{R}^n$.

Proof. The estimate

$$\begin{aligned} |f(x + jh)| &= \frac{|f(x + jh)|}{1 + |jbh|^a} \cdot (1 + |jbh|^a) \leq (1 + |Mbh|^a) \sup_{z \in \mathbb{R}^n} \frac{f(x - z)}{1 + |bz|^a} \\ &\lesssim \max(1, |bh|^a) P_{b,a} f(x), \quad j = 0, \dots, M, \end{aligned}$$

holds for all the admissible parameters even without the assumption on \hat{f} .

Hence we need to prove only

$$|\Delta_h^M f(x)| \leq C \max(1, |bh|^a) \cdot |bh|^M \cdot P_{b,a} f(x). \quad (85)$$

Using the Taylor formula for the (analytic) function f , we obtain by direct calculation

$$\begin{aligned} |\Delta_h^M f(x)| &\leq c|h|^M \sup_{|\alpha|=M} \sup_{|y| \leq M|h} \frac{|(D^\alpha f)(x - y)|}{1 + |by|^a} \cdot (1 + |by|^a) \\ &\leq c'|h|^M \max(1, |bh|^a) \cdot \sup_{|\alpha|=M} \sup_{|y| \leq M|h} \frac{|(D^\alpha f)(x - y)|}{1 + |by|^a}. \end{aligned}$$

If $\text{supp } \hat{g} \subset \{\xi \in \mathbb{R}^n : |\xi| \leq 1\}$, then this may be combined with the Nikol'skij inequality, cf. [51, Section 1.3.1], in the form

$$\sup_{|\alpha|=M} \sup_{z \in \mathbb{R}^n} \frac{|(D^\alpha g)(x - z)|}{1 + |z|^a} \lesssim \sup_{z \in \mathbb{R}^n} \frac{|g(x - z)|}{1 + |z|^a}$$

to obtain

$$|\Delta_h^M g(x)| \leq c''|h|^M \max(1, |h|^a) \cdot \sup_{z \in \mathbb{R}^n} \frac{|g(x - z)|}{1 + |z|^a}. \quad (86)$$

If $\text{supp } \hat{f} \subset \{\xi \in \mathbb{R}^n : |\xi| \leq b\}$, we define $g(x) = f(x/b)$, apply (86) together with $\Delta_h^M f(x) = \Delta_{bh}^M g(x)$ and obtain

$$|\Delta_h^M f(x)| \lesssim |bh|^M \max(1, |bh|^a) \sup_{z \in \mathbb{R}^n} \frac{|g(bx - z)|}{1 + |z|^a}.$$

From this (85) follows and the proof is then complete. \square

Following lemma resembles Lemma A.3 of [17].

Lemma 23. *Let $k \in \mathbb{Z}$, $l \in \mathbb{N}_0$ and $m > n$. Then*

$$\eta_{k+l,m} * [2^{kn} \chi_{2^{-k}B}] \lesssim \eta_{k,m}.$$

Proof. Using dilations, we may suppose that $k = 0$. If $|x| \leq 2$, then

$$\int_{\{y:|x-y|\leq 1\}} 2^{nl}(1+2^l|y|)^{-m} dy \leq \int_{y \in \mathbb{R}^n} 2^{nl}(1+2^l|y|)^{-m} dy \lesssim (1+|x|)^{-m}.$$

If $|x| > 2$ and $|x - y| \leq 1$, we obtain $1 + 2^l|y| \gtrsim 1 + 2^l|x|$ and $2^{nl}(1 + 2^l|x|)^{-m} \lesssim (1 + |x|)^{-m}$. This immediately implies that

$$\int_{\{y:|x-y|\leq 1\}} 2^{nl}(1+2^l|y|)^{-m} dy \lesssim \int_{\{y:|x-y|\leq 1\}} (1+|x|)^{-m} dy \lesssim (1+|x|)^{-m}.$$

\square

Remark 10. Another way, how to prove Lemma 23 is to use the inequality $\chi_B(x) \leq 2^m \eta_{0,m}(x)$ and apply Lemma A.3 of [17].

The following Lemma is quite simple and we leave out its proof.

Lemma 24. *Let $0 < q < \infty$, $\delta > 0$ and let $(a_l)_{l \in \mathbb{N}}$ be a sequence of non-negative real numbers. Then*

$$\left(\sum_{l=1}^{\infty} 2^{-l\delta q} a_l \right)^{1/q} \lesssim \sum_{l=1}^{\infty} 2^{-l\delta/2} a_l^{1/q},$$

where the constant involved depends only on δ and q .

Finally, we shall need a certain version of Hölder's inequality for $\ell_{q(\cdot)}(L_{p(\cdot)})$ spaces.

Lemma 25. *Let $p, q \in \mathcal{P}(\mathbb{R}^n)$ and let $0 < \lambda < 1$. Then*

$$\|f_k \cdot g_k\|_{\ell_{q(\cdot)}(L_{p(\cdot)})} \leq 2^{1/q^-} \|f_k^{1/(1-\lambda)}\|_{\ell_{q(\cdot)}(L_{p(\cdot)})}^{1-\lambda} \cdot \|g_k^{1/\lambda}\|_{\ell_{q(\cdot)}(L_{p(\cdot)})}^{\lambda} \quad (87)$$

holds for all sequences of non-negative functions $(f_k)_{k \in \mathbb{N}_0}$ and $(g_k)_{k \in \mathbb{N}_0}$.

Proof. Due to the homogeneity, we may assume that

$$\left\| f_k^{1/(1-\lambda)} \right|_{\ell_{q(\cdot)}(L_{p(\cdot)})} = \left\| g_k^{1/\lambda} \right|_{\ell_{q(\cdot)}(L_{p(\cdot)})} = 1.$$

Then to every $\varepsilon > 0$, there exist two sequences of positive real numbers $(\lambda_k)_{k \in \mathbb{N}_0}$ and $(\mu_k)_{k \in \mathbb{N}_0}$, such that

$$\sum_{k=0}^{\infty} \lambda_k < 1 + \varepsilon, \quad \sum_{k=0}^{\infty} \mu_k < 1 + \varepsilon$$

and

$$\varrho_{p(\cdot)} \left(\frac{f_k^{1/(1-\lambda)}}{\lambda_k^{1/q(\cdot)}} \right) \leq 1, \quad \varrho_{p(\cdot)} \left(\frac{g_k^{1/\lambda}}{\mu_k^{1/q(\cdot)}} \right) \leq 1.$$

We put

$$c := 2^{1/q^-} \quad \text{and} \quad \gamma_k := \frac{\lambda_k + \mu_k}{2} \geq \frac{\lambda_k + \mu_k}{c^{q(x)}}$$

and use the Young inequality in the form

$$[f_k(x)g_k(x)]^{p(x)} \leq (1-\lambda)f_k(x)^{p(x)/(1-\lambda)} + \lambda g_k(x)^{p(x)/\lambda}$$

to obtain

$$\begin{aligned} \int_{\mathbb{R}^n} \left(\frac{f_k(x)g_k(x)}{c\gamma_k^{1/q(\cdot)}} \right)^{p(x)} dx &\leq (1-\lambda) \int_{\mathbb{R}^n} \frac{f_k(x)^{p(x)/(1-\lambda)}}{c^{p(x)}\gamma_k^{p(x)/q(x)}} dx + \lambda \int_{\mathbb{R}^n} \frac{g_k(x)^{p(x)/\lambda}}{c^{p(x)}\gamma_k^{p(x)/q(x)}} dx \\ &\leq (1-\lambda) \int_{\mathbb{R}^n} \frac{f_k(x)^{p(x)/(1-\lambda)}}{\lambda_k^{p(x)/q(x)}} dx + \lambda \int_{\mathbb{R}^n} \frac{g_k(x)^{p(x)/\lambda}}{\mu_k^{p(x)/q(x)}} dx \leq 1. \end{aligned}$$

Furthermore, the estimate

$$\sum_{k=0}^{\infty} \gamma_k < \frac{2(1+\varepsilon)}{2} = 1 + \varepsilon$$

finishes the proof of (87) with the constant $c = 2^{1/q^-}$. \square

Acknowledgement: The first author acknowledges the financial support provided by the DFG project HA 2794/5-1 "Wavelets and function spaces on domains". Furthermore, the first author thanks the RICAM for his hospitality and support while a short term visit in Linz.

The second author acknowledges the financial support provided by the FWF project Y 432-N15 START-Preis "Sparse Approximation and Optimization in High Dimensions".

References

- [1] A. Almeida, P. Hästö: *Besov spaces with variable smoothness and integrability*, J. Funct. Anal. **258** (2010), no. 5, 1628–1655.
- [2] A. Almeida, S. Samko: *Characterization of Riesz and Bessel potentials on variable Lebesgue spaces*, J. Function Spaces Appl. **4** (2006), no. 2, 113–144.
- [3] A. Almeida, S. Samko: *Pointwise inequalities in variable Sobolev spaces and applications*, Z. Anal. Anwend. **26** (2007), no. 2, 179–193.
- [4] A. Almeida, S. Samko: *Embeddings of variable Hajlasz-Sobolev spaces into Hölder spaces of variable order*, J. Math. Anal. Appl. **353** (2009), no. 2, 489–496.
- [5] P. Andersson: *Two-microlocal spaces, local norms and weighted spaces*, Paper 2 in PhD Thesis (1997), 35–58.
- [6] T. Aoki: *Locally bounded linear topological spaces*, Proc. Imp. Acad. Tokyo **18** (1942), 588–594.
- [7] B. Beauzamy: *Espaces de Sobolev et de Besov d'ordre variable définis sur L^p* , C.R. Acad. Sci. Paris (Ser. A) **274** (1972), 1935–1938.
- [8] O. V. Besov: *Equivalent normings of spaces of functions of variable smoothness*, Proc. Steklov Inst. Math. **243** (2003), no. 4, 80–88.
- [9] J.-M. Bony: *Second microlocalization and propagation of singularities for semi-linear hyperbolic equations*, Taniguchi Symp. HERT. Katata (1984), 11–49.
- [10] F. Cobos, D. L. Fernandez: *Hardy-Sobolev spaces and Besov spaces with a function parameter* Proc. Lund Conf. 1986, Lect. Notes Math. **1302** (1986), Berlin: Springer, 158–170.
- [11] D. Cruz-Uribe, A. Fiorenza, C. J. Neugebauer: *The maximal function on variable L^p spaces*, Ann. Acad. Sci. Fenn. Math. **28** (2003), 223–238.
- [12] D. Cruz-Uribe, A. Fiorenza, J. M. Martell, C. Pérez: *The boundedness of classical operators in variable L^p -spaces*, Ann. Acad. Sci. Fenn. Math. **31** (2006), 239–264.
- [13] L. Diening: *Maximal function on generalized Lebesgue spaces $L^{p(\cdot)}$* , Math. Inequal. Appl. **7** (2004), no. 2, 245–254.
- [14] L. Diening: *private communication*.
- [15] L. Diening, P. Harjulehto, P. Hästö, Y. Mizuta, T. Shimomura: *Maximal functions in variable exponent spaces: limiting cases of the exponent*, Ann. Acad. Sci. Fenn. Math. **34** (2009), (2), 503–522.

- [16] L. Diening, P. Harjulehto, P. Hästö, M. Růžička: *Lebesgue and Sobolev Spaces with Variable Exponents*, Springer, Lecture Notes in Mathematics **2017**, Springer (2011).
- [17] L. Diening, P. Hästö, S. Roudenko: *Function spaces of variable smoothness and integrability*, J. Funct. Anal. **256** (2009), (6), 1731–1768.
- [18] W. Farkas, H.-G. Leopold: *Characterisations of function spaces of generalised smoothness*, Ann. Mat. Pura Appl. **185** (2006), no. 1, 1–62.
- [19] M. L. Goldman: *A description of the traces of some function spaces*, Trudy Mat. Inst. Steklov **150** (1979), 99–127; English transl.: Proc. Steklov Inst. Math. **150** (1981), no. 4.
- [20] M. L. Goldman: *A method of coverings for describing general spaces of Besov type*, Trudy Mat. Inst. Steklov **156** (1980), 47–81; English transl.: Proc. Steklov Inst. Math. **156** (1983), no. 2.
- [21] M. L. Goldman: *Imbedding theorems for anisotropic Nikolskij-Besov spaces with moduli of continuity of general type*, Trudy Mat. Inst. Steklov **170** (1984), 86–104; English transl.: Proc. Steklov Inst. Math. **170** (1987), no. 1.
- [22] P. Gurka, P. Harjulehto, A. Nekvinda: *Bessel potential spaces with variable exponent*, Math. Inequal. Appl. **10** (2007), no. 3, 661–676.
- [23] S. Jaffard: *Pointwise smoothness, two-microlocalisation and wavelet coefficients*, Publications Mathematiques **35**, (1991), 155–168.
- [24] S. Jaffard, Y. Meyer: *Wavelet methods for pointwise regularity and local oscillations of functions*, Memoirs of the AMS, vol. **123**, (1996).
- [25] G. A. Kalyabin: *Characterization of spaces of generalized Liouville differentiation*, Mat. Sb. Nov. Ser. **104** (1977), 42–48.
- [26] G. A. Kalyabin: *Description of functions in classes of Besov-Lizorkin-Triebel type* Trudy Mat. Inst. Steklov **156** (1980), 82–109; English transl.: Proc. Steklov Institut Math. **156** (1983), no. 2.
- [27] G. A. Kalyabin, P. I. Lizorkin: *Spaces of functions of generalized smoothness*, Math. Nachr. **133** (1987), 7–32.
- [28] G. A. Kalyabin, *Characterization of spaces of Besov-Lizorkin and Triebel type by means of generalized differences*, Trudy Mat. Inst. Steklov **181**, (1988), 95–116; English transl.: Proc. Steklov Inst. Math. **181** (1989), no. 4.
- [29] H. Kempka: *2-microlocal Besov and Triebel-Lizorkin spaces of variable integrability*, Rev. Mat. Complut. **22** (2009), no. 1, 227–251.
- [30] H. Kempka: *Atomic, molecular and wavelet decomposition of 2-microlocal Besov and Triebel-Lizorkin spaces with variable integrability*, Funct. Approx. **43** (2010), (2), 171–208.

- [31] H. Kempka, J. Vybíral: *A note on the spaces of variable integrability and summability of Almeida and Hästö*, submitted.
- [32] O. Kováčik, J. Rákosník: *On spaces $L^{p(x)}$ and $W^{1,p(x)}$* , Czechoslovak Math. J. **41** (1991), no. 4, 592–618.
- [33] H.-G. Leopold: *On function spaces of variable order of differentiation*, Forum Math. **3** (1991), 1–21.
- [34] J. Lévy Véhel, S. Seuret: *A time domain characterization of 2-microlocal Spaces*, J. Fourier Analysis and Appl. **9** (2003), (5), 473–495.
- [35] J. Lévy Véhel, S. Seuret: *The 2-microlocal formalism*, Fractal Geometry and Applications: A Jubilee of Benoit Mandelbrot, Proceedings of Symposia in Pure Mathematics, PSPUM, vol. **72** (2004), part2, 153–215.
- [36] C. Merucci: *Applications of interpolation with a function parameter to Lorentz, Sobolev and Besov spaces*. Proc. Lund Conf. 1983, Lect. Notes Math. **1070** (1983), Berlin: Springer, 183–201.
- [37] Y. Meyer: *Wavelets, vibrations and scalings*, CRM Monograph Series **9**, AMS (1998).
- [38] S. Moritoh, T. Yamada: *Two-microlocal Besov spaces and wavelets*, Rev. Mat. Iberoamericana **20**, (2004), 277–283.
- [39] S. Moura: *Function spaces of generalised smoothness*, Diss. Math. **398** (2001), 1–87.
- [40] A. Nekvinda: *Hardy-Littlewood maximal operator on $L^{p(x)}(\mathbb{R}^n)$* , Math. Inequal. Appl. **7** (2004), no. 2, 255–266.
- [41] W. Orlicz: *Über konjugierte Exponentenfolgen*, Studia Math. **3**, (1931), 200–212.
- [42] J. Peetre: *On spaces of Triebel-Lizorkin type*, Ark. Math. **13**, (1975), 123–130.
- [43] S. Rolewicz: *On a certain class of linear metric spaces*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astrono. Phys. **5**, (1957), 471–473.
- [44] B. Ross, S. Samko: *Fractional integration operator of variable order in the spaces H^λ* , Int. J. Math. Sci. **18** (1995), no. 4, 777–788.
- [45] M. Ružička: *Electrorheological fluids: modeling and mathematical theory*, Lecture Notes in Mathematics **1748**, Springer, Berlin, (2000).
- [46] V. S. Rychkov: *On a theorem of Bui, Paluszynski and Taibleson*, Steklov Institute of Mathematics **227**, (1999), 280–292.
- [47] B. Scharf: *Atomare Charakterisierungen vektorwertiger Funktionenräume*, Diploma Thesis, Jena (2009).

- [48] C. Schneider: *On dilation operators in Besov spaces*, Rev. Mat. Complut. **22** (2009), no. 1, 111–128.
- [49] C. Schneider, J. Vybíral: *On dilation operators in Triebel-Lizorkin spaces*, Funct. Approx. Comment. Math. **41** (2009), part 2, 139–162.
- [50] J. Schneider: *Function spaces of varying smoothness I*, Math. Nachr. **280** (2007), no. 16, 1801–1826.
- [51] H. Triebel: *Theory of function spaces*, Basel, Birkhäuser, (1983).
- [52] H. Triebel: *Theory of function spaces II*, Basel, Birkhäuser, (1992).
- [53] T. Ullrich, *Function spaces with dominating mixed smoothness, characterization by differences*, Technical report, Jenaer Schriften zur Math. und Inform., Math/Inf/05/06, (2006).
- [54] T. Ullrich: *Continuous characterizations of Besov-Lizorkin-Triebel spaces and new interpretations as coorbit spaces*, to appear in J. Funct. Spaces Appl.
- [55] T. Ullrich, H. Rauhut: *Generalized coorbit space theory and inhomogeneous function spaces of Besov-Lizorkin-Triebel type*, to appear in J. Funct. Anal.
- [56] A. Unterberger, J. Bokobza: *Les opérateurs pseudodifférentiels d'ordre variable*, C.R. Acad. Sci. Paris **261** (1965), 2271–2273.
- [57] A. Unterberger: *Sobolev Spaces of Variable Order and Problems of Convexity for Partial Differential Operators with Constant Coefficients*, Astérisque **2 et 3** SOC. Math. France (1973), 325–341.
- [58] M. I. Višik, G. I. Eskin: *Convolution equations of variable order* (russ.), Trudy Moskov Mat. Obsc. **16** (1967), 26–49.
- [59] J. Vybíral: *Sobolev and Jawerth embeddings for spaces with variable smoothness and integrability*, Ann. Acad. Sci. Fenn. Math. **34** (2009), no. 2, 529–544.
- [60] H. Xu: *Généralisation de la théorie des chirps à divers cadres fonctionnels et application à leur analyse par ondelettes*, Ph. D. thesis, Université Paris IX Dauphine (1996).
- [61] J.-S. Xu: *Variable Besov and Triebel-Lizorkin spaces*, Ann. Acad. Sci. Fenn. Math. **33** (2008), no. 2, 511–522.
- [62] J.-S. Xu: *An atomic decomposition of variable Besov and Triebel-Lizorkin spaces*, Armenian J. Math. **2** (2009), no. 1, 1–12.
- [63] W. Yuan, W. Sickel, D. Yang: *Morrey and Campanato meet Besov, Lizorkin and Triebel*, Lecture Notes in Mathematics **2005**, Springer (2010).