

Minimal Set Theorems

Andreas Hamel*

Andreas Löhne**

Abstract

We present two existence principles for minimal points of subsets of the product space $X \times 2^Y$, where X stands for a separated uniform space and Y a topological vector space. The two principles are distinct with respect to the involved ordering structure in 2^Y .

We derive from them new variants of Ekeland's principle for set-valued maps as well as a minimal point theorem in $X \times Y$ and Ekeland's principle for vector-valued functions.

Keywords and phrases: minimal set theorem, set-valued variational principle, minimal point theorem, set-valued optimization, natural criteria

Mathematical Subject Classification. 58E30, 46N10.

*Department of Mathematics and Computer Science Martin-Luther-University Halle-Wittenberg 06099 Halle (Saale), Germany, e-mail: hamel@mathematik.uni-halle.de

**Department of Mathematics and Computer Science Martin-Luther-University Halle-Wittenberg 06099 Halle (Saale), Germany, e-mail: loehne@mathematik.uni-halle.de

1 Introduction

Ekeland's variational principle and its equivalent formulations belong to the cornerstones of Nonlinear Functional Analysis with applications in many fields of analysis, optimization and operations research. During the last years, an increasing interest could be observed for versions involving a set-valued function, compare e.g. [2], [11], [12], [25].

A set-valued mapping from a set X to a set Y is usually understood to be a relation $F \subset X \times Y$ not necessarily satisfying the uniqueness property, i.e. we have not that $(x, y_1), (x, y_2) \in F$ implies $y_1 = y_2$.

In contrast to this, we understand a set-valued mapping to be a function from X to 2^Y , i.e. a relation F on $X \times 2^Y$ satisfying the uniqueness property (i.e. $(x, V_1), (x, V_2) \in F$ implies $V_1 = V_2$). This leads in a natural way to new definitions of concepts like graph, domain and minimal points of set-valued maps as well as new results in set-valued optimization theory.

Investigating an optimization problem with set-valued cost function it is necessary to compare its values. In this paper, we use reflexive and transitive relations on 2^Y to compare two values $F(x_1)$ and $F(x_2)$ of a map $F : X \rightarrow 2^Y$. This is not the common approach to set-valued optimization up to now. Usually, a set-valued optimization problem is reduced to a vector-valued problem by looking for minimal (efficient) points of the set $\bigcup_{x \in X} F(x)$.

We start considering subsets of $X \times 2^Y$ and looking for minimal elements of them with respect to appropriate ordering relations on 2^Y . We present existence results called *Minimal Set Theorems (Generalized Minimal Point Theorems)*, for such elements.

We shall draw several conclusions of the minimal set theorems.

First, we derive new variants of Ekeland's variational principle for vector-valued set-valued maps. Our Ekeland-type theorems are much more general than and cover most of the known results of the field as special cases.

Secondly, we conclude a minimal point theorem in $X \times Y$. Such theorems are well-established and useful tools in vector optimization and related fields, cf. [22], [10], [9], [11] for example.

Finally, the well-known Ekeland-type principle for vector-valued functions $f : X \rightarrow Y$ (e.g. [23], [13], [10]) turns out to be a consequence of our new Ekeland-type principles for set-valued maps as well as of the minimal point theorem.

The paper is organized as follows. In the next section we introduce two ordering relations for elements of 2^Y where Y is a linear topological space as well as related boundedness concepts in the space 2^Y . In Section 3 we present scalarization methods for subsets of 2^Y . These methods are essentially used for the proofs of the minimal set theorems and may be of independent interest. In Section 5 our main results, the minimal set theorems, are presented. Section 6 contains the new set-valued variational principles and several conclusions.

2 Ordering Relations and Boundedness in 2^Y

Let Y be a topological vector space. We denote by 2^Y the set of all subsets of Y including the empty set \emptyset . As usual, the sum of two sets $V_1, V_2 \in 2^Y$ is defined by $V_1 + V_2 := \{v_1 + v_2 : v_1 \in V_1, v_2 \in V_2\}$. We use the convention $\emptyset + V_1 = \emptyset$. The product of $\alpha \in \mathbb{R}$ and $V \in 2^Y \setminus \{\emptyset\}$ is defined by $\alpha V := \{\alpha v : v \in V\}$. Moreover, we define $\alpha \cdot \emptyset = \emptyset$ for $\alpha \neq 0$ and $0 \cdot \emptyset = \{0\}$.

This section is concerned with ordering relations for sets as well as with order boundedness concepts in 2^Y . Such relations has been introduced for the case $Y = \mathbb{R}$ in a paper [27] by R. C. Young from 1931. Kuroiwa gave in [15], [16] from 1998 a more general definition and started developing a new approach to set-valued optimization.

We define two relations \preceq, \preceq_K on 2^Y as follows. These relations are two out of six being natural generalizations of partial orderings on a linear space Y to relations on 2^Y , compare [17] for details.

Definition 2.1 *Let $K \subset Y$ be a convex cone and $V_1, V_2 \in 2^Y$. We define:*

$$\begin{aligned} V_1 \preceq_K V_2 &\iff V_2 \subset V_1 + K; \\ V_1 \preceq V_2 &\iff V_1 \subset V_2 - K. \end{aligned}$$

If there is no risk of confusion the relations are simply denoted by \preceq and \preceq_K .

Note that Luc [20], Chapters 2.5 and 5.1 implicitly used these relations describing the constraints for a set-valued optimization problem. This can be seen observing that $V \preceq_K \{0\}$ iff $V \cap (-K) \neq \emptyset$ and $V \preceq \{0\}$ iff $V \subset -K$.

Both relations can also be expressed by the ordering \leq_K in Y which is defined by $y_1 \leq_K y_2$ iff $y_2 - y_1 \in K$:

$$V_1 \preceq V_2 \iff \forall v_2 \in V_2 \exists v_1 \in V_1 : v_1 \leq_K v_2; \quad (1)$$

$$V_1 \preceq_K V_2 \iff \forall v_1 \in V_1 \exists v_2 \in V_2 : v_1 \leq_K v_2. \quad (2)$$

Furthermore, the following relationships are easy to show:

$$V_1 \preceq V_2 \iff -V_2 \preceq -V_1 \iff V_2 \preceq_{-K} V_1. \quad (3)$$

In the following, we study the properties of \preceq having in mind that by (3) we are able to obtain the same properties for \preceq with K replaced by $-K$ if necessary.

The relations \preceq and \preceq_K are reflexive and transitive. We have no antisymmetry but

$$(V_1 \preceq V_2, V_2 \preceq V_1) \iff V_1 + K = V_2 + K. \quad (4)$$

Introducing the equivalence relation $V_1 \sim V_2$ iff $V_1 \preceq V_2$ and $V_2 \preceq V_1$ we may generate an partial ordering on the set of equivalence classes. Furthermore, it can be shown that for $\alpha_1, \alpha_2 \geq 0$ we have

$$(V_1 \preceq V_2, V_3 \preceq V_4) \implies \alpha_1 V_1 + \alpha_2 V_3 \preceq \alpha_1 V_2 + \alpha_2 V_4. \quad (5)$$

Note that K has not to be pointed (K is pointed iff $K \cap -K = \{0\}$) for proving (1)-(5). Let $V \in 2^Y$ be a subset of Y . We say that $\bar{v} \in V$ is a \leq_K -minimal element of V if $v \in V$, $v \leq_K \bar{v}$ implies $\bar{v} \leq_K v$. The set of all \leq_K -minimal elements of V is denoted by $\text{Min } V$. If K is a convex pointed cone then we have $\bar{v} \in \text{Min } V$ iff $v \in V$, $v \leq_K \bar{v}$ implies $\bar{v} = v$. In this case,

$$(V_1 \preceq V_2, V_2 \preceq V_1) \implies \text{Min}(V_1) = \text{Min}(V_2). \quad (6)$$

A subset $V \subset Y$ is said to be *lower externally stable* iff $V \subset \text{Min } V + K$. This property is called *domination property* by several authors. Compare Luc [20] and the references therein. By direct calculation, one may find for $V_1, V_2 \in 2^Y$ being lower externally stable sets that

$$V_1 \preceq V_2 \iff \text{Min } V_1 \preceq \text{Min } V_2. \quad (7)$$

A similar assertion follows for \succcurlyeq replacing \leq_K -minimal by \leq_K -maximal elements and lower by upper external stability.

The following relationships can easily be verified:

$$\begin{aligned} \forall V \in 2^Y : \quad & V \preceq \emptyset, Y \preceq V, \emptyset \succcurlyeq V, V \succcurlyeq Y; \\ & \emptyset \preceq V \Rightarrow V = \emptyset, \quad V \preceq Y \Rightarrow Y = V + K, \\ & V \succcurlyeq \emptyset \Rightarrow V = \emptyset, \quad Y \succcurlyeq V \Rightarrow Y = V - K. \end{aligned}$$

These relationships motivate the following boundedness concepts for subsets of 2^Y .

Definition 2.2 A subset $\mathcal{V} \subset 2^Y$ is said to be \preceq -bounded below if there exists some topologically bounded subset $\tilde{V} \subset Y$ such that $\tilde{V} \preceq V$ holds for all $V \in \mathcal{V}$. The set \tilde{V} is called a lower \preceq -bound of \mathcal{V} . A subset $\mathcal{V} \subset 2^Y$ is said to be \succcurlyeq -bounded above and \tilde{V} is called an upper \succcurlyeq -bound of \mathcal{V} if $-\mathcal{V} := \{-V : V \in \mathcal{V}\}$ is \preceq -bounded below with the lower \preceq -bound $-\tilde{V}$.

Definition 2.3 A subset $\mathcal{V} \subset 2^Y$ is said to be \succcurlyeq -bounded below if there exists some nonempty subset $\tilde{V} \subset Y$ such that $\tilde{V} \succcurlyeq V$ holds for all $V \in \mathcal{V}$. The set \tilde{V} is called a lower \succcurlyeq -bound of \mathcal{V} . A subset $\mathcal{V} \subset 2^Y$ is said to be \preceq -bounded above and \tilde{V} is called an upper \preceq -bound of \mathcal{V} if $-\mathcal{V}$ is \succcurlyeq -bounded below with the lower \succcurlyeq -bound $-\tilde{V}$.

Using the complementary relations of \preceq and \succcurlyeq , denoted by $\not\preceq$ and $\not\succcurlyeq$ we introduce further boundedness concepts.

Definition 2.4 A subset $\mathcal{V} \subset 2^Y$ is said to be weakly \preceq -bounded below if there exists some topologically bounded subset $\tilde{V} \subset Y$ such that $V \not\preceq \tilde{V}$ holds for all $V \in \mathcal{V}$. The set \tilde{V} is called a weak lower \preceq -bound of \mathcal{V} . A subset $\mathcal{V} \subset 2^Y$ is said to be weakly \succcurlyeq -bounded above and \tilde{V} is called a weak upper \succcurlyeq -bound of \mathcal{V} if $-\mathcal{V}$ is weakly \preceq -bounded below with the weak lower \preceq -bound $-\tilde{V}$.

Definition 2.5 A subset $\mathcal{V} \subset 2^Y$ is said to be weakly \preceq -bounded below if there exists some nonempty subset $\tilde{V} \subset Y$ such that $V \not\preceq \tilde{V}$ holds for all $V \in \mathcal{V}$. The set \tilde{V} is called a weak lower \preceq -bound of \mathcal{V} . A subset $\mathcal{V} \subset 2^Y$ is said to be weakly \preceq -bounded above and \tilde{V} is called a weak upper \preceq -bound of \mathcal{V} if $-\mathcal{V}$ is weakly \preceq -bounded below with the weak lower \preceq -bound $-\tilde{V}$.

Note that we have a kind of duality between nonempty sets and topologically bounded sets in the definition of the above boundedness concepts. This duality can be observed throughout the paper. Further, we have to take care using the symbol \emptyset : It denotes an element of 2^Y but of course a subset $\mathcal{V} \subset 2^Y$ can also be the empty set.

Remark 2.6 Let $\text{cl} K \neq Y$. If $\mathcal{V} \subset 2^Y$ is \preceq -bounded below then \mathcal{V} is weakly \preceq -bounded below. Indeed, let \mathcal{V} be \preceq -bounded below, i.e. there exists some topologically bounded set $\tilde{V} \subset Y$ such that $\tilde{V} \preceq V$ for all $V \in \mathcal{V}$. We show that $\tilde{V} + y$ is a weak lower \preceq -bound of \mathcal{V} , where $y \notin \text{cl} K$. Assuming the contrary, i.e. $V \preceq \tilde{V} + y$ for some $V \in \mathcal{V}$ it follows $\tilde{V} + y \subset V + K \subset \tilde{V} + K$. An induction argument yields $\tilde{V} + ny \subset \tilde{V} + K$ for all $n \in \mathbb{N}$. Dividing by n and letting $n \rightarrow \infty$ implies $y \in \text{cl} K$, a contradiction.

Example 2.7 Let $K = \mathbb{R}_+^2$ and $\mathcal{V} = \{Y \setminus -K\}$. Then \mathcal{V} is weakly \preceq -bounded below but \mathcal{V} is not \preceq -bounded below.

Remark 2.8 Let $\text{cl} K \neq Y$. If $\mathcal{V} \subset 2^Y$ is \preceq -bounded below then \mathcal{V} is weakly \preceq -bounded below. Indeed, let \mathcal{V} be \preceq -bounded below, i.e. there exists some nonempty set $\tilde{V} \subset Y$ such that $\tilde{V} \preceq V$ for all $V \in \mathcal{V}$. We show that $\tilde{v} - y$ is a weak lower \preceq -bound of \mathcal{V} , where $\tilde{v} \in \tilde{V}$ and $y \notin \text{cl} K$. Assuming the contrary, i.e. $V \preceq \tilde{v} - y$ for some $V \in \mathcal{V}$ it follows $\tilde{v} \in \tilde{V} \subset V + K \subset \tilde{v} - y + K$. An induction argument yields $\tilde{v} \in \tilde{v} - ny + K$ for all $n \in \mathbb{N}$. Dividing by n and letting $n \rightarrow \infty$ implies $y \in \text{cl} K$, a contradiction.

Example 2.9 Let $K = \mathbb{R}_+^2$ and $\mathcal{V} = \{\{y\} : y \in Y \setminus -K\}$. Then \mathcal{V} is weakly \preceq -bounded below but \mathcal{V} is not \preceq -bounded below.

3 Scalarization Methods on 2^Y

We present several nonlinear scalarization functionals defined on 2^Y . They are generalizations of the functionals introduced in [6] and extensively studied in [26], [7], [10] and [9]. The most important property of our functionals turns out to be the monotonicity with respect to the relations \preceq , \preceq . Let $V_1, V_2 \in 2^Y$. We call a functional $z : 2^Y \rightarrow \mathbb{R} \cup \{\pm\infty\}$ \preceq -monotone iff $V_1 \preceq V_2$ implies $z(V_1) \leq z(V_2)$. It is called \preceq -monotone iff $V_1 \preceq V_2$ implies $z(V_1) \leq z(V_2)$.

Theorem 3.1 Let Y be a topological vector space, $K \subset Y$ a convex cone and $k^0 \in K \setminus \text{cl} K$. Let $\mathcal{V} \subset 2^Y$ be nonempty and \preceq -bounded, i.e. there is a topological bounded set $V' \subset Y$ and a nonempty set $V'' \subset Y$ such that

$$\forall V \in \mathcal{V} : V' \preceq V \preceq V''.$$

Then, the functional $z^l : 2^Y \rightarrow \mathbb{R} \cup \{\pm\infty\}$ defined by

$$z^l(V) := \inf\{t \in \mathbb{R} : tk^0 + V'' \subset V + \text{cl } K\}$$

has the following properties:

- (i) z^l is bounded on \mathcal{V} ;
- (ii) $V \in \mathcal{V}, \alpha \in \mathbb{R} \Rightarrow z^l(V + \alpha k^0) = z^l(V) + \alpha$;
- (iii) z^l is \preceq -monotone.

Proof. Since V'' is an upper \preceq -bound, we have $V'' \neq \emptyset$ and $V'' \subset V + K$. Hence $V \neq \emptyset$ and $z^l(V) \leq 0$ for all $V \in \mathcal{V}$.

Assume that z^l is not bounded below. Then for each $n \in \mathbb{N}$, we can find some $t_n < -n$ and some $V_n \in \mathcal{V}$ such that $-nk^0 + V'' = (-n - t_n)k^0 + t_n k^0 + V'' \subset K \setminus -\text{cl } K + V_n + \text{cl } K \subset V_n + \text{cl } K$. Since \mathcal{V} is \preceq -bounded below by $V' \subset Y$ we have $-nk^0 + V'' \subset V_n + \text{cl } K \subset V' + \text{cl } K$ for all $n \in \mathbb{N}$. Hence $-nk^0 + v_0 \in V' + \text{cl } K$ for arbitrary $v_0 \in V''$. Dividing by n and letting $n \rightarrow \infty$ we get $k^0 \in -\text{cl } K$ since V' is bounded. This contradicts the assumption $k^0 \in K \setminus -\text{cl } K$.

Assertion (ii) is obvious. To show (iii) let $V_1 \preceq V_2$. Then $V_2 + \text{cl } K \subset V_1 + K + \text{cl } K \subset V_1 + \text{cl } K$. This implies $z^l(V_1) \leq z^l(V_2)$ by definition of z^l . \blacksquare

An analogous result for the relation \preceq is an immediate conclusion.

Corollary 3.2 *Let Y, K, k^0 be as in Theorem 3.1. Let $\mathcal{V} \subset 2^Y$ be nonempty and \preceq -bounded, i.e. there is a nonempty set $W' \subset Y$ and a topologically bounded set $W'' \subset Y$ such that*

$$\forall V \in \mathcal{V} : W' \preceq V \preceq W''.$$

Then, the functional $z^u : 2^Y \rightarrow \mathbb{R} \cup \{\pm\infty\}$ defined by

$$z^u(V) := -\inf\{t \in \mathbb{R} : t(-k^0) + W' \subset V - \text{cl } K\}$$

is bounded on \mathcal{V} , satisfies $z^u(V + \alpha k^0) = z^u(V) + \alpha$ for all $V \in \mathcal{V}$ and is \preceq -monotone.

Proof. Note that \mathcal{V} is \preceq -bounded iff it is \preceq_{-K} -bounded with upper bound W' . Taking into account that $-z^u(V)$ coincides with $z^l(V)$ replacing V'' by W' , K by $-K$ and k^0 by $-k^0$ we may apply Theorem 3.1 to obtain the assertions of the corollary. \blacksquare

Let Y be a topological vector space and $K \subset Y$ a convex cone. We use the following assumption for weakening the boundedness condition.

- (C) There exists a proper closed convex cone $C \subset Y$ with nonempty interior satisfying $K \subset C$.

Remark 3.3 If Y is a locally convex space and $K \subset Y$ a convex cone such that $K \setminus -\text{cl} K \neq \emptyset$ then assumption (C) is satisfied and there exists an element $k^0 \in K \cap \text{int} C$. Indeed, if $k^0 \in K \setminus -\text{cl} K$ then we have $\{-k^0\} \cap \text{cl} K = \emptyset$ and we can apply a classical separation theorem on the convex compact set $\{-k^0\}$ and the closed convex set $\text{cl} K$. We obtain the existence of a linear continuous functional $y^* \in Y^*$ such that $y^*(-k^0) < 0 \leq y^*(k)$ for all $k \in \text{cl} K$. The desired cone C can be defined by $C := \{y \in Y : y^*(y) \geq 0\}$.

Theorem 3.4 Let Y be a topological vector space, $K \subset Y$ a convex cone satisfying assumption (C). Let $k^0 \in K \cap \text{int} C$ and let $\mathcal{V} \subset 2^Y$ be nonempty, \preceq_C -bounded above and weakly \preceq_C -bounded below. Then, the functional $c^l : 2^Y \rightarrow \mathbb{R} \cup \{\pm\infty\}$ defined by

$$c^l(V) := \inf \{t \in \mathbb{R} : V \preceq_C \{tk^0\}\}$$

has the following properties:

- (i) c^l is bounded on \mathcal{V} ;
- (ii) $\forall V \in \mathcal{V} \forall \alpha \in \mathbb{R} : c^l(V + \alpha k^0) = c^l(V) + \alpha$;
- (iii) $V_1 \preceq V_2 \Rightarrow c^l(V_1) \leq c^l(V_2)$;
- (iv) If \mathcal{V} consists of compact sets $V \subset Y$ and $K \setminus \{0\} \subset \text{int} C$ then

$$\left(V_1 \preceq V_2, V_1 \cap V_2 = \emptyset \right) \Rightarrow c^l(V_1) < c^l(V_2).$$

Proof. Let V'' be an upper \preceq_C -bound of \mathcal{V} , i.e. $V'' \neq \emptyset$ and $V'' \subset V + C$ for all $V \in \mathcal{V}$. Let $v'' \in V''$ be given. Because of $k^0 \in \text{int} C$ there exists a neighborhood U of zero such that $U \subset -k^0 + \text{int} C$. Choosing some $\sigma > 0$ such that $-v'' \in \sigma U$ we obtain $v'' \in -\sigma U \subset \sigma(k^0 - \text{int} C) \subset \sigma k^0 - C$. Hence $\sigma k^0 \in V'' + C \subset V + C$ for all $V \in \mathcal{V}$. This means $c^l(V) \leq \sigma$ for all $V \in \mathcal{V}$, i.e. c^l is bounded above.

Assume that c^l is not bounded below. Then, for all $n \in \mathbb{N}$, we can find $t_n < -n$ and $V_n \in \mathcal{V}$ such that $V_n \preceq_C \{t_n k^0\}$. Hence $-nk^0 = (-n - t_n)k^0 + t_n k^0 \in C + V_n + C \subset V_n + C$. Thus

$$\forall n \in \mathbb{N} \exists v_n \in V_n : -nk^0 \in v_n + C. \quad (8)$$

Since \mathcal{V} is supposed to be weakly \preceq -bounded below there exists some topologically bounded set $V' \subset Y$ such that $V' \not\subset V + C$ for all $V \in \mathcal{V}$. Hence for each $n \in \mathbb{N}$ there exists $v'_n \in V'$ such that $v'_n \notin v_n + C$. It follows

$$-k^0 - v'_n/n \stackrel{(8)}{\in} (v_n - v'_n)/n + C \subset (Y \setminus -\text{int} C) + C \subset Y \setminus -\text{int} C.$$

Letting $n \rightarrow \infty$ we get $k^0 \notin \text{int} C$ which contradicts the assumption $k^0 \in K \cap \text{int} C$. Hence c^l is bounded on \mathcal{V} .

Assertions (ii) and (iii) are obvious. Let us prove (iv). By definition of the infimum, for each $n \in \mathbb{N}$ there exists $v_n \in V_2$ such that $(c^l(V_2) + 1/n)k^0 \in v_n + C$. Since V_2 is compact, we can find a subnet of the sequence $\{v_n\}_{n \in \mathbb{N}}$ converging to some $\bar{v} \in V_2$. Hence $c^l(V_2)k^0 \in \bar{v} + C$. Let $V_1 \preccurlyeq V_2$ and $V_1 \cap V_2 = \emptyset$. We have $\bar{v} \in V_2 \subset V_1 + K \setminus \{0\} \subset V_1 + \text{int } C$. Hence there is some $\delta > 0$ such that $\bar{v} - \delta k^0 \in V_1 + \text{int } C$. It follows $c^l(V_2)k^0 \in \bar{v} + C \subset \delta k^0 + V_1 + C$. Applying assertion (ii) we obtain $c^l(V_1) + \delta = c^l(V_1 + \delta k^0) \leq c^l(V_2)$. \blacksquare

The analogous result using the relation \preccurlyeq can not be deduced from Theorem 3.4 by a construction similar to that of Corollary 3.2. Instead, we define a new scalarization functional. Compare the remark following the proof of Theorem 3.5.

Theorem 3.5 *Let Y be a topological vector space, $K \subset Y$ a convex cone satisfying assumption (C). Let $k^0 \in K \cap \text{int } C$ and let $\mathcal{V} \subset 2^Y$ be nonempty, \preccurlyeq_C -bounded above and weakly \preccurlyeq_C -bounded below. Then, the functional $c^u : 2^Y \rightarrow \mathbb{R} \cup \{\pm\infty\}$*

$$c^u(V) := \inf \{t \in \mathbb{R} : V \preccurlyeq_C \{tk^0\}\}$$

has the following properties:

(i) c^u is bounded on \mathcal{V} ;

(ii) $\forall V \in \mathcal{V} \forall \alpha \in \mathbb{R} : c^u(V + \alpha k^0) = c^u(V) + \alpha$;

(iii) $V_1 \preccurlyeq V_2 \Rightarrow c^u(V_1) \leq c^u(V_2)$;

(iv) If \mathcal{V} consists of compact sets $V \subset Y$ and $K \setminus \{0\} \subset \text{int } C$ then

$$\left(V_1 \preccurlyeq V_2, V_1 \cap V_2 = \emptyset \right) \Rightarrow c^u(V_1) < c^u(V_2).$$

Proof. Let V_0 be an upper \preccurlyeq -bound of \mathcal{V} , i.e. $W'' \subset Y$ is topologically bounded and $V \subset W'' - C$ for all $V \in \mathcal{V}$. Let U be a neighborhood of zero such that $U \subset -k^0 + \text{int } C$. Choosing $\sigma > 0$ such that $-W'' \subset \sigma U$ we obtain $V \subset W'' - C \subset -\sigma U - C \subset \sigma(k^0 - \text{int } C) - C \subset \sigma k^0 - C$. Hence $c^u(V) \leq \sigma$ for all $V \in \mathcal{V}$ i.e. c^u is bounded above on \mathcal{V} .

Assuming that c^u is not bounded below, for all $n \in \mathbb{N}$ we can find some $V_n \in \mathcal{V}$ such that $V_n \subset -nk^0 - C$. Since \mathcal{V} is weakly \preccurlyeq -bounded below there exists some nonempty set $W' \subset Y$ such that $V \not\subset W' - C$, hence $V \not\subset w' - C$ where $w' \in W'$ is arbitrarily chosen. Hence for all $n \in \mathbb{N}$ there exists $v_n \in V_n$ such that $v_n - w' \notin -C$ and $-v_n - nk^0 \in C$. We obtain

$$-k^0 - w'/n = -k^0 - v_n/n + (v_n - w')/n \subset C + (Y \setminus -\text{int } C) \subset Y \setminus -\text{int } C.$$

Letting $n \rightarrow \infty$ we get $k^0 \notin \text{int } C$ which contradicts the assumption $k^0 \in K \cap \text{int } C$.

Assertions (ii) and (iii) are obvious. It remains to prove (iv). By the definition of the infimum there exists a sequence $\{v_n\}_{n \in \mathbb{N}} \subset V_1$ such that $v_n \notin (c^u(V_1) - 1/n)k^0 - C$.

Since V_1 is compact, there is a subnet of $\{v_n\}_{n \in \mathbb{N}} \subset V_1$ converging to some $\bar{v} \in V_1$. Hence

$$\bar{v} \notin c^u(V_1)k^0 - \text{int } C. \quad (9)$$

Let $V_1 \preceq V_2$ and $V_1 \cap V_2 = \emptyset$. We have $\bar{v} \in V_2 - K \setminus \{0\} \subset V_2 - \text{int } C$. Choose $v \in V_2$ such that $\bar{v} \in v - \text{int } C$. Then there exists some $\delta > 0$ such that

$$\bar{v} \in v - \delta k^0 - \text{int } C. \quad (10)$$

Assuming that $v \in (c^u(V_1) + \delta)k^0 - C$ we obtain

$$\bar{v} \stackrel{(10)}{\in} v - \delta k^0 - \text{int } C \subset c^u(V_1)k^0 - C - \text{int } C \subset c^u(V_1)k^0 - \text{int } C.$$

This contradicts (9). Hence we have $v \notin (c^u(V_1) + \delta)k^0 - C$. Since $v \in V_2$ it follows $c^u(V_1) + \delta \leq c^u(V_2)$, i.e. $c^u(V_1) < c^u(V_2)$. \blacksquare

Let us discuss why the construction of Corollary 3.2 fails in the present setting. This is due to the fact that the boundedness assumptions are not longer symmetric: In Theorem 3.5 we supposed \mathcal{V} to be \preceq_C -bounded above and weakly \preceq_C -bounded below. This is true if and only if $-\mathcal{V}$ is \preceq_C -bounded below and weakly \preceq_C -bounded above. However, the following example shows that the weak \preceq_C -boundedness from above does not imply the boundedness of the functional c^l of Theorem 3.4.

Example 3.6 Let $Y = \mathbb{R}^2$, $C = \mathbb{R}_+^2$, $k^0 = (1, 1)$ and consider the set $\mathcal{V} = \{\{(-1, 1)\}, \{(-1, 2)\}, \{(-1, 3)\}, \dots\} \subset 2^Y$ consisting of singletons. Then $\{(0, 0)\}$ is a weak upper \preceq_C -bound and $\{(-1, 1)\}$ is a lower \preceq_C -bound of \mathcal{V} . But $c^l(\{(-1, n)\}) = n$ for all $n \in \mathbb{N}$, i.e. c^l is not bounded above on \mathcal{V} .

4 Basic Tools

For the convenience of the reader we present two basic tools for the proof of our minimal set theorems.

4.1 The Brézis-Browder Principle

The first tool is a very general existence principle for minimal points in quasi-ordered sets due to Brézis and Browder [1], 1976.

Theorem 4.1 *Let (W, \preceq) be a quasi-ordered set (i.e. \preceq is a reflexive and transitive relation on W) and let $\phi : W \rightarrow \mathbb{R}$ be a function satisfying*

(A1) ϕ is bounded below;

(A2) $w_1 \preceq w_2 \Rightarrow \phi(w_1) \leq \phi(w_2)$;

(A3) For every \preceq -decreasing sequence $\{w_n\}_{n \in \mathbb{N}} \subset W$ there exists some $w \in W$ such that $w \preceq w_n$ for all $n \in \mathbb{N}$.

Then, for every $w_0 \in W$ there exists some $\bar{w} \in W$ such that

(i) $\bar{w} \preceq w_0$;

(ii) $\hat{w} \preceq \bar{w} \Rightarrow \phi(\hat{w}) = \phi(\bar{w})$.

Proof. See [1, Corollary 1]. ■

4.2 Uniform Spaces

Our results involve a uniform space X (cf. [14]). Examples for uniform spaces not being necessarily metrizable are topological vector spaces and K -metric spaces (see [21] or [19]). If the reader is only interested in results for metric spaces the following considerations can be skipped. Then, one has to replace the “families of quasi-metrics”, in the following denoted by $\{q_\lambda\}_{\lambda \in \Lambda}$ or by q_Λ , by the metric.

In [11] we presented a characterization of uniform spaces via *families of quasi-metrics* introduced by Fang [4]. We shall give a short summary of these results.

Definition 4.2 Let X be a nonempty set and let (Λ, \prec) be a directed set. A system $\{q_\lambda\}_{\lambda \in \Lambda}$ of functions $q_\lambda : X \times X \rightarrow [0, \infty)$ satisfying

$$(Q1) \quad \forall \lambda \in \Lambda \quad \forall x \in X : q_\lambda(x, x) = 0;$$

$$(Q2) \quad \forall \lambda \in \Lambda \quad \forall x, y \in X : q_\lambda(x, y) = q_\lambda(y, x);$$

$$(Q3) \quad \forall \lambda \in \Lambda \quad \exists \mu \in \Lambda \text{ with } \lambda \prec \mu : \forall x, y, z \in X : q_\lambda(x, y) \leq q_\mu(x, z) + q_\mu(z, y);$$

$$(Q4) \quad \lambda \prec \mu \Rightarrow \left(\forall x, y \in X : q_\lambda(x, y) \leq q_\mu(x, y) \right)$$

is called a family of quasi-metrics. If, in addition, the condition

$$(Q5) \quad \left(\forall \lambda \in \Lambda : q_\lambda(x, y) = 0 \right) \Rightarrow x = y$$

is satisfied, the family of quasi-metrics is said to be separating.

Theorem 4.3 A topological space (X, τ) is a (separated) uniform space iff its topology τ can be generated by a (separating) family of quasi-metrics.

Proof. See [11]. ■

Convention 4.4 For an easy dealing with uniform spaces we introduce the following notation. Let $\{q_\lambda\}_{\lambda \in \Lambda}$ the family of quasi-metrics which generates the topology of the uniform space X . We write q_Λ iff an assertion holds for all $\lambda \in \Lambda$. If X is a metric space, then q_Λ is its metric.

5 Minimal Set Theorems

This section contains the main results of the paper. We present two minimal set theorems with respect to the set ordering relations introduced in Section 2. Let us consider a subset \mathcal{A} of $X \times 2^Y$, where X is a separated uniform space and Y is a topological vector space. We introduce the following notation:

$$\mathcal{V}(\mathcal{A}) := \{V \in 2^Y : \exists x \in X : (x, V) \in \mathcal{A}\}.$$

5.1 Minimal Set Theorem I

Using the relation \preceq we introduce the following ordering relation on $X \times 2^Y$:

$$(x_1, V_1) \preceq_{k^0} (x_2, V_2) \iff V_1 + k^0 q_\Lambda(x_1, x_2) \preceq V_2.$$

According to Convention 4.4, the last inequality has to be read as

$$\forall \lambda \in \Lambda : V_1 + k^0 q_\lambda(x_1, x_2) \preceq V_2.$$

The relation \preceq_{k^0} is a reflexive and transitive relation on $X \times 2^Y$. We present our *Minimal Set Theorem* involving \preceq_{k^0} .

Theorem 5.1 *Let X be a separated uniform space, Y a topological vector space, $K \subset Y$ a convex cone and $k^0 \in K \setminus -\text{cl} K$. Let \mathcal{A} be a nonempty subset of $X \times 2^Y$ such that for some $(x_0, V_0) \in \mathcal{A}$ and for $\mathcal{A}_0 := \{(x, V) \in \mathcal{A} : (x, V) \preceq_{k^0} (x_0, V_0)\}$ the following conditions are satisfied:*

- (M1) $\mathcal{V}(\mathcal{A}_0)$ is \preceq -bounded above, i.e. V_0 is nonempty;
- (M2) $\mathcal{V}(\mathcal{A}_0)$ is \preceq -bounded below;
- (M3) For every \preceq_{k^0} -decreasing sequence $\{(x_n, V_n)\}_{n \in \mathbb{N}} \subset \mathcal{A}_0$ there exists some $(x, V) \in \mathcal{A}_0$ such that $(x, V) \preceq_{k^0} (x_n, V_n)$ for all $n \in \mathbb{N}$.

Then, there exists $(\bar{x}, \bar{V}) \in \mathcal{A}$ such that

- (i) $(\bar{x}, \bar{V}) \preceq_{k^0} (x_0, V_0)$;
- (ii) $\left((\hat{x}, \hat{V}) \in \mathcal{A}, (\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V}) \right) \Rightarrow \hat{x} = \bar{x}$.

Under the additional assumption (C) and if $k^0 \in K \cap \text{int} C$, (M2) can be relaxed to

- (M2') $\mathcal{V}(\mathcal{A}_0)$ is weakly \preceq_C -bounded below.

If, in addition, $k^0 \in K \setminus \{0\} \subset \text{int} C$ and if for each $(x, V) \in \mathcal{A}_0$, V is compact, then (ii) can be strengthened to

- (ii') $\left((\hat{x}, \hat{V}) \in \mathcal{A}, (\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V}) \right) \Rightarrow \left(\hat{x} = \bar{x} \text{ and } \hat{V} \cap \bar{V} \neq \emptyset \right)$.

Proof. We shall apply the Brézis-Browder principle to the quasi-ordered set $(\mathcal{A}_0, \preceq_{k^0})$ and the functional

$$\phi : \mathcal{A}_0 \rightarrow \mathbb{R}, \quad \phi(x, V) := z^l(V),$$

where $z^l : \mathcal{V}(\mathcal{A}_0) \rightarrow \mathbb{R}$ is the scalarization functional of Theorem 3.1. In the definition of z^l the upper \preceq -bound V'' has to be replaced by V_0 .

We have to check the assumptions of Theorem 4.1. By (M1), (M2) and Theorem 3.1 (i), ϕ is well-defined and bounded. Theorem 3.1 (ii) and (iii) yield

$$(x_1, V_1) \preceq_{k^0} (x_2, V_2) \Rightarrow \phi(x_1, V_1) + q_\Lambda(x_1, x_2) \leq \phi(x_2, V_2). \quad (11)$$

Hence, ϕ is \preceq_{k^0} -monotone on \mathcal{A}_0 , i.e. assumption (A2) of Theorem 4.1 is satisfied. Of course, (M3) implies assumption (A3).

Theorem 4.1 yields the existence of an element $(\bar{x}, \bar{V}) \in \mathcal{A}_0$ (i.e. (i) holds) such that

$$\left((\hat{x}, \hat{V}) \in \mathcal{A}_0, (\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V}) \right) \Rightarrow \phi(\hat{x}, \hat{V}) = \phi(\bar{x}, \bar{V}). \quad (12)$$

Let $(\hat{x}, \hat{V}) \in \mathcal{A}$ such that $(\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V})$. The transitivity of \preceq_{k^0} yields $(\hat{x}, \hat{V}) \in \mathcal{A}_0$. Applying (12) and (11) we obtain $q_\Lambda(\hat{x}, \bar{x}) = 0$. Since X is separated, we have $\hat{x} = \bar{x}$, i.e. (ii) holds.

To see that (M2) can be replaced by (M2') we proceed as above but using the functional c^l of Theorem 3.4 instead of z^l in the Definition of ϕ . To prove (ii') assume that $\hat{V} \cap \bar{V} = \emptyset$. Then, (iv) of Theorem 3.4 yields $\phi(\hat{x}, \hat{V}) < \phi(\bar{x}, \bar{V})$. This contradicts (12). ■

5.2 Minimal Set Theorem II

Using the \preceq -relation we present a second minimal set theorem. We introduce the following ordering relation on $X \times 2^Y$:

$$(x_1, V_1) \preceq_{k^0} (x_2, V_2) \Leftrightarrow V_1 + k^0 q_\Lambda(x_1, x_2) \preceq V_2$$

which is also a reflexive and transitive relation.

Theorem 5.2 *Let X be a separated uniform space, Y a topological vector space, $K \subset Y$ a convex cone and $k^0 \in K \setminus -\text{cl} K$. Let \mathcal{A} be a nonempty subset of $X \times 2^Y$ such that for some $(x_0, V_0) \in \mathcal{A}$ and for $\mathcal{A}_0 := \{(x, V) \in \mathcal{A} : (x, V) \preceq_{k^0} (x_0, V_0)\}$ the following conditions are satisfied:*

(M1) $\mathcal{V}(\mathcal{A}_0)$ is \preceq -bounded above;

(M2) $\mathcal{V}(\mathcal{A}_0)$ is \preceq -bounded below;

(M3) For every \preceq_{k^0} -decreasing sequence $\{(x_n, V_n)\}_{n \in \mathbb{N}} \subset \mathcal{A}_0$ there exists some $(x, V) \in \mathcal{A}_0$ such that $(x, V) \preceq_{k^0} (x_n, V_n)$ for all $n \in \mathbb{N}$.

Then, there exists $(\bar{x}, \bar{V}) \in \mathcal{A}$ such that

- (i) $(\bar{x}, \bar{V}) \preceq_{k^0} (x_0, V_0)$;
- (ii) $\left((\hat{x}, \hat{V}) \in \mathcal{A}, (\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V}) \right) \Rightarrow \hat{x} = \bar{x}$.

Under the additional assumption (C) and if $k^0 \in K \cap \text{int } C$, (M1), (M2) can be relaxed to

- (M1') $\{V_0\}$ is \preceq_C -bounded above;
- (M2') $\mathcal{V}(\mathcal{A}_0)$ is weakly \preceq_C -bounded below.

If, in addition, $k^0 \in K \setminus \{0\} \subset \text{int } C$ and if for each $(x, V) \in \mathcal{A}_0$, V is compact, then (ii) can be strengthened to

- (ii') $\left((\hat{x}, \hat{V}) \in \mathcal{A}, (\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V}) \right) \Rightarrow (\hat{x} = \bar{x} \text{ and } \hat{V} \cap \bar{V} \neq \emptyset)$.

Proof. We shall apply the Brézis-Browder principle to the quasi-ordered set $(\mathcal{A}_0, \preceq_{k^0})$ and the functional

$$\phi : \mathcal{A}_0 \rightarrow \mathbb{R}, \phi(x, V) := z_u(V)$$

where $z_u : \mathcal{V}(\mathcal{A}_0) \rightarrow \mathbb{R}$ is the scalarization functional of Corollary 3.2. In the definition of z_u , the set W' has to be a lower \preceq -bound of $\mathcal{V}(\mathcal{A}_0)$ which exists according to (M2).

We have to check the assumptions of Theorem 4.1. By (M1), (M2) and Corollary 3.2, ϕ is well-defined and bounded on \mathcal{A}_0 . Corollary 3.2 yields

$$(x_1, V_1) \preceq_{k^0} (x_2, V_2) \Rightarrow \phi(x_1, V_1) + q_\Lambda(x_1, x_2) \leq \phi(x_2, V_2). \quad (13)$$

It follows that ϕ is \preceq_{k^0} -monotone on \mathcal{A}_0 , i.e. assumption (A2) of Theorem 4.1 is satisfied. Of course, (M3) implies assumption (A3). Theorem 4.1 yields the existence of an element $(\bar{x}, \bar{V}) \in \mathcal{A}_0$ (i.e. (i) holds) such that

$$\left((\hat{x}, \hat{V}) \in \mathcal{A}_0, (\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V}) \right) \Rightarrow \phi(\hat{x}, \hat{V}) = \phi(\bar{x}, \bar{V}). \quad (14)$$

Let $(\hat{x}, \hat{V}) \in \mathcal{A}$ such that $(\hat{x}, \hat{V}) \preceq_{k^0} (\bar{x}, \bar{V})$. The transitivity of \preceq_{k^0} yields $(\hat{x}, \hat{V}) \in \mathcal{A}_0$. Applying (14) and (13) we obtain $q_\Lambda(\hat{x}, \bar{x}) = 0$. Since X is separated, we have $\hat{x} = \bar{x}$, i.e. (ii) holds.

To see that (M1) and (M2) can be replaced by (M1') and (M2') proceed as above but using the functional c^u of Theorem 3.5 instead of z^u in the definition of ϕ .

To prove (ii') assume that $\hat{V} \cap \bar{V} = \emptyset$. Theorem 3.5 (iv) yields $\phi(\hat{x}, \hat{V}) < \phi(\bar{x}, \bar{V})$. This contradicts (14). ■

Note that Theorem 5.2 can be transformed into a *Maximal Set Theorem* with respect to the reflexive and transitive relation \preceq_{-k^0} on $X \times 2^Y$ defined by

$$(x_1, V_1) \preceq_{-k^0} (x_2, V_2) \iff V_1 - k^0 q_\Lambda(x_1, x_2) \preceq_{-K} V_2$$

observing that a sequence $\{(x_n, V_n)\}_{n \in \mathbb{N}} \subset \mathcal{A}_0$ is \preceq_{k^0} -decreasing if and only if it is \preceq_{-k^0} -increasing. One can see that it is not possible to obtain Theorem 5.2 from Theorem 5.1 by replacing \preceq_{k^0} by \preceq_{-k^0} . We face the alternative either to state minimal set theorems for both the relations \preceq_{k^0} , \preceq_{-k^0} or to state a minimal as well as a maximal set theorem involving one of the relations.

6 Set-Valued Ekeland's Principles

In this section, we present several conclusions of Theorem 5.1 and 5.2. Recently, Truong [24] proved a variant of Ekeland's variational principle involving only the \preceq -relation. A similar, but more general variant can be obtained from Theorem 5.1. A new variational principle will be derived from Theorem 5.2. Another variant of Ekeland's principle for set-valued maps (cf. [2], [11]) as well as a minimal point theorem in $X \times Y$ turn out to be consequences of Theorem 5.1 as well as of Theorem 5.2.

Let X be a set, Y a linear topological space, $F : X \rightarrow 2^Y$ a set-valued map and K a convex cone in Y . In contrast to known definitions we call the set

$$\text{graph } F := \{(x, V) \in X \times 2^Y : V = F(x)\}$$

the *graph* of F and the *image* of a subset $M \subset X$ (see also [24]) is defined by

$$F(M) := \{F(x) : x \in M\}.$$

Note that $\text{graph } F$ is a subset of $X \times 2^Y$, not of $X \times Y$ and the image $F(M)$ is a subset of 2^Y , not of Y !

Moreover, we shall introduce the concept of the *domain* of a set-valued map F in a suitable way for each of the relations \preceq , \preceq . We define

$$\begin{aligned} \preceq\text{-dom } F &:= \{x \in X : F(x) \preceq V \text{ for some nonempty } V \subset Y\} \\ \preceq\text{-dom } F &:= \{x \in X : F(x) \preceq V \text{ for some topologically bounded } V \subset Y\}. \end{aligned}$$

Clearly, $x \in \preceq\text{-dom } F$ means that the set $\{F(x)\} \subset 2^Y$ consisting of just one element is \preceq -bounded above which is equivalent to $F(x) \neq \emptyset$. Similarly, $x \in \preceq\text{-dom } F$ means that the set $\{F(x)\} \subset 2^Y$ consisting of just one element is \preceq -bounded above.

6.1 Ekeland's Principle for Set-Valued Maps I

First, we state a variational principle involving the ordering relation \preceq .

Theorem 6.1 *Let X be a separated uniform space, Y a topological vector space, $K \subset Y$ a convex cone and $k^0 \in K \setminus -\text{cl } K$. Let $F : X \rightarrow 2^Y$ be a set-valued mapping, $x_0 \in \preceq\text{-dom } F$, $S(x_0) := \{x \in X : F(x) + k^0 q_\Lambda(x, x_0) \preceq F(x_0)\}$ and $\mathcal{A}_0 := \{(x, V) \in \text{graph } F : x \in S(x_0)\}$ such that the following conditions are satisfied:*

(E1) $F(S(x_0))$ is \preceq -bounded below;

(E2) For every \preceq_{k^0} -decreasing sequence $\{(x_n, V_n)\}_{n \in \mathbb{N}} \subset \mathcal{A}_0$ there exists some $(x, V) \in \mathcal{A}_0$ such that $(x, V) \preceq_{k^0} (x_n, V_n)$ for all $n \in \mathbb{N}$.

Then, there exists $\bar{x} \in \preceq\text{-dom } F$ such that

- (i) $F(\bar{x}) + k^0 q_\Lambda(\bar{x}, x_0) \preceq F(x_0)$;
- (ii) $\forall x \neq \bar{x} : \exists \lambda \in \Lambda : F(x) + k^0 q_\Lambda(x, \bar{x}) \not\preceq F(\bar{x})$.

Under the additional assumption (C) and if $k^0 \in K \cap \text{int } C$, (E1) can be relaxed to

(E1') $F(S(x_0))$ is weakly \preceq_C -bounded below;

Proof. Set $\mathcal{A} := \text{graph } F$ and apply Theorem 5.1. It only remains to note that $x_0 \in \preceq\text{-dom } F$ implies condition (M1) of Theorem 5.1. \blacksquare

Condition (ii) tells us that there does not exist an $x \in X \setminus \{\bar{x}\}$ such that $F_\lambda(x) \preceq F_\lambda(\bar{x}) = F(\bar{x})$ where

$$F_\lambda : X \rightarrow 2^Y, \quad F_\lambda(x) := F(x) + k^0 q_\Lambda(x, \bar{x}).$$

This means, \bar{x} is an s-minimizer in the sense of [25] of F_λ . Of course, $F_\lambda(\bar{x})$ is also a minimal element of $\{F_\lambda(x) : x \in X\}$ with respect to \preceq .

The assumptions of Theorem 6.1 may look somewhat artificial. We give a sufficient condition for (E2).

Theorem 6.2 *Let $X, Y, K, k^0, F, x_0, S(x_0), \mathcal{A}_0$ be as in Theorem 6.1 and let (E1) be satisfied. Then (E2) is in force if the following condition is satisfied:*

(E2') For every $x \in X$ the set $S(x) := \{x' \in X : F(x') + k^0 q_\Lambda(x', x) \preceq F(x)\}$ is sequentially complete in X .

Proof. Let $\{(x_n, V_n)\}_{n \in \mathbb{N}} \subset \mathcal{A}_0$ be a \preceq_{k^0} -decreasing sequence, i.e. $V_n = F(x_n)$ and

$$F(x_{n+1}) + k^0 q_\Lambda(x_{n+1}, x_n) \preceq F(x_n) \tag{15}$$

for all $n \in \mathbb{N}$. By the transitivity of \preceq_{k^0} we get

$$F(x_m) + k^0 q_\Lambda(x_m, x_n) \preceq F(x_n) \tag{16}$$

for all $m \geq n, m \in \mathbb{N}$. Applying the functional $\phi : \mathcal{A}_0 \rightarrow \mathbb{R}$ from the proof of Theorem 5.1 to relation (15) we obtain

$$\phi(x_{n+1}, F(x_{n+1})) + q_\Lambda(x_{n+1}, x_n) \leq \phi(x_n, F(x_n)).$$

The sequence $\{\phi(x_n, F(x_n))\}_{n \in \mathbb{N}}$ is nonincreasing and bounded below by (E1) and the corresponding properties of ϕ , hence convergent and a fortiori a Cauchy sequence. Applying ϕ to (16) we may conclude that $\{x_n\}_{n \in \mathbb{N}}$ is Cauchy as well. Since $S(x_0)$ is sequentially complete, $\{x_n\}_{n \in \mathbb{N}}$ converges to some $x \in S(x_0)$. Moreover, (16) implies that $x_m \in S(x_n)$ for all $m \geq n, m \in \mathbb{N}$. Since by (E2') $S(x_n)$ is sequentially complete we have $x \in S(x_n)$ for all $n \in \mathbb{N}$. This means $(x, F(x)) \preceq_{k^0} (x_n, F(x_n))$ as desired. \blacksquare

Remark 6.3 We indicate a special situation where (E2') is satisfied. Let X be a sequentially complete separated uniform space. Ferro [5] introduced the concept of lower \preceq -semicontinuity (D-lower semicontinuity in [5]) and Truong [24] proved an Ekeland-type theorem using this continuity property. A set-valued map $F : X \rightarrow 2^Y$ is said to be sequentially lower \preceq -semicontinuous iff for every $V \in 2^Y$ the set $\{x' \in X : F(x') \preceq V\}$ is a sequentially closed subset of X . We claim that (E2') is satisfied if F is sequentially lower \preceq -semicontinuous with \preceq -closed values (for each $x \in X$, $F(x) + K$ is a closed set). Indeed, let $\{x_n\}_{n \in \mathbb{N}} \subset S(x)$ be a sequence such that $x_n \rightarrow x'$. Fix $\lambda \in \Lambda$. Then there exists $\mu \in \Lambda$ such that $q_\lambda(x', x) \leq q_\mu(x', x_n) + q_\mu(x_n, x)$. Since $x_n \rightarrow x'$ we can find for each $\varepsilon > 0$ a number $n_\varepsilon \in \mathbb{N}$ such that $q_\mu(x', x_n) \leq \varepsilon$ for all $n \in \mathbb{N}$, $n > n_\varepsilon$. Hence $q_\mu(x_n, x) \geq q_\lambda(x', x) - \varepsilon$. Since $F(x_n) + k^0 q_\mu(x_n, x) \preceq F(x)$, for $n > n_\varepsilon$ holds $F(x_n) + k^0 (q_\lambda(x', x) - \varepsilon) \preceq F(x)$. This implies $F(x_n) \preceq F(x) - k^0 (q_\lambda(x', x) - \varepsilon)$ for $n > n_\varepsilon$. Since F is lower \preceq -semicontinuous we have $F(x') \preceq F(x) - k^0 (q_\lambda(x', x) - \varepsilon)$, hence

$$F(x) \subset F(x') + k^0 (q_\lambda(x', x) - \varepsilon) + K. \quad (17)$$

Take $y \in F(x)$. Then, from (17) it follows that $y + \varepsilon k^0 \in F(x') + k^0 q_\lambda(x', x) + K$. Letting $\varepsilon \rightarrow 0$ we obtain $y \in F(x') + k^0 q_\lambda(x', x) + K$ since the latter set is closed. Hence $x' \in S(x)$.

Remark 6.4 In [24] the concept of K -boundedness (cf. [20]) is defined as follows: A subset A of a linear topological space Y is said to be K -bounded if there is a topologically bounded set $M \subset Y$ such that $A \subset M + K$. It can easily be seen that the boundedness condition in [24] coincides with condition (E1) of Theorem 6.1.

In [24], Y is supposed to be a locally convex space, K is a closed pointed convex cone and $k^0 \in K \setminus \{0\}$. Therefore we have $k^0 \in K \setminus -\text{cl} K$. Remark 3.3 yields that assumption (C) is satisfied and $k^0 \in K \cap \text{int} C$. Therefore, these assumptions have not to be proposed additionally. Hence, our boundedness condition (E1') is in fact weaker than the boundedness condition in [24] (cf. Remark 2.6 and Example 2.7). This extends the area of applicability of Ekeland's principle as the following example shows.

Example 6.5 Let $X = [0, \infty)$, $Y = \mathbb{R}^2$ with Euclidean norm, $B_r(y)$ be the closed ball of radius $r \geq 0$ centered at $y \in Y$, $K = C = \mathbb{R}_+^2$ and

$$F : X \rightarrow 2^Y, \quad F(x) := B_x(0) \setminus -\text{int} K.$$

One may check that F is not \preceq -bounded, but $(-1, -1)^T$ is a weak lower \preceq -bound. Hence the results of [24] are not applicable. Moreover, there does not exist a \preceq -minimal value of F , but it is \preceq -lower semicontinuous. Thus, we can apply Theorem 6.2 in combination with Remark 6.3.

Theorem 6.2 and Remark 6.3–6.6 show that Theorem 5.1 of [24] is a very special case of Theorem 6.1 concerning the properties of F : We use (E2) instead of K-lower

semicontinuity and also weaker boundedness assumptions, cf. Remark 6.4. Of course, we deal with larger classes of spaces X (uniform instead of metric spaces) and Y (linear topological instead of locally convex spaces) as well as of cones in Y (not necessarily closed and pointed).

Remark 6.6 If $x_0 \in X$ is an εk^0 -minimal point of F in the sense of [24], i.e. $F(x) \not\preceq F(x_0) - \varepsilon k^0$ then relation (i) of Theorem of 6.1 can be split into the two relations (i₁) $F(\bar{x}) \preceq F(x_0)$; (i₂) $q_\Lambda(\bar{x}, x_0) \leq \varepsilon$. Indeed, while (i₁) is immediate, (i) is equivalent to $F(x_0) \subset F(\bar{x}) + k^0 q_\Lambda(\bar{x}, x_0) + K$. If $q_\mu(\bar{x}, x_0) > \varepsilon$ for some $\mu \in \Lambda$, we have

$$F(x_0) \subset F(\bar{x}) + k^0 (q_\mu(\bar{x}, x_0) - \varepsilon) + \varepsilon k^0 + K \subset F(\bar{x}) + \varepsilon k^0 + K$$

which contradicts the εk^0 -minimality of x_0 .

The following lemma tells us that there are always \preceq - k^0 -minimal solution if F is \preceq -bounded below.

Lemma 6.7 *Let X, Y, K and k_0 as in Theorem 6.1, $F : X \rightarrow 2^Y$. If F is \preceq -bounded below, then there exists $x_0 \in X$ such that*

$$\forall x \in X : F(x) \not\preceq F(x_0) - k^0.$$

Proof. Assume the contrary, namely, for all $x_0 \in X$ there exists some $x \in X$ such that $F(x) \preceq F(x_0) - k^0$. By induction we can construct a sequence $\{x_n\}_{n \in \mathbb{N}}$ such that $F(x_n) \preceq F(x_0) - nk^0$ for all $n \in \mathbb{N}$. Since $F(x_0) - nk^0 \preceq F(x_0)$ and by assumption we conclude that $\mathcal{V} := \{F(x_n) : n \in \mathbb{N}\}$ is \preceq -bounded. Applying the functional z^l of Theorem 3.1 to the above inequality we get $z^l(F(x_n)) \leq z^l(F(x_0)) - n$ for all $n \in \mathbb{N}$, i.e. z^l is not bounded below on \mathcal{V} . This contradicts (i) of Theorem 3.1 saying that z^l is bounded on \mathcal{V} . ■

Note that, involving assumption (C), it is enough to suppose that F is weakly \preceq -bounded below (compare Theorem 6.1). The same applies for the following theorem, the classical form of Ekeland's principle, see Theorem 1bis in [3].

Theorem 6.8 *Let the assumptions of Theorem 6.1 be satisfied. Then there exists $\bar{x} \in X$ such that*

$$\forall x \in X : F(x) \not\preceq F(\bar{x}) - k^0 \tag{18}$$

$$\forall x \in X, x \neq \bar{x}, \exists \lambda \in \Lambda : F(x) + k^0 q_\lambda(x, \bar{x}) \not\preceq F(\bar{x}). \tag{19}$$

Proof. According to Lemma 6.7, (1) there exists $x_0 \in X$ such that

$$\forall x \in X : F(x) \not\preceq F(x_0) - k^0. \tag{20}$$

Applying Theorem 6.1 and taking into account Remark 6.6 we obtain $\bar{x} \in X$ satisfying (19) as well as $F(\bar{x}) \preceq F(x_0)$. If $x' \in X$, $F(x') \preceq F(\bar{x}) - k^0$ we obtain by transitivity $F(x') \preceq F(x_0) - k^0$ contradicting (20). Hence (18) is true. ■

6.2 Ekeland's Principle for Set-Valued Maps II

In this section, we shall prove an analogous variational principle for the relation \preceq .

Theorem 6.9 *Let X be a separated uniform space, Y a topological vector space, $K \subset Y$ a convex cone and $k^0 \in K \setminus -\text{cl}K$. Let $F : X \rightarrow 2^Y$ be a set-valued mapping, $x_0 \in \preceq\text{-dom} F$, $S(x_0) := \{x \in X : F(x) + k^0 q_\Lambda(x, x_0) \preceq F(x_0)\}$ and $\mathcal{A}_0 := \{(x, V) \in \text{graph } F : x \in S(x_0)\}$, such that the following conditions are satisfied:*

(E1) $F(S(x_0))$ is \preceq -bounded below;

(E2) For every \preceq_{k^0} -decreasing sequence $\{(x_n, V_n)\}_{n \in \mathbb{N}} \subset \mathcal{A}_0$ there exists some $(x, V) \in \mathcal{A}_0$ such that $(x, V) \preceq_{k^0} (x_n, V_n)$ for all $n \in \mathbb{N}$.

Then, there exists $\bar{x} \in \preceq\text{-dom} F$ such that

(i) $F(\bar{x}) + k^0 q_\Lambda(\bar{x}, x_0) \preceq F(x_0)$;

(ii) $\forall x \neq \bar{x} : \exists \lambda \in \Lambda : F(x) + k^0 q_\Lambda(x, \bar{x}) \not\preceq F(\bar{x})$.

Under the additional assumption (C) and if $k^0 \in K \cap \text{int} C$, then we can even allow $x_0 \in \preceq_C\text{-dom} F$ and (E1) can be replaced by

(E1') $F(S(x_0))$ is weakly \preceq_C -bounded below.

Proof. Set $A := \text{graph } F$ and apply Theorem 5.2 noting that $x_0 \in \preceq\text{-dom} F$ and $x_0 \in \preceq_C\text{-dom} F$ imply (M1) and (M1') of Theorem 5.2, respectively. ■

Theorem 6.10 *Let $X, Y, K, k^0, F, S(x_0), \mathcal{A}_0$ be as in Theorem 6.9 and let (E1) be satisfied. Then (E2) is in force if the following condition is satisfied:*

(E2') For every $x \in X$ the set $S(x) := \{x' \in X : F(x') + k^0 q_\Lambda(x', x) \preceq F(x)\}$ is sequentially complete in X .

Proof. Follow the lines of the proof of Theorem 6.2. ■

Remark 6.11 A map $F : X \rightarrow 2^Y$ is called sequentially lower \preceq -semicontinuous iff for every $V \in 2^Y$ the set $\{x' \in X : F(x') \preceq V\}$ is a sequentially closed subset of X . Following arguments similar to those in Remark 6.3, we can prove that (E2') of Remark 6.10 is satisfied if F is sequentially lower \preceq -semicontinuous with \preceq -closed values (for each $x \in X$, $F(x) - K$ is a closed set).

Remark 6.12 Considerations similar to Remark 6.6, Lemma 6.7 and Theorem 6.8 can be done.

6.3 Ekeland's Principle with Point Relation and a Minimal Point Theorem

In [11] we proved a minimal point theorem and its equivalence to a variant of Ekeland's principle for set-valued maps as well as many conclusions of them (e.g. results of [2], [10], [9], [13], [18]). In this subsection we show that these theorems are corollaries of Theorem 5.1 as well as of Theorem 5.2. We are concerned with elements $w = (w_X, w_Y) = (x, y)$ of the product space $W = X \times Y$, X a separated uniform space, Y a topological vector space. We introduce the ordering relation \preceq_{k^0} on W using an element $k^0 \in K \setminus -\text{cl } K$:

$$(x_1, y_1) \preceq_{k^0} (x_2, y_2) \Leftrightarrow y_1 + k^0 q_\Lambda(x_1, x_2) \leq_K y_2. \quad (21)$$

If K is a convex cone, \preceq_{k^0} is a reflexive and transitive relation. If K additionally is pointed, the relation is also antisymmetric. See e.g. [10], [11]. Identifying an element $(x, y) \in X \times Y$ with $(x, \{y\}) \in X \times 2^Y$, it is easy to see that \preceq_{k^0} coincides with the above defined ordering \preceq_{k^0} as well as with \preccurlyeq_{k^0} .

The following corollary is a variant of Ekeland's variational principle for set-valued maps involving the ordering relation \preceq_{k^0} applied to elements of the set

$$\text{gr } F := \{(x, y) \in X \times Y : y \in F(x)\}$$

usually denoting the graph of F . Note the difference to the definition at the beginning of this section.

Corollary 6.13 *Let X be a separated uniform space, Y a topological vector space, $K \subset Y$ a convex cone and $k_0 \in K \setminus -\text{cl } K$. For the set-valued mapping $F : X \rightarrow 2^Y$, let $w_0 = (x_0, y_0) \in \text{gr } F$ be given such that for the set $A_0 := \{w \in \text{gr } F : w \preceq_{k^0} w_0\}$ the following assumptions are satisfied:*

- (E1) *The set $(A_0)_Y := \{y \in Y : \exists x \in X : w = (x, y) \in A_0\}$ is \leq_K -bounded below;*
- (E2) *For every \preceq_{k^0} -decreasing sequence $\{(x_n, y_n)\}_{n \in \mathbb{N}} \subset A_0$ there exists some point $(x, y) \in A_0$ such that $(x, y) \preceq_{k^0} (x_n, y_n)$ for all $n \in \mathbb{N}$.*

Then, there exists some point $(\bar{x}, \bar{y}) \in \text{gr } F$ such that

- (i) $\bar{y} + k^0 q_\Lambda(\bar{x}, x_0) \leq_K y_0$;
- (ii) $\forall (x, y) \in \text{gr } F$ with $x \neq \bar{x} \exists \lambda \in \Lambda : y + k^0 q_\Lambda(x, \bar{x}) \not\leq_K \bar{y}$.

Under the additional assumption (C) and if $k^0 \in K \cap \text{int } C$, (E1) can be relaxed to

- (E1') *There exists some $\tilde{y} \in Y$ such that $(A_0)_Y \cap (\tilde{y} - \text{int } C) = \emptyset$;*

and if $k^0 \in K \setminus \{0\} \subset \text{int } C$ then \bar{y} is \leq_K -minimal in $F(\bar{x})$.

Proof. Choose $\mathcal{A} = \{(x, \{y\}) : (x, y) \in \text{gr } F\}$ and apply Theorem 5.1 or 5.2. ■

The following corollary is a minimal point theorem in $X \times Y$. Compare Theorem 9 of [11].

Corollary 6.14 *Let X be a separated uniform space, Y a topological vector space, $K \subset Y$ a convex cone and $k^0 \in K \setminus -\text{cl} K$. Let $A \subset W = X \times Y$ such that for some $w_0 \in A$ and for $A_0 := \{w \in A : w \preceq_{k^0} w_0\}$ the following conditions are satisfied:*

(M2) *The set $(A_0)_Y := \{y \in Y : \exists x \in X : (x, y) \in A_0\}$ is \leq_K -bounded below;*

(M3) *For every \preceq_{k^0} -decreasing sequence $\{w_n\}_{n \in \mathbb{N}} \subset A_0$ there exists some $w \in A_0$ such that $w \preceq_{k^0} w_n$ for all $n \in \mathbb{N}$.*

Then, there exists $\bar{w} \in A$ such that

(i) $\bar{w} \preceq_{k^0} w_0$;

(ii) $(\hat{w} \in A, \hat{w} \preceq_{k^0} \bar{w}) \Rightarrow \hat{w}_X = \bar{w}_X$.

Under the additional assumption (C) and if $k^0 \in K \cap \text{int} C$, (M2) can be replaced by

(M2') *There exists some $\tilde{y} \in Y$ such that $(A_0)_Y \cap (\tilde{y} - \text{int} C) = \emptyset$.*

If, in addition, $k^0 \in K \setminus \{0\} \subset \text{int} C$ then (ii) can be strengthened to

(ii') $(\hat{w} \in A, \hat{w} \preceq_{k^0} \bar{w}) \Rightarrow \hat{w} = \bar{w}$, (i.e. \bar{w} is \preceq_{k^0} -minimal in A).

Proof. Choose $\mathcal{A} = \{(x, \{y\}) : (x, y) \in A\}$ and apply Theorem 5.1 or 5.2. ■

The equivalence of Corollary 6.13 and Corollary 6.14 as well as a fixed point theorem of Kirk-Caristi type has been established in [11]. Because of this equivalence the minimal point theorem in $X \times Y$ can be understood as Ekeland's principle for set-valued maps with respect to an ordering relation for elements of $\text{gr} F \subset X \times Y$.

6.4 Ekeland's Principle for Single-valued Maps

Finally, we present a conclusion for the case of a single-valued map $f : X \rightarrow Y$. As above, identifying an element $y \in Y$ with $\{y\} \in 2^Y$ the ordering relations \preceq and \preccurlyeq coincide, hence the following corollary may be deduced from Theorem 6.1 as well as from Theorem 6.9. It is also possible to derive it from Corollary 6.13. As proposed in [10], [9], we extend the space Y by an element ∞ such that $y \leq_K \infty$ for all $y \in Y$. As usually, the *domain* of f is said to be the set $\text{dom} f := \{x \in X : f(x) \neq \infty\}$.

Corollary 6.15 *Let X be a separated uniform space, Y a topological vector space, $K \subset Y$ a convex cone and $k_0 \in K \setminus -\text{cl} K$. Let $f : X \rightarrow Y \cup \{\infty\}$ be a proper function satisfying the conditions*

(E1) *f is \leq_K -bounded below;*

(E2) *For every $x \in \text{dom} f$ the set $S(x) := \{x' \in X : f(x') + k_0 q_\Lambda(x', x) \leq_K f(x)\}$ is sequentially complete.*

Then, for each $x_0 \in \text{dom } f$ there exists $\bar{x} \in X$ such that

(i) $f(\bar{x}) + k^0 q_\Lambda(\bar{x}, x_0) \leq_K f(x_0)$;

(ii) $\forall x \in X \setminus \{\bar{x}\} : \exists \lambda \in \Lambda : f(x) + k^0 q_\lambda(x, \bar{x}) \not\leq_K f(\bar{x})$.

Under the additional assumption (C) and if $k^0 \in K \cap \text{int } C$, (E1) can be replaced by

(E1') There exists some $\tilde{y} \in Y$ such that $f(S(x_0)) \cap (\tilde{y} - \text{int } C) = \emptyset$.

Proof. To apply Corollary 6.13 or Theorem 6.1 (and Remark 6.2) we may define a set-valued function $F : X \rightarrow 2^Y$ by setting $F(x) := \{f(x)\}$ if $f(x) \neq \infty$ and $F(x) := \emptyset$ else.

A third variant of a proof can be given by setting $F(x) := \{f(x)\}$ if $f(x) \neq \infty$ and $F(x) := Y$ else and applying Theorem 6.9 (and Remark 6.10). ■

Taking into account Remark 6.6 and 6.12, respectively, we see that Corollary 6.15 is a generalization of Theorem 4 of [13] as well as Theorem 4 of [18]. It also covers Corollary 2 of [10] as well as Corollary 2 of [9]. This shows that our minimal set theorems are powerful generalizations of the minimal point theorems of [22], [9], [10], [11].




Figure 1: Relationships between the main results

References

- [1] Brézis, H., Browder, F. E., *A General Principle on Ordered Sets in Nonlinear Functional Analysis*, Advances in Mathematics 21, 355-364, (1976)
- [2] Chen, G. Y., Huang, X. X., Hou, S. H., *General Ekeland's Variational Principle for Set-Valued Mappings*, JOTA, Vol. 106, No. 1, pp. 151-164, (2000)
- [3] Ekeland, I., *Nonconvex Minimization Problems*, Bull. Americ. Math. Soc., Vol. 1, No. 3, pp. 443-474 (1979)
- [4] Fang, X.-J., *The Variational Principle and Fixed Point Theorems in Certain Topological Spaces*, J. Math. Anal. Appl. 202, 398-412, (1996)
- [5] Ferro, F., *An Optimization Result for Set-valued Mappings and a Stability Property in Vector Problems with Constraints*, JOTA, Vol. 90, pp. 63-77, (1996)
- [6] Gerstewitz (Tammer), Chr., *Nichtkonvexe Dualität in der Vektoroptimierung*, Wiss. Z. TH Leuna-Merseburg, 25(3), 357-364, (1983)
- [7] Gerth (Tammer), Chr., Weidner, P., *Nonconvex Separation Theorems and Some Applications in Vector Optimization*, JOTA, Vol. 67, No. 2, 297-320, (1990)
- [8] Göpfert, A., Tammer, Chr., *A New Maximal Point Theorem*, ZAA 14 (2), 379-390, (1995)
- [9] Göpfert, A., Tammer, Chr., Zălinescu, C., *A New Minimal Point Theorem in Product Spaces*, ZAA 18 (3), 767-770, (1999)
- [10] Göpfert, A., Tammer, Chr., Zălinescu, C., *On the Vectorial Ekeland's Variational Principle and Minimal Points in Product Spaces*, Nonlinear Analysis 39, 909-922, (2000)
- [11] Hamel, A., Löhne, A., *A Minimal Point Theorem in Uniform Spaces*, Report 09, University of Halle-Wittenberg, Institute of Optimization and Stochastics, (2002)
- [12] Huang, X. X., *A New Variant of Ekeland's Variational Principle for Set-Valued Maps*, Optimization Vol. 52, No. 1, pp.53-63, (2003)
- [13] Isac, G., *The Ekeland's Principle and the Pareto ε -Efficiency*, in: M.Tamiz (ed.), Multi-Objective Programming and Goal Programming: Theories and Applications, LN in Econ. Math. Systems 432, Springer-Verlag, Berlin, 148-163, (1996) preprint, University of Erlangen-Nürnberg, (2000)
- [14] Kelly, J. L., *General Topology*, Springer-Verlag, New York, (1955)

- [15] Kuroiwa, D., *The Natural Criteria in Set-Valued Optimization*, RIMS Kokyuroku 1031, 85-90, (1998)
- [16] Kuroiwa, D., *On Natural Criteria in Set-Valued Optimization*, RIMS Kokyuroku 1048, 86-92, (1998)
- [17] Kuroiwa, Tanaka, T., D., Truong, X. D. H., *On Cone Convexity of Set-Valued Maps*, *Nonlinear Analysis* 30, No. 3, 1487-1496, (1997)
- [18] Li, S. J., Yang, X. Q., Chen, G., *Vector Ekeland Variational Principle*, in Giannessi, F. (ed.), *Vector Variational Inequalities and Vector Equilibria*, Kluwer Academic Publishers, Dordrecht, (2000)
- [19] Löhne, A., *Minimalpunkttheoreme in uniformen Räumen und verwandte Aussagen*, Diplomarbeit, University of Halle-Wittenberg, Halle, (2001)
- [20] Luc, D. T., *Theory of Vector Optimization*, LN in Econ. Math. Systems 319, Springer-Verlag, Berlin, 1989
- [21] Nemeth, A. B., *A Nonconvex Vector Minimization Problem*, *Nonlinear Analysis* 10, 669-678, (1986)
- [22] Phelps, R. R., *Convex Functions, Monotone Operators and Differentiability*, LN in Mathematics 1364, Springer-Verlag, Berlin, (1989)
- [23] Tammer, Chr., *A Generalization of Ekeland's Variational Principle*, *Optimization*, 25, 129-141, (1992)
- [24] Truong, X. D. H., *Ekeland's Principle for a Set-Valued Map Studied with the Set Optimization Approach*, manuscript, (2002)
- [25] Truong, X. D. H., *Optimal Solution for Set-Valued Optimization Problems*, manuscript, (2002)
- [26] Weidner, P., *Ein Trennungskonzept und seine Anwendung auf Vektoroptimierungsverfahren*, Habilitation Thesis, Halle, 1987
- [27] Young, R.C., *The Algebra of Many-Valued Quantities*, *Math. Ann.* 104, 260-290, (1931).